

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
15 November 2001 (15.11.2001)

PCT

(10) International Publication Number  
**WO 01/85942 A2**

(51) International Patent Classification<sup>7</sup>: **C12N 15/12**,  
15/10, C12Q 1/68, C07K 14/47, 14/705, 16/18, 16/28,  
A01K 67/027, A61K 38/17, 39/395, G01N 33/53, 33/577

(21) International Application Number: PCT/US01/14355

(22) International Filing Date: 3 May 2001 (03.05.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/201,960 5 May 2000 (05.05.2000) US  
60/202,729 8 May 2000 (08.05.2000) US  
60/209,705 5 June 2000 (05.06.2000) US  
60/210,149 7 June 2000 (07.06.2000) US  
60/213,215 21 June 2000 (21.06.2000) US

(71) Applicant (*for all designated States except US*): **INCYTE GENOMICS, INC.** [US/US]; 3160 Porter Drive, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **YUE, Henry** [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). **TANG, Y., Tom** [US/US]; 4230 Ranwick Court, San Jose, CA 94304 (US). **AU-YOUNG, Janice** [US/US]; 233 Golden Eagle Lane, Brisbane, CA 94005 (US). **LU, Dyung, Aina, M.** [US/US]; 233 Coy Drive, San Jose, CA 95123 (US). **BAUGHN, Mariah, R.** [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). **HILLMAN, Jennifer, L.** [US/US]; 230 Monroe Drive #17, Mountain View, CA 94040 (US). **AZIMZAI, Yalda** [US/US]; 5518 Boulder Canyon Drive, Castro Valley, CA 94552 (US).

**LAL, Preeti** [IN/US]; P.O. Box 5142, Santa Clara, CA 95056 (US). **YAO, Monique, G.** [US/US]; 111 Frederick Court, Mountain View, CA 94043 (US). **BANDMAN, Olga** [US/US]; 366 Anna Avenue, Mountain View, CA 94043 (US). **BURFORD, Neil** [US/US]; 105 Wildwood Circle, Durham, CT 06422 (US). **BATRA, Sajeew** [US/US]; 555 El Camino Real #709, Sunnyvale, CA 94087 (US). **KEARNEY, Liam** [IE/US]; 50 Woodside Avenue, San Francisco, CA 94127 (US). **POLICKY, Jennifer, L.** [US/US]; 1511 Jarvis Court, San Jose, CA 95118 (US).

(74) Agents: **HAMLET-COX, Diana** et al.; Incyte Genomics, Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— *without international search report and to be republished upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: CYTOSKELETON-ASSOCIATED PROTEINS

(57) Abstract: The invention provides human cytoskeleton-associated proteins (CYSKP) and polynucleotides which identify and encode CYSKP. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with aberrant expression of CYSKP.



WO 01/85942 A2

## CYTOSKELETON-ASSOCIATED PROTEINS

### TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of cytoskeleton-associated proteins and to the use of these sequences in the diagnosis, treatment, and prevention of cell proliferative, autoimmune/inflammatory, vesicle trafficking, neurological, cell motility, reproductive, and muscle disorders, and in the assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid sequences of cytoskeleton-associated proteins.

### BACKGROUND OF THE INVENTION

The cytoskeleton, a cytoplasmic system of protein fibers, mediates cell shape, structure, and movement. The cytoskeleton supports the cell membrane and forms tracks along which organelles and other elements move in the cytosol. The cytoskeleton is a dynamic structure that allows cells to adopt various shapes and to carry out directed movements. Additionally, molecules can be sequestered to a specific cellular location through interaction with cytoskeleton associated proteins. Major cytoskeletal fibers are the microfilaments, the microtubules, and the intermediate filaments. Motor proteins, including myosin, dynein, and kinesin, drive movement of, or along, the fibers. Accessory or associated proteins modify the structure or activity of the fibers while cytoskeletal membrane anchors connect the fibers to the cell membrane. Other proteins associated with the cytoskeleton have roles in processes such as secretion and intracellular signaling. (The cytoskeleton is reviewed in Lodish, H. et al. (1995) Molecular Cell Biology Scientific American Books, New York NY.)

#### Microtubules and Associated Proteins

##### Tubulins

Microtubules, cytoskeletal fibers with a diameter of 24 nm, have multiple roles in the cell. Bundles of microtubules form cilia and flagella, which are whip-like extensions of the cell membrane that are necessary for sweeping materials across an epithelium and for swimming of sperm, respectively. Marginal bands of microtubules in red blood cells and platelets are important for these cells' pliability. Organelles, membrane vesicles, and proteins are transported in the cell along tracks of microtubules. For example, microtubules run through nerve cell axons, allowing bi-directional transport of materials and membrane vesicles between the cell body and the nerve terminal. Failure to supply the nerve terminal with these vesicles blocks the transmission of neural signals. Microtubules, in the form of the spindle, are also critical to chromosomal movement during cell division. Both stable and short-lived populations of microtubules exist in the cell.

Microtubules are a polymer of GTP-binding tubulin protein subunits. Each subunit is a

heterodimer of  $\alpha$ - and  $\beta$ - tubulin, multiple isoforms of which exist. Alpha-tubulin undergoes a number of post-translational modifications, including acetylation, polyglutamylation, truncation of two amino acids (forming  $\Delta 2$  tubulin), and tyrosination. In some cases, these modifications can affect microtubule stability. The hydrolysis of GTP is linked to the addition of tubulin subunits at the end of a  
5 microtubule. The subunits interact head to tail to form protofilaments; the protofilaments interact side to side to form a microtubule. A microtubule is polarized, one end ringed with  $\alpha$ -tubulin and the other with  $\beta$ -tubulin, and the two ends differ in their rates of assembly. Each microtubule is generally composed of 13 protofilaments although 11 or 15 protofilament-microtubules are sometimes found. Cilia and flagella contain doublet microtubules. Microtubules grow from specialized structures known  
10 as centrosomes or microtubule-organizing centers (MTOCs). MTOCs may contain one or two centrioles, which are pinwheel arrays of triplet microtubules. The basal body, the organizing center located at the base of a cilium or flagellum, contains one centriole.  $\gamma$ - tubulin present in the MTOC is important for nucleating the polymerization of  $\alpha$ - and  $\beta$ - tubulin heterodimers but does not polymerize into microtubules. The protein pericentrin is found in the MTOC and has a role in microtubule  
15 assembly.

#### Microtubule-Associated Proteins

Microtubule-associated proteins (MAPs) have roles in the assembly and stabilization of microtubules. One major family of MAPs, assembly MAPs, can be identified in neurons as well as non-neuronal cells. Assembly MAPs are responsible for cross-linking microtubules in the cytosol.  
20 These MAPs are organized into two domains: a basic microtubule-binding domain and an acidic projection domain. The projection domain is the binding site for membranes, intermediate filaments, or other microtubules. Based on sequence analysis, assembly MAPs can be further grouped into two types: Type I and Type II.

Type I MAPs, which include MAP1A and MAP1B, are large, filamentous molecules that co-  
25 purify with microtubules and are abundantly expressed in brain and testis. They contain several repeats of a positively-charged amino acid sequence motif that binds and neutralizes negatively charged tubulin, leading to stabilization of microtubules. MAP1A and MAP1B are each derived from a single precursor polypeptide that is subsequently proteolytically processed to generate one heavy chain and one light chain.

30 Another light chain, LC3, is a 16.4 kDa molecule that binds MAP1A, MAP1B, and microtubules. It is suggested that LC3 is synthesized from a source other than the MAP1A or MAP1B transcripts, and the expression of LC3 may be important in regulating the microtubule binding activity of MAP1A and MAP1B during cell proliferation (Mann, S. S. et al. (1994) J. Biol. Chem. 269:11492-11497).

Type II MAPs, which include MAP2a, MAP2b, MAP2c, MAP4, and Tau, are characterized by three to four copies of an 18-residue sequence in the microtubule-binding domain. MAP2a, MAP2b, and MAP2c are found only in dendrites, MAP4 is found in non-neuronal cells, and Tau is found in axons and dendrites of nerve cells. Alternative splicing of the Tau mRNA leads to the existence of multiple forms of Tau protein. Tau phosphorylation is altered in neurodegenerative disorders such as Alzheimer's disease, Pick's disease, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia and Parkinsonism linked to chromosome 17. The altered Tau phosphorylation leads to a collapse of the microtubule network and the formation of intraneuronal Tau aggregates (Spillantini, M.G. and Goedert, M. (1998) Trends Neurosci. 21:428-433).

Microtubule stability may also be regulated by severing the microtubule along its length. The protein katanin, a member of the AAA adenosine triphosphatase (ATPase) superfamily, uses ATP hydrolysis energy to sever and disassemble stable microtubules. Katanin may play a role in releasing microtubules from centrosomes, regulating assembly of the mitotic spindle, and accelerating microtubule turnover during cell cycle transitions (Hartman, J.J. and Vale, R.D. (1999) Science 286:782-785).

Microtubular aggregates are associated with several disorders. An extraskeletal myxoid chondrosarcoma tumor from human contained parallel arrays of microtubules within the rough endoplasmic reticulum (Suzuki, T. et al. (1988) J. Pathol. 156:51-57). Microtubular aggregates were also found in hepatocytes from chimpanzees infected with hepatitis C virus. Monoclonal antibodies prepared to these aggregates detect a protein called p44 (or microtubular aggregates protein) (Maeda, T. et al. (1989) J. Gen. Virol. 70:1401-1407). A human homolog of p44 is inducible by interferon- $\alpha$  and interferon- $\beta$ , but not by interferon- $\gamma$ . p44 protein may be a mediator in the antiviral action of interferon (Kitamura, A. et al. (1994) Eur. J. Biochem. 224:877-883).

#### Dynein-related Motor Proteins

Dyneins are (-) end-directed motor proteins which act on microtubules. Two classes of dyneins exist, cytosolic and axonemal. Cytosolic dyneins are responsible for translocation of materials along cytoplasmic microtubules; for example, transport from the nerve terminal to the cell body and transport of endocytic vesicles to lysosomes. Cytoplasmic dyneins are also reported to play a role in mitosis. Axonemal dyneins are responsible for the beating of flagella and cilia. Dynein on one microtubule doublet walks along the adjacent microtubule doublet. This sliding force produces bending forces that cause the flagellum or cilium to beat. Dyneins have a native mass between 1000 and 2000 kDa and contain either two or three force-producing heads driven by the hydrolysis of ATP. The heads are linked via stalks to a basal domain which is composed of a highly variable number of accessory intermediate and light chains.



### Kinesin-related Motor Proteins

Kinesins are (+) end-directed motor proteins which act on microtubules. The prototypical kinesin molecule is involved in the transport of membrane-bound vesicles and organelles. This function is particularly important for axonal transport in neurons. Kinesin is also important in all cell types for the transport of vesicles from the Golgi complex to the endoplasmic reticulum. This role is critical for maintaining the identity and functionality of these secretory organelles.

Kinesins define a ubiquitous, conserved family of over 50 proteins that can be classified into at least 8 subfamilies based on primary amino acid sequence, domain structure, velocity of movement, and cellular function. (Reviewed in Moore, J.D. and S.A. Endow (1996) *Bioessays* 18:207-219; and Hoyt, A.M. (1994) *Curr. Opin. Cell Biol.* 6:63-68.) The prototypical kinesin molecule is a heterotetramer comprised of two heavy polypeptide chains (KHCs) and two light polypeptide chains (KLCs). The KHC subunits are typically referred to as "kinesin." KHC is about 1000 amino acids in length, and KLC is about 550 amino acids in length. Two KHCs dimerize to form a rod-shaped molecule with three distinct regions of secondary structure. At one end of the molecule is a globular motor domain that functions in ATP hydrolysis and microtubule binding. Kinesin motor domains are highly conserved and share over 70% identity. Beyond the motor domain is an  $\alpha$ -helical coiled-coil region which mediates dimerization. At the other end of the molecule is a fan-shaped tail that associates with molecular cargo. The tail is formed by the interaction of the KHC C-termini with the two KLCs.

Members of the more divergent subfamilies of kinesins are called kinesin-related proteins (KRPs), many of which function during mitosis in eukaryotes (Hoyt, *supra*). Some KRPs are required for assembly of the mitotic spindle. *In vivo* and *in vitro* analyses suggest that these KRPs exert force on microtubules that comprise the mitotic spindle, resulting in the separation of spindle poles. Phosphorylation of KRP is required for this activity. Failure to assemble the mitotic spindle results in abortive mitosis and chromosomal aneuploidy, the latter condition being characteristic of cancer cells. In addition, a unique KRP, centromere protein E, localizes to the kinetochore of human mitotic chromosomes and may play a role in their segregation to opposite spindle poles.

### Microfilaments and Associated Proteins

#### Actins

Microfilaments, cytoskeletal filaments with a diameter of 7-9 nm, are vital to cell locomotion, cell shape, cell adhesion, cell division, and muscle contraction. Assembly and disassembly of the microfilaments allow cells to change their morphology. Microfilaments are the polymerized form of actin, the most abundant intracellular protein in the eukaryotic cell. Human cells contain six isoforms of actin. The three  $\alpha$ -actins are found in different kinds of muscle, nonmuscle  $\beta$ -actin and nonmuscle  $\gamma$ -actin are found in nonmuscle cells, and another  $\gamma$ -actin is found in intestinal smooth muscle cells. G-

actin, the monomeric form of actin, polymerizes into polarized, helical F-actin filaments, accompanied by the hydrolysis of ATP to ADP. Actin filaments associate to form bundles and networks, providing a framework to support the plasma membrane and determine cell shape. These bundles and networks are connected to the cell membrane. In muscle cells, thin filaments containing actin slide past thick  
5 filaments containing the motor protein myosin during contraction. Other actin-related filaments are not part of the actin cytoskeleton, but rather associate with microtubules and dyenin.

#### Actin-Associated Proteins

Actin-associated proteins have roles in cross-linking, severing, and stabilization of actin filaments and in sequestering actin monomers. Several of the actin-associated proteins have multiple  
10 functions. Bundles and networks of actin filaments are held together by actin cross-linking proteins. These proteins have two actin-binding sites, one for each filament. Short cross-linking proteins promote bundle formation while longer, more flexible cross-linking proteins promote network formation. Calmodulin-like calcium-binding domains in actin cross-linking proteins allow calcium regulation of cross-linking. Group I cross-linking proteins have unique actin-binding domains and include the 30 Kd  
15 protein, EF-1a, fascin, and scruin. Group II cross-linking proteins have a 7,000-MW actin-binding domain and include villin and dematin. Group III cross-linking proteins have pairs of a 26,000-MW actin-binding domain and include alpha-actinin, fimbrin, spectrin, dystrophin, ABP 120, and filamin.

Severing proteins regulate the length of actin filaments by breaking them into short pieces or by blocking their ends. Severing proteins include gCAP39, severin (fragmin), gelsolin, and villin.

20 Capping proteins can cap the ends of actin filaments, but cannot break filaments. Capping proteins include CapZ, tropomodulin, and tensin.

Tensin, which is found in focal adhesions, also crosslinks actin filaments. Integrin activation by the extracellular matrix leads to the phosphorylation of tensin on tyrosine, serine, and threonine residues; this phosphorylation also occurs in cells transformed with oncogenes. Tensin has an SH2  
25 domain and may bind to other tyrosine-phosphorylated proteins. (Lo, S.H. et al. (1997) J. Cell Biol. 136:1349-1361.) The N-terminus of tensin contains a region homologous to the catalytic domain of a putative tyrosine phosphatase (PTP) from *Saccharomyces cerevisiae*. This PTP domain in tensin may mediate binding interactions with phosphorylated polypeptides (Haynie, D.T. and Ponting, C.P. (1996) Protein Sci. 5:2643-2646). Mice which lack the tensin gene have kidney abnormalities, indicating that  
30 the loss of tensin leads to weakening of focal adhesions in the kidney (Lo, supra).

The proteins thymosin and profilin sequester actin monomers in the cytosol, allowing a pool of unpolymerized actin to exist. Profilin may also stimulate F-actin formation by effectively lowering the critical concentration required for actin monomer addition (Gertler, F.B. et al. (1996) Cell 87:227-239).

The actin-associated proteins tropomyosin, troponin, and caldesmon regulate muscle

contraction in response to calcium. The tropomyosin proteins, found in muscle and nonmuscle cells, are  $\alpha$ -helical and form coiled-coil dimers. Striated muscle tropomyosin mediates the interactions between the troponin complex and actin, regulating muscle contraction (PROSITE PDOC00290 Tropomyosins signature). The troponin complex is composed of troponin-T, troponin-I, and troponin-C. Troponin-T binds tropomyosin, linking troponin-I and troponin-C to tropomyosin.

The actin-associated proteins tropomyosin, troponin, and caldesmon regulate muscle contraction in response to calcium. The tropomyosin proteins, found in muscle and nonmuscle cells, are  $\alpha$ -helical and form coiled-coil dimers. Striated muscle tropomyosin mediates the interactions between the troponin complex and actin, regulating muscle contraction. (PROSITE PDOC00290 Tropomyosins signature.) The troponin complex is composed of troponin-T, troponin-I, and troponin-C. Troponin-T binds tropomyosin, linking troponin-I and troponin-C to tropomyosin.

Many proteins involved in the regulation of actin assembly have characteristic protein-protein interaction domains, such as for example the calponin homology (CH) domain found in actin cross-linking proteins including alpha-actinin, spectrin, and fimbrin. Other proteins which are localized primarily in focal adhesions, macromolecular complexes which mediate the contact between extracellular matrix receptors and the cytoskeleton, contain protein-protein interaction motifs known as LIM domains. For example, zyxin is a protein that plays a role in the spatial control of actin assembly and contains three tandem LIM domains. Zyxin also interacts with alpha-actinin through its proline rich N-terminus (Beckerle, M. C. (1997) BioEssays 19:949-957).

Cytoskeletal proteins are implicated in several diseases. Pathologies such as muscular dystrophy, nephrotic syndrome, and dilated cardiomyopathy have been associated with differential expression of alpha-actinin-3 (Vainzof, M. et al. (1997) Neuropediatrics 28:223-228; Smoyer, W.E. and Mundel, P. (1998) J. Mol. Med. 76:172-183; and Sussman, M.A. et al. (1998) J. Clin. Invest. 101:51-61). Alpha-actinin and several MAPs are present in Hirano bodies, which are observed more frequently in the elderly and in patients with neurodegenerative diseases such as Alzheimer's disease (Maciver, S.K. and Harrington, C.R. (1995) Neuroreport. 6:1985-1988). Actinin-4, a novel actin-bundling protein, appears to be associated with the cell motility of metastatic cancer cells. Other disease associations include premature chromosome condensation which is frequently observed in dividing cells from tumor tissue (Murnane, J.P. (1995) Cancer Metastasis Rev. 14:17-29) and the significant roles of axonemal and assembly MAPs in viral pathogenesis (Sodeik, B. et al. (1997) J. Cell Biol. 136:1007-1021).

#### Intermediate Filaments and Associated Proteins

Intermediate filaments (IFs) are cytoskeletal fibers with a diameter of 10 nm, intermediate between that of microfilaments and microtubules. They serve structural roles in the cell, reinforcing cells and organizing cells into tissues. IFs are particularly abundant in epidermal cells and in neurons.

IFs are extremely stable, and, in contrast to microfilaments and microtubules, do not function in cell motility. IF proteins include acidic keratins, basic keratins, desmin, glial fibrillary acidic protein, vimentin, peripherin, neurofilaments, nestin, and lamins.

IFs have a central  $\alpha$ -helical rod region interrupted by short nonhelical linker segments. The rod region is bracketed, in most cases, by non-helical head and tail domains. The rod regions of intermediate filament proteins associate to form a coiled-coil dimer. A highly ordered assembly process leads from the dimers to the IFs. Neither ATP nor GTP is needed for IF assembly, unlike that of microfilaments and microtubules.

IF-associated proteins (IFAPs) mediate the interactions of IFs with one another and with other cell structures. IFAPs cross-link IFs into a bundle, into a network, or to the plasma membrane, and may cross-link IFs to the microfilament and microtubule cytoskeleton. Microtubules and IFs in particular are closely associated. IFAPs include BPAG1, plakoglobin, desmoplakin I, desmoplakin II, plectin, ankyrin, filaggrin, and lamin B receptor.

The N-terminal portion of ankyrin consists of a repeated 33-amino acid motif, the ankyrin repeat, which is involved in specific protein-protein interactions. Variable regions within the motif are responsible for specific protein binding, such that different ankyrin repeats are involved in binding to tubulin, anion exchange protein, voltage-gated sodium channel,  $\text{Na}^+/\text{K}^+$ -ATPase, and neurofascin. The ankyrin motif is also found in transcription factors, such as NF- $\kappa$ -B, and in the yeast cell cycle proteins CDC10, SW14, and SW16. Proteins involved in tissue differentiation, such as Drosophila Notch and C. elegans LIN-12 and GLP-1, also contain ankyrin-like repeats. Lux et al. (1990; Nature 344:36-42) suggest that ankyrin-like repeats function as 'built-in' ankyrins and form binding sites for integral membrane proteins, tubulin, and other proteins.

#### Cytoskeletal-Membrane Anchors

Cytoskeletal fibers are attached to the plasma membrane by specific proteins. These attachments are important for maintaining cell shape and for muscle contraction. In erythrocytes, the spectrin-actin cytoskeleton is attached to cell membrane by three proteins, band 4.1, ankyrin, and adducin. Defects in this attachment result in abnormally shaped cells which are more rapidly degraded by the spleen, leading to anemia. In platelets, the spectrin-actin cytoskeleton is also linked to the membrane by ankyrin; a second actin network is anchored to the membrane by filamin. In muscle cells the protein dystrophin links actin filaments to the plasma membrane; mutations in the dystrophin gene lead to Duchenne muscular dystrophy. In adherens junctions and adhesion plaques the peripheral membrane proteins  $\alpha$ -actinin and vinculin attach actin filaments to the cell membrane.

IFs are also attached to membranes by cytoskeletal-membrane anchors. The nuclear lamina is attached to the inner surface of the nuclear membrane by the lamin B receptor. Vimentin IFs are

attached to the plasma membrane by ankyrin and plectin. Desmosome and hemidesmosome membrane junctions hold together epithelial cells of organs and skin. These membrane junctions allow shear forces to be distributed across the entire epithelial cell layer, thus providing strength and rigidity to the epithelium. IFs in epithelial cells are attached to the desmosome by plakoglobin and desmoplakins.

- 5 The proteins that link IFs to hemidesmosomes are not known. Desmin IFs surround the sarcomere in muscle and are linked to the plasma membrane by paranemin, synemin, and ankyrin.

#### Proteins of the Erythrocyte Membrane Skeleton

- Distribution of oxygen throughout the vertebrate body is effected by red blood cells (erythrocytes). Oxygen diffuses from surrounding water or from the atmosphere through either gill  
10 epithelium or pulmonary epithelial type I cells. Oxygen then diffuses through the blood capillary endothelium directly to the blood circulatory system and through the erythrocyte membrane and is stored as soluble oxyhemoglobin in the cytoplasm. Oxygen is released from hemoglobin at sites throughout the organism and diffuses out from the erythrocyte to other target cells. The structure of the erythrocyte membrane as well as that of other non-erythrocyte cells must be maintained to enable  
15 efficient diffusion of oxygen to intracellular compartments.

- The erythrocyte membrane is comprised of i) a cholesterol-rich phospholipid bilayer in which many trans-bilayer proteins are embedded, ii) external glycosylphosphatidylinositol-anchored proteins (GPI-proteins), and iii) the erythrocyte or membrane skeleton that laminates the inner surface of the bilayer. The trans-bilayer proteins include anion exchangers, glycophorins, glucose transporters, and a  
20 variety of cation transporters and pumps. The erythrocyte GPI-proteins include acetylcholinesterase and decay-accelerating factor (CD 55). The skeletal proteins are organized on the cortical, or cytoplasmic, face of the plasma membrane. These proteins include protein 4.1, protein p55,  $\alpha$ - and  $\beta$ -spectrin, actin, and actin-binding proteins such as dematin, tropomyosin, and tropomodulin.  $\alpha$ - and  $\beta$ -spectrin combine to form a heterotetramer in vivo. The spectrin heterotetramer organizes into a cortical  
25 bidimensional network with a hexagonal mesh. The network is linked to trans-bilayer proteins through a protein complex comprising  $\beta$ -spectrin, ankyrin, anion exchanger, and protein 4.2 and through the “triangular” interaction between protein 4.1, glycophorin C, and protein p55. Structural and functional variants of erythrocyte membrane proteins have been found in a variety of tissues. Variants may be transcribed from multigene families, e.g., anion exchanger, ankyrin, or spectrin, or from single  
30 gene families, e.g., protein 4.1 or protein 4.2. mRNA transcripts undergo tissue-specific alternative splicing. Many congenital hemolytic anemias result from mutations in the above-mentioned genes encoding erythrocyte membrane proteins. For example, hereditary elliptocytosis stems from an array of mutations in the spectrin genes at or near the head-to-head self-association region of the spectrin tetramer, or from mutations in the protein 4.1 gene which reduce levels of protein 4.1. In another

example, hereditary spherocytosis is associated with mutations in the ankyrin gene, the anion exchanger gene, the protein 4.2 gene, or the  $\alpha$ - and  $\beta$ -spectrin genes. (Delaunay J. (1995) Transfus. Clin. Biol. 2:207-216.)

Protein 4.1 is an 80 kDa erythrocyte membrane protein with four functional domains. These domains include: i) a 30 kDa basic N-terminal domain, homologous to the ERM (Ezrin/Radixin/Moesin) family of actin- and transmembrane protein-binding proteins (Tsukita, S. et al. (1997) Trends Biochem. Sci. 22:53-58); ii) a 16 kDa hydrophilic domain containing a protein kinase C phosphorylation site; iii) a 10 kDa highly charged domain containing a cAMP-dependent protein kinase phosphorylation site critical for the interaction with spectrin and actin; and iv) a 22/24 kDa acidic domain. Protein 4.1 is a member of a structurally and functionally related protein 4.1 family. The protein 4.1 family is part of an evolutionarily related protein superfamily that includes many tyrosine phosphatases. (Baklouti, F. et al. (1997) Genomics 39:289-302.)

In contrast to the strictly cortical localization of protein 4.1 in mature enucleate erythrocytes, protein 4.1 epitopes have been observed throughout the cytoplasmic compartment and the nucleoskeleton in nucleated cells. In particular, protein 4.1 is present in the nucleoskeleton during interphase, in the mitotic spindle during mitosis, in perichromatin during telophase, and in the midbody during cytokinesis. (Krauss, S.W. et al. (1997) J. Cell Biol. 137:275-289.)

Differential expression of the protein 4.1 gene resulting in a number of mRNA splice variants has been observed in various human and rodent tissues. Comparison of the gene structure and mRNA splice variants revealed the extreme genomic sequence conservation of protein 4.1 between different species. The 5' UTR of both the human and rodent mRNA species has not been successfully identified and sequenced, possibly due to GC-rich regions therein which give rise to technical complications during nucleotide sequencing reactions. (Baklouti, supra; Conboy, J.G. (1988) Proc. Natl. Acad. Sci. 85:9062-9065.)

Analysis of proteins included in the ERM family of proteins has indicated that the N-terminal domain interacts with intracellular domains of transmembrane proteins such as CD44 and the C-terminal domain binds actin. Both interactions involve interactions with Rho-GTP protein complex, polyphosphoinositides, and serine/threonine kinase and tyrosine kinase activities. Many of the phosphorylation sites on ERM proteins are conserved. Although expression of ERM proteins in vivo is restricted to tissues such as endothelium, repression of ERM protein gene expression is released under conditions of cell culture. (Tsukita, supra.)

The cortical actin cytoskeleton participates in various membrane-based processes which necessitate a large amount of functional plasticity in the molecular components involved. A family of proteins homologous to band 4.1 is involved in the reorganization of the actin cytoskeleton in response

to various stimuli and probably plays a role in transmembrane signaling. This family includes tyrosine phosphatases, substrates of tyrosine kinases and a candidate for a tumor-suppressor gene. (Arpin M, et al. (1994) Curr. Opin. Cell Biol. 6:136-141.)

Disruptions in cytoskeletal protein interaction have been identified in a number of disease  
 5 conditions or disorders. Neurofibromatosis type 2 is an autosomal dominant disease of the nervous system. Schwann cells isolated from patients with neurofibromatosis type 2 have characteristic morphology and growth parameters which differ from control Schwann cells. A gene associated with neurofibromatosis type 2 has been identified and is termed NF2. The NF2 gene product, known as schwannomin or merlin, is a member of the protein 4.1 superfamily, and mutations in the NF2 gene  
 10 have been shown to be associated with the disease. (Rosenbaum, C. et al. (1998) Neurobiol. Dis. 5:55-64.) In addition, a form of psoriasis may be due to altered expression or distribution in epidermal epithelium of analogs of erythrocyte protein 4.1. (Shimizu, T. (1996) Histol. Histopathol. 11:495-501.) Erythrocytes carrying mutations in spectrin and protein 4.1 showed differing sensitivities to invasion by Plasmodium falciparum. (Facer, C.A. (1995) Parasitol Res. 81:52-57.) Furthermore, antibodies raised  
 15 against erythrocyte protein 4.1 stained the majority of neurofibrillary tangles in the prefrontal cortex and hippocampus of brain tissue from patients with Alzheimer's disease. A 68 kDa protein was identified as the most likely brain analog of erythrocyte protein 4.1. (Sihag, R.K. et al. (1994) Brain Res. 656:14-26.)

The discovery of new cytoskeleton-associated proteins and the polynucleotides encoding them  
 20 satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of cell proliferative, autoimmune/inflammatory, vesicle trafficking, neurological, cell motility, reproductive, and muscle disorders, and in the assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid sequences of cytoskeleton-associated proteins.

25

## SUMMARY OF THE INVENTION

The invention features purified polypeptides, cytoskeleton-associated proteins, referred to collectively as "CYSKP" and individually as "CYSKP-1," "CYSKP-2," "CYSKP-3," "CYSKP-4," "CYSKP-5," "CYSKP-6," "CYSKP-7," "CYSKP-8," "CYSKP-9," "CYSKP-10," "CYSKP-11,"  
 30 "CYSKP-12," "CYSKP-13," "CYSKP-14," "CYSKP-15," "CYSKP-16," "CYSKP-17," "CYSKP-18," "CYSKP-19," "CYSKP-20," "CYSKP-21," "CYSKP-22," "CYSKP-23," "CYSKP-24," "CYSKP-25," "CYSKP-26," "CYSKP-27," "CYSKP-28," "CYSKP-29," "CYSKP-30," "CYSKP-31," "CYSKP-32," "CYSKP-33," and "CYSKP-34." In one aspect, the invention provides an isolated polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence

selected from the group consisting of SEQ ID NO:1-34, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, and d) an immunogenic  
5 fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34. In one alternative, the invention provides an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:1-34.

The invention further provides an isolated polynucleotide encoding a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group  
10 consisting of SEQ ID NO:1-34, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34. In one  
15 alternative, the polynucleotide encodes a polypeptide selected from the group consisting of SEQ ID NO:1-34. In another alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:35-68.

Additionally, the invention provides a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding a polypeptide selected from the group consisting  
20 of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, and d) an immunogenic fragment of a polypeptide having an amino  
25 acid sequence selected from the group consisting of SEQ ID NO:1-34. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

The invention also provides a method for producing a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of  
30 SEQ ID NO:1-34, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34. The method comprises a)



culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

Additionally, the invention provides an isolated antibody which specifically binds to a  
5 polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, and d) an immunogenic  
10 fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34.

The invention further provides an isolated polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:35-68, b) a naturally occurring polynucleotide comprising a polynucleotide sequence at least 90%  
15 identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:35-68, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

Additionally, the invention provides a method for detecting a target polynucleotide in a sample,  
20 said target polynucleotide having a sequence of a polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:35-68, b) a naturally occurring polynucleotide comprising a polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:35-68, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the  
25 polynucleotide of b), and e) an RNA equivalent of a)-d). The method comprises a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and b) detecting the presence or absence of  
30 said hybridization complex, and optionally, if present, the amount thereof. In one alternative, the probe comprises at least 60 contiguous nucleotides.

The invention further provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID

NO:35-68, b) a naturally occurring polynucleotide comprising a polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:35-68, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). The method comprises a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

The invention further provides a composition comprising an effective amount of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, and a pharmaceutically acceptable excipient. In one embodiment, the composition comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-34. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional CYSKP, comprising administering to a patient in need of such treatment the composition.

The invention also provides a method for screening a compound for effectiveness as an agonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. In one alternative, the invention provides a composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with decreased expression of functional CYSKP, comprising administering to a patient in need of such treatment the composition.

Additionally, the invention provides a method for screening a compound for effectiveness as an antagonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, b) a naturally occurring

polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting antagonist activity in the sample. In one alternative, the invention provides a composition comprising an antagonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with overexpression of functional CYSKP, comprising administering to a patient in need of such treatment the composition.

The invention further provides a method of screening for a compound that specifically binds to a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34. The method comprises a) combining the polypeptide with at least one test compound under suitable conditions, and b) detecting binding of the polypeptide to the test compound, thereby identifying a compound that specifically binds to the polypeptide.

The invention further provides a method of screening for a compound that modulates the activity of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34. The method comprises a) combining the polypeptide with at least one test compound under conditions permissive for the activity of the polypeptide, b) assessing the activity of the polypeptide in the presence of the test compound, and c) comparing the activity of the polypeptide in the presence of the test compound with the activity of the polypeptide in the absence of the test compound, wherein a change in the activity of the polypeptide in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide.

The invention further provides a method for screening a compound for effectiveness in

altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence selected from the group consisting of SEQ ID NO:35-68, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, and b) detecting altered expression of the target polynucleotide.

5           The invention further provides a method for assessing toxicity of a test compound, said method comprising a) treating a biological sample containing nucleic acids with the test compound; b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide selected from the group consisting of i) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:35-68, ii) a  
10 naturally occurring polynucleotide comprising a polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:35-68, iii) a polynucleotide having a sequence complementary to i), iv) a polynucleotide complementary to the polynucleotide of ii), and v) an RNA equivalent of i)-iv). Hybridization occurs under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in  
15 the biological sample, said target polynucleotide selected from the group consisting of i) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:35-68, ii) a naturally occurring polynucleotide comprising a polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:35-68, iii) a polynucleotide complementary to the polynucleotide of i), iv) a polynucleotide complementary to the  
20 polynucleotide of ii), and v) an RNA equivalent of i)-iv). Alternatively, the target polynucleotide comprises a fragment of a polynucleotide sequence selected from the group consisting of i)-v) above; c) quantifying the amount of hybridization complex; and d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated  
25 biological sample is indicative of toxicity of the test compound.

#### BRIEF DESCRIPTION OF THE TABLES

Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide sequences of the present invention.

30           Table 2 shows the GenBank identification number and annotation of the nearest GenBank homolog for polypeptides of the invention. The probability score for the match between each polypeptide and its GenBank homolog is also shown.

Table 3 shows structural features of polypeptide sequences of the invention, including predicted motifs and domains, along with the methods, algorithms, and searchable databases used for analysis of  
35 the polypeptides.

Table 4 lists the cDNA and genomic DNA fragments which were used to assemble polynucleotide sequences of the invention, along with selected fragments of the polynucleotide sequences.

Table 5 shows the representative cDNA library for polynucleotides of the invention.

5 Table 6 provides an appendix which describes the tissues and vectors used for construction of the cDNA libraries shown in Table 5.

Table 7 shows the tools, programs, and algorithms used to analyze the polynucleotides and polypeptides of the invention, along with applicable descriptions, references, and threshold parameters.

## 10 DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will  
15 be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so  
20 forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described.  
25 All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

## DEFINITIONS

30 "CYSKP" refers to the amino acid sequences of substantially purified CYSKP obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which intensifies or mimics the biological activity of CYSKP. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other

compound or composition which modulates the activity of CYSKP either by directly interacting with CYSKP or by acting on components of the biological pathway in which CYSKP participates.

An "allelic variant" is an alternative form of the gene encoding CYSKP. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

"Altered" nucleic acid sequences encoding CYSKP include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as CYSKP or a polypeptide with at least one functional characteristic of CYSKP. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding CYSKP, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding CYSKP. The encoded protein may also be "altered," and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent CYSKP. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of CYSKP is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms "amino acid" and "amino acid sequence" refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where "amino acid sequence" is recited to refer to a sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

"Amplification" relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity of

CYSKP. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of CYSKP either by directly interacting with CYSKP or by acting on components of the biological pathway in which CYSKP participates.

5           The term “antibody” refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')<sub>2</sub>, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind CYSKP polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or  
10           synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

          The term “antigenic determinant” refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to  
15           immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

          The term “antisense” refers to any composition capable of base-pairing with the “sense”  
20           (coding) strand of a specific nucleic acid sequence. Antisense compositions may include DNA; RNA; peptide nucleic acid (PNA); oligonucleotides having modified backbone linkages such as phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethoxy sugars; or oligonucleotides having modified bases such as 5-methyl cytosine, 2'-deoxyuracil, or 7-deaza-2'-deoxyguanosine. Antisense  
25           molecules may be produced by any method including chemical synthesis or transcription. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or translation. The designation “negative” or “minus” can refer to the antisense strand, and the designation “positive” or “plus” can refer to the sense strand of a reference DNA molecule.

30           The term “biologically active” refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, “immunologically active” or “immunogenic” refers to the capability of the natural, recombinant, or synthetic CYSKP, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

“Complementary” describes the relationship between two single-stranded nucleic acid sequences that anneal by base-pairing. For example, 5'-AGT-3' pairs with its complement, 3'-TCA-5'.

A “composition comprising a given polynucleotide sequence” and a “composition comprising a given amino acid sequence” refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding CYSKP or fragments of CYSKP may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

“Consensus sequence” refers to a nucleic acid sequence which has been subjected to repeated DNA sequence analysis to resolve uncalled bases, extended using the XL-PCR kit (Applied Biosystems, Foster City CA) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from one or more overlapping cDNA, EST, or genomic DNA fragments using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI) or Phrap (University of Washington, Seattle WA). Some sequences have been both extended and assembled to produce the consensus sequence.

“Conservative amino acid substitutions” are those substitutions that are predicted to least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

	Original Residue	Conservative Substitution
25	Ala	Gly, Ser
	Arg	His, Lys
	Asn	Asp, Gln, His
	Asp	Asn, Glu
	Cys	Ala, Ser
30	Gln	Asn, Glu, His
	Glu	Asp, Gln, His
	Gly	Ala
	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
35	Leu	Ile, Val
	Lys	Arg, Gln, Glu
	Met	Leu, Ile
	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr



Thr	Ser, Val
Trp	Phe, Tyr
Tyr	His, Phe, Trp
Val	Ile, Leu, Thr

---

5

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

10 A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to a chemically modified polynucleotide or polypeptide. Chemical modifications of a polynucleotide can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one  
 15 biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

A "detectable label" refers to a reporter molecule or enzyme that is capable of generating a measurable signal and is covalently or noncovalently joined to a polynucleotide or polypeptide.

20 "Differential expression" refers to increased or upregulated; or decreased, downregulated, or absent gene or protein expression, determined by comparing at least two different samples. Such comparisons may be carried out between, for example, a treated and an untreated sample, or a diseased and a normal sample.

A "fragment" is a unique portion of CYSKP or the polynucleotide encoding CYSKP which is  
 25 identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 16, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid  
 30 residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50%) of a polypeptide as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present  
 35 embodiments.

A fragment of SEQ ID NO:35-68 comprises a region of unique polynucleotide sequence that

specifically identifies SEQ ID NO:35-68, for example, as distinct from any other sequence in the genome from which the fragment was obtained. A fragment of SEQ ID NO:35-68 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:35-68 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:35-68 and the region of SEQ ID NO:35-68 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A fragment of SEQ ID NO:1-34 is encoded by a fragment of SEQ ID NO:35-68. A fragment of SEQ ID NO:1-34 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-34. For example, a fragment of SEQ ID NO:1-34 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-34. The precise length of a fragment of SEQ ID NO:1-34 and the region of SEQ ID NO:1-34 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A “full length” polynucleotide sequence is one containing at least a translation initiation codon (e.g., methionine) followed by an open reading frame and a translation termination codon. A “full length” polynucleotide sequence encodes a “full length” polypeptide sequence.

“Homology” refers to sequence similarity or, interchangeably, sequence identity, between two or more polynucleotide sequences or two or more polypeptide sequences.

The terms “percent identity” and “% identity,” as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and “diagonals saved”=4. The “weighted” residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the “percent similarity” between aligned polynucleotide sequences.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search

Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis programs including “blastn,” that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called “BLAST 2 Sequences” that is used for direct pairwise comparison of two nucleotide sequences. “BLAST 2 Sequences” can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The “BLAST 2 Sequences” tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the “BLAST 2 Sequences” tool Version 2.0.12 (April-21-2000) set at default parameters. Such default parameters may be, for example:

*Matrix: BLOSUM62*

*Reward for match: 1*

*Penalty for mismatch: -2*

*Open Gap: 5 and Extension Gap: 2 penalties*

*Gap x drop-off: 50*

*Expect: 10*

*Word Size: 11*

*Filter: on*

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

The phrases “percent identity” and “% identity,” as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some alignment methods take into account conservative amino acid substitutions. Such conservative substitutions,

explained in more detail above, generally preserve the charge and hydrophobicity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and “diagonals saved”=5. The PAM250 matrix is selected as the default residue weight table. As with polynucleotide alignments, the percent identity is reported by CLUSTAL V as the “percent similarity” between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the “BLAST 2 Sequences” tool Version 2.0.12 (April-21-2000) with blastp set at default parameters. Such default parameters may be, for example:

*Matrix: BLOSUM62*

*Open Gap: 11 and Extension Gap: 1 penalties*

*Gap x drop-off: 50*

*Expect: 10*

*Word Size: 3*

*Filter: on*

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

“Human artificial chromosomes” (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size and which contain all of the elements required for chromosome replication, segregation and maintenance.

The term “humanized antibody” refers to an antibody molecule in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

“Hybridization” refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of complementarity.

Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the “washing” step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml sheared, denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Such wash temperatures are typically selected to be about 5°C to 20°C lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength and pH. The  $T_m$  is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating  $T_m$  and conditions for nucleic acid hybridization are well known and can be found in Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2<sup>nd</sup> ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, sheared and denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term “hybridization complex” refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g.,  $C_0t$  or  $R_0t$  analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

An "immunogenic fragment" is a polypeptide or oligopeptide fragment of CYSKP which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment of CYSKP which is useful in any of the antibody production methods disclosed herein or known in the art.

The term "microarray" refers to an arrangement of a plurality of polynucleotides, polypeptides, or other chemical compounds on a substrate.

The terms "element" and "array element" refer to a polynucleotide, polypeptide, or other chemical compound having a unique and defined position on a microarray.

The term "modulate" refers to a change in the activity of CYSKP. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of CYSKP.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with a second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

"Post-translational modification" of an CYSKP may involve lipidation, glycosylation, phosphorylation, acetylation, racemization, proteolytic cleavage, and other modifications known in the art. These processes may occur synthetically or biochemically. Biochemical modifications will vary by

cell type depending on the enzymatic milieu of CYSKP.

"Probe" refers to nucleic acid sequences encoding CYSKP, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical  
5 labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes. "Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

10 Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the  
15 specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2<sup>nd</sup> ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel, F.M. et al. (1987) Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis, M. et al. (1990) PCR  
20 Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such  
25 purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South  
30 West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for

microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that  
5 hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary  
10 polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the  
15 artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, supra. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

20 Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

A "regulatory element" refers to a nucleic acid sequence usually derived from untranslated regions of a gene and includes enhancers, promoters, introns, and 5' and 3' untranslated regions (UTRs).  
25 Regulatory elements interact with host or viral proteins which control transcription, translation, or RNA stability.

"Reporter molecules" are chemical or biochemical moieties used for labeling a nucleic acid, amino acid, or antibody. Reporter molecules include radionuclides; enzymes; fluorescent, chemiluminescent, or chromogenic agents; substrates; cofactors; inhibitors; magnetic particles; and  
30 other moieties known in the art.

An "RNA equivalent," in reference to a DNA sequence, is composed of the same linear sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.



The term "sample" is used in its broadest sense. A sample suspected of containing CYSKP, nucleic acids encoding CYSKP, or fragments thereof may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

5       The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide comprising the epitope  
10   A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

      The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which  
15   they are naturally associated.

      A "substitution" refers to the replacement of one or more amino acid residues or nucleotides by different amino acid residues or nucleotides, respectively.

      "Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers,  
20   microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

      A "transcript image" refers to the collective pattern of gene expression by a particular cell type or tissue under given conditions at a given time.

      "Transformation" describes a process by which exogenous DNA is introduced into a recipient  
25   cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, bacteriophage or viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed cells"  
30   includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

      A "transgenic organism," as used herein, is any organism, including but not limited to animals and plants, in which one or more of the cells of the organism contains heterologous nucleic

acid introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or in vitro fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The transgenic organisms contemplated in accordance with the present invention include bacteria, cyanobacteria, fungi, plants and animals. The isolated DNA of the present invention can be introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention into such organisms are widely known and provided in references such as Sambrook et al. (1989), supra.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternative splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides will generally have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or greater sequence identity over a

certain defined length of one of the polypeptides.

## THE INVENTION

The invention is based on the discovery of new human cytoskeleton-associated proteins (CYSKP), the polynucleotides encoding CYSKP, and the use of these compositions for the diagnosis, treatment, or prevention of cell proliferative, autoimmune/inflammatory, vesicle trafficking, neurological, cell motility, reproductive, and muscle disorders.

Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide sequences of the invention. Each polynucleotide and its corresponding polypeptide are correlated to a single Incyte project identification number (Incyte Project ID). Each polypeptide sequence is denoted by both a polypeptide sequence identification number (Polypeptide SEQ ID NO:) and an Incyte polypeptide sequence number (Incyte Polypeptide ID) as shown. Each polynucleotide sequence is denoted by both a polynucleotide sequence identification number (Polynucleotide SEQ ID NO:) and an Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) as shown.

Table 2 shows sequences with homology to the polypeptides of the invention as identified by BLAST analysis against the GenBank protein (genpept) database. Columns 1 and 2 show the polypeptide sequence identification number (Polypeptide SEQ ID NO:) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for polypeptides of the invention. Column 3 shows the GenBank identification number (Genbank ID NO:) of the nearest GenBank homolog. Column 4 shows the probability score for the match between each polypeptide and its GenBank homolog. Column 5 shows the annotation of the GenBank homolog along with relevant citations where applicable, all of which are expressly incorporated by reference herein.

Table 3 shows various structural features of the polypeptides of the invention. Columns 1 and 2 show the polypeptide sequence identification number (SEQ ID NO:) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for each polypeptide of the invention. Column 3 shows the number of amino acid residues in each polypeptide. Column 4 shows potential phosphorylation sites, and column 5 shows potential glycosylation sites, as determined by the MOTIFS program of the GCG sequence analysis software package (Genetics Computer Group, Madison WI). Column 6 shows amino acid residues comprising signature sequences, domains, and motifs. Column 7 shows analytical methods for protein structure/function analysis and in some cases, searchable databases to which the analytical methods were applied.

Together, Tables 2 and 3 summarize the properties of polypeptides of the invention, and these properties establish that the claimed polypeptides are cytoskeleton-associated proteins. For example, SEQ ID NO:31 is 34% identical to a *Caenorhabditis elegans* protein similar to mouse ankyrin

(GenBank ID g3879121) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is  $1.1\text{e-}146$ , which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:31 also contains Ank repeats as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. As a second example, SEQ ID NO:34 is 96% identical over 97 amino acids to human Intermediate Filament Associated Protein (GenBank ID 1333846) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is  $8.2\text{e-}45$ , which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. Data from BLAST analyses using the PRODOM database provide further corroborative evidence that SEQ ID NO:34 is a cytoskeleton protein. (See Table 3.) SEQ ID NO:1-30 and SEQ ID NO:32-33 were analyzed and annotated in a similar manner. The algorithms and parameters for the analysis of SEQ ID NO:1-34 are described in Table 7.

As shown in Table 4, the full length polynucleotide sequences of the present invention were assembled using cDNA sequences or coding (exon) sequences derived from genomic DNA, or any combination of these two types of sequences. Columns 1 and 2 list the polynucleotide sequence identification number (Polynucleotide SEQ ID NO:) and the corresponding Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) for each polynucleotide of the invention. Column 3 shows the length of each polynucleotide sequence in basepairs. Column 4 lists fragments of the polynucleotide sequences which are useful, for example, in hybridization or amplification technologies that identify SEQ ID NO:35-68 or that distinguish between SEQ ID NO:35-68 and related polynucleotide sequences. Column 5 shows identification numbers corresponding to cDNA sequences, coding sequences (exons) predicted from genomic DNA, and/or sequence assemblages comprised of both cDNA and genomic DNA. These sequences were used to assemble the full length polynucleotide sequences of the invention. Columns 6 and 7 of Table 4 show the nucleotide start (5') and stop (3') positions of the cDNA and genomic sequences in column 5 relative to their respective full length sequences.

The identification numbers in Column 5 of Table 4 may refer specifically, for example, to Incyte cDNAs along with their corresponding cDNA libraries. For example, 3824958H1 is the identification number of an Incyte cDNA sequence, and BRAXNOT01 is the cDNA library from which it is derived. Incyte cDNAs for which cDNA libraries are not indicated were derived from pooled cDNA libraries (e.g., 71263527V1). Alternatively, the identification numbers in column 5 may refer to GenBank cDNAs or ESTs (e.g., g2276318) which contributed to the assembly of the full length polynucleotide sequences. Alternatively, the identification numbers in column 5 may refer to coding regions predicted by Genscan analysis of genomic DNA. The Genscan-predicted coding sequences may

have been edited prior to assembly. (See Example IV.) Alternatively, the identification numbers in column 5 may refer to assemblages of both cDNA and Genscan-predicted exons brought together by an “exon stitching” algorithm. (See Example V.) Alternatively, the identification numbers in column 5 may refer to assemblages of both cDNA and Genscan-predicted exons brought together by an “exon-stretching” algorithm. (See Example V.) In some cases, Incyte cDNA coverage redundant with the sequence coverage shown in column 5 was obtained to confirm the final consensus polynucleotide sequence, but the relevant Incyte cDNA identification numbers are not shown.

Table 5 shows the representative cDNA libraries for those full length polynucleotide sequences which were assembled using Incyte cDNA sequences. The representative cDNA library is the Incyte cDNA library which is most frequently represented by the Incyte cDNA sequences which were used to assemble and confirm the above polynucleotide sequences. The tissues and vectors which were used to construct the cDNA libraries shown in Table 5 are described in Table 6.

The invention also encompasses CYSKP variants. A preferred CYSKP variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the CYSKP amino acid sequence, and which contains at least one functional or structural characteristic of CYSKP.

The invention also encompasses polynucleotides which encode CYSKP. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:35-68, which encodes CYSKP. The polynucleotide sequences of SEQ ID NO:35-68, as presented in the Sequence Listing, embrace the equivalent RNA sequences, wherein occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The invention also encompasses a variant of a polynucleotide sequence encoding CYSKP. In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding CYSKP. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:35-68 which has at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:35-68. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of CYSKP.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding CYSKP, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the

invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring CYSKP, and all such variations are to be considered as being specifically disclosed.

5           Although nucleotide sequences which encode CYSKP and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring CYSKP under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding CYSKP or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide  
10 occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding CYSKP and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

15           The invention also encompasses production of DNA sequences which encode CYSKP and CYSKP derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding CYSKP or any fragment thereof.

20           Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:35-68 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in  
25 "Definitions."

          Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Applied Biosystems), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or  
30 combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (Applied Biosystems). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing

system (Applied Biosystems), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding CYSKP may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) PCR Methods Applic. 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) Nucleic Acids Res. 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) PCR Methods Applic. 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-3060). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 primer analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-

specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Applied Biosystems), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode CYSKP may be cloned in recombinant DNA molecules that direct expression of CYSKP, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express CYSKP.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter CYSKP-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

The nucleotides of the present invention may be subjected to DNA shuffling techniques such as MOLECULARBREEDING (Maxygen Inc., Santa Clara CA; described in U.S. Patent Number 5,837,458; Chang, C.-C. et al. (1999) Nat. Biotechnol. 17:793-797; Christians, F.C. et al. (1999) Nat. Biotechnol. 17:259-264; and Cramer, A. et al. (1996) Nat. Biotechnol. 14:315-319) to alter or improve the biological properties of CYSKP, such as its biological or enzymatic activity or its ability to bind to other molecules or compounds. DNA shuffling is a process by which a library of gene variants is produced using PCR-mediated recombination of gene fragments. The library is then subjected to selection or screening procedures that identify those gene variants with the desired properties. These preferred variants may then be pooled and further subjected to recursive rounds of DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial" breeding and rapid molecular evolution. For example, fragments of a single gene containing random point mutations may be recombined, screened, and then reshuffled until the desired properties are optimized. Alternatively, fragments of a given gene may be recombined with fragments of homologous genes in the same gene family, either from the same or different species, thereby maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.



In another embodiment, sequences encoding CYSKP may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) Nucleic Acids Symp. Ser. 7:215-223; and Horn, T. et al. (1980) Nucleic Acids Symp. Ser. 7:225-232.) Alternatively, CYSKP itself or a fragment thereof may be synthesized using chemical methods. For example, peptide  
5 synthesis can be performed using various solution-phase or solid-phase techniques. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY, pp. 55-60; and Roberge, J.Y. et al. (1995) Science 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Applied Biosystems). Additionally, the amino acid sequence of CYSKP, or any part thereof, may be altered during direct synthesis and/or combined with sequences  
10 from other proteins, or any part thereof, to produce a variant polypeptide or a polypeptide having a sequence of a naturally occurring polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) Methods Enzymol. 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing.  
15 (See, e.g., Creighton, supra, pp. 28-53.)

In order to express a biologically active CYSKP, the nucleotide sequences encoding CYSKP or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and  
20 inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding CYSKP. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding CYSKP. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding CYSKP and its initiation codon and upstream regulatory sequences are inserted  
25 into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers  
30 appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) Results Probl. Cell Differ. 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding CYSKP and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in

vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding CYSKP. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. (See, e.g., Sambrook, supra; Ausubel, supra; Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509; Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945; Takamatsu, N. (1987) EMBO J. 6:307-311; The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196; Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659; and Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.) Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. (See, e.g., Di Nicola, M. et al. (1998) Cancer Gen. Ther. 5(6):350-356; Yu, M. et al. (1993) Proc. Natl. Acad. Sci. USA 90(13):6340-6344; Buller, R.M. et al. (1985) Nature 317(6040):813-815; McGregor, D.P. et al. (1994) Mol. Immunol. 31(3):219-226; and Verma, I.M. and N. Somia (1997) Nature 389:239-242.) The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding CYSKP. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding CYSKP can be achieved using a multifunctional E. coli vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1 plasmid (Life Technologies). Ligation of sequences encoding CYSKP into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for in vitro transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of CYSKP are needed, e.g. for the production of antibodies, vectors which direct high level expression of CYSKP may be used. For example, vectors containing the strong, inducible SP6 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of CYSKP. A number of vectors

containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast Saccharomyces cerevisiae or Pichia pastoris. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, supra;  
5 Bitter, G.A. et al. (1987) Methods Enzymol. 153:516-544; and Scorer, C.A. et al. (1994) Bio/Technology 12:181-184.)

Plant systems may also be used for expression of CYSKP. Transcription of sequences encoding CYSKP may be driven by viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J.  
10 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al. (1984) Science 224:838-843; and Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill,  
15 New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding CYSKP may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain  
20 infective virus which expresses CYSKP in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of  
25 DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of CYSKP in cell lines is preferred. For example, sequences encoding CYSKP can be transformed into  
30 cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the

introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk<sup>-</sup>* and *apr<sup>-</sup>* cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; 5 Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) 10 Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech),  $\beta$  glucuronidase and its substrate  $\beta$ -glucuronide, or luciferase and its substrate luciferin may be used. 15 These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the 20 sequence encoding CYSKP is inserted within a marker gene sequence, transformed cells containing sequences encoding CYSKP can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding CYSKP under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

25 In general, host cells that contain the nucleic acid sequence encoding CYSKP and that express CYSKP may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

30 Immunological methods for detecting and measuring the expression of CYSKP using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on CYSKP is preferred, but a competitive binding

assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

5           A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding CYSKP include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding CYSKP, or any fragments thereof, may be cloned into a vector  
10       for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease  
15       of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding CYSKP may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence  
20       and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode CYSKP may be designed to contain signal sequences which direct secretion of CYSKP through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the  
25       polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture  
30       Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding CYSKP may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric CYSKP protein

containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of CYSKP activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose  
5 binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion  
10 proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the CYSKP encoding sequence and the heterologous protein sequence, so that CYSKP may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, supra, ch. 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

15 In a further embodiment of the invention, synthesis of radiolabeled CYSKP may be achieved in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, <sup>35</sup>S-methionine.

20 CYSKP of the present invention or fragments thereof may be used to screen for compounds that specifically bind to CYSKP. At least one and up to a plurality of test compounds may be screened for specific binding to CYSKP. Examples of test compounds include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

In one embodiment, the compound thus identified is closely related to the natural ligand of  
25 CYSKP, e.g., a ligand or fragment thereof, a natural substrate, a structural or functional mimetic, or a natural binding partner. (See, e.g., Coligan, J.E. et al. (1991) Current Protocols in Immunology 1(2): Chapter 5.) Similarly, the compound can be closely related to the natural receptor to which CYSKP binds, or to at least a fragment of the receptor, e.g., the ligand binding site. In either case, the compound can be rationally designed using known techniques. In one embodiment, screening for  
30 these compounds involves producing appropriate cells which express CYSKP, either as a secreted protein or on the cell membrane. Preferred cells include cells from mammals, yeast, Drosophila, or E. coli. Cells expressing CYSKP or cell membrane fractions which contain CYSKP are then contacted with a test compound and binding, stimulation, or inhibition of activity of either CYSKP or the compound is analyzed.

An assay may simply test binding of a test compound to the polypeptide, wherein binding is detected by a fluorophore, radioisotope, enzyme conjugate, or other detectable label. For example, the assay may comprise the steps of combining at least one test compound with CYSKP, either in solution or affixed to a solid support, and detecting the binding of CYSKP to the compound.

5 Alternatively, the assay may detect or measure binding of a test compound in the presence of a labeled competitor. Additionally, the assay may be carried out using cell-free preparations, chemical libraries, or natural product mixtures, and the test compound(s) may be free in solution or affixed to a solid support.

CYSKP of the present invention or fragments thereof may be used to screen for compounds  
10 that modulate the activity of CYSKP. Such compounds may include agonists, antagonists, or partial or inverse agonists. In one embodiment, an assay is performed under conditions permissive for CYSKP activity, wherein CYSKP is combined with at least one test compound, and the activity of CYSKP in the presence of a test compound is compared with the activity of CYSKP in the absence of the test compound. A change in the activity of CYSKP in the presence of the test compound is  
15 indicative of a compound that modulates the activity of CYSKP. Alternatively, a test compound is combined with an in vitro or cell-free system comprising CYSKP under conditions suitable for CYSKP activity, and the assay is performed. In either of these assays, a test compound which modulates the activity of CYSKP may do so indirectly and need not come in direct contact with the test compound. At least one and up to a plurality of test compounds may be screened.

20 In another embodiment, polynucleotides encoding CYSKP or their mammalian homologs may be "knocked out" in an animal model system using homologous recombination in embryonic stem (ES) cells. Such techniques are well known in the art and are useful for the generation of animal models of human disease. (See, e.g., U.S. Patent Number 5,175,383 and U.S. Patent Number 5,767,337.) For example, mouse ES cells, such as the mouse 129/SvJ cell line, are derived from the early mouse embryo  
25 and grown in culture. The ES cells are transformed with a vector containing the gene of interest disrupted by a marker gene, e.g., the neomycin phosphotransferase gene (neo; Capecchi, M.R. (1989) Science 244:1288-1292). The vector integrates into the corresponding region of the host genome by homologous recombination. Alternatively, homologous recombination takes place using the Cre-loxP system to knockout a gene of interest in a tissue- or developmental stage-specific manner (Marth, J.D.  
30 (1996) Clin. Invest. 97:1999-2002; Wagner, K.U. et al. (1997) Nucleic Acids Res. 25:4323-4330). Transformed ES cells are identified and microinjected into mouse cell blastocysts such as those from the C57BL/6 mouse strain. The blastocysts are surgically transferred to pseudopregnant dams, and the resulting chimeric progeny are genotyped and bred to produce heterozygous or homozygous strains. Transgenic animals thus generated may be tested with potential therapeutic or toxic agents.

Polynucleotides encoding CYSKP may also be manipulated in vitro in ES cells derived from human blastocysts. Human ES cells have the potential to differentiate into at least eight separate cell lineages including endoderm, mesoderm, and ectodermal cell types. These cell lineages differentiate into, for example, neural cells, hematopoietic lineages, and cardiomyocytes (Thomson, J.A. et al. (1998) Science 282:1145-1147).

Polynucleotides encoding CYSKP can also be used to create “knockin” humanized animals (pigs) or transgenic animals (mice or rats) to model human disease. With knockin technology, a region of a polynucleotide encoding CYSKP is injected into animal ES cells, and the injected sequence integrates into the animal cell genome. Transformed cells are injected into blastulae, and the blastulae are implanted as described above. Transgenic progeny or inbred lines are studied and treated with potential pharmaceutical agents to obtain information on treatment of a human disease. Alternatively, a mammal inbred to overexpress CYSKP, e.g., by secreting CYSKP in its milk, may also serve as a convenient source of that protein (Janne, J. et al. (1998) Biotechnol. Annu. Rev. 4:55-74).

#### THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of CYSKP and cytoskeleton-associated proteins. In addition, the expression of CYSKP is closely associated with lung, reproductive (including placenta), neural (including brain), adrenal, endothelial, kidney, and spleen tissue, as well as with ovarian, breast, and testicular tumor tissue. Therefore, CYSKP appears to play a role in cell proliferative, autoimmune/inflammatory, vesicle trafficking, neurological, cell motility, reproductive, and muscle disorders. In the treatment of disorders associated with increased CYSKP expression or activity, it is desirable to decrease the expression or activity of CYSKP. In the treatment of disorders associated with decreased CYSKP expression or activity, it is desirable to increase the expression or activity of CYSKP.

Therefore, in one embodiment, CYSKP or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of CYSKP. Examples of such disorders include, but are not limited to, a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; an autoimmune/inflammatory disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory



distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with

5 lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus,

10 systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a vesicle trafficking disorder such as cystic fibrosis, glucose-galactose malabsorption syndrome, hypercholesterolemia, diabetes mellitus, diabetes insipidus, hyper- and hypoglycemia, Grave's disease, goiter, Cushing's disease, and Addison's disease,

15 gastrointestinal disorders including ulcerative colitis, gastric and duodenal ulcers, other conditions associated with abnormal vesicle trafficking, including acquired immunodeficiency syndrome (AIDS), allergies including hay fever, asthma, and urticaria (hives), autoimmune hemolytic anemia, proliferative glomerulonephritis, inflammatory bowel disease, multiple sclerosis, myasthenia gravis, rheumatoid and osteoarthritis, scleroderma, Chediak-Higashi and Sjogren's syndromes, systemic lupus erythematosus,

20 toxic shock syndrome, traumatic tissue damage, and viral, bacterial, fungal, helminthic, and protozoal infections; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and

25 other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal

30 syndrome, mental retardation and other developmental disorders of the central nervous system including Down syndrome, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety,

and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; a cell motility disorder such as ankylosing spondylitis, Chediak-Higashi syndrome, Duchenne and Becker muscular dystrophy, intrahepatic cholestasis, myocardial hyperplasia, cardiomyopathy, 5 early onset peridontitis, cancers such as adenocarcinoma, ovarian carcinoma, and chronic myelogenous leukemia, and bacterial and helminthic infections; a reproductive disorder such as a disorder of prolactin production, infertility, including tubal disease, ovulatory defects, endometriosis, a disruption of the estrous cycle, a disruption of the menstrual cycle, polycystic ovary syndrome, ovarian 10 hyperstimulation syndrome, an endometrial or ovarian tumor, a uterine fibroid, autoimmune disorders, ectopic pregnancy, teratogenesis, cancer of the breast, fibrocystic breast disease, galactorrhea, a disruption of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the male breast, gynecomastia, hypergonadotropic and hypogonadotropic hypogonadism, pseudohermaphroditism, 15 azoospermia, premature ovarian failure, acrosin deficiency, delayed puberty, retrograde ejaculation and anejaculation, haemangioblastomas, cystsphaeochromocytomas, paraganglioma, cystadenomas of the epididymis, and endolymphatic sac tumours; and a muscle disorder such as myocarditis, Duchenne's muscular dystrophy, Becker's muscular dystrophy, myotonic dystrophy, central core disease, nemaline myopathy, centronuclear myopathy, lipid myopathy, mitochondrial myopathy, infectious myositis, 20 polymyositis, dermatomyositis, inclusion body myositis, thyrotoxic myopathy, and ethanol myopathy, angina, anaphylactic shock, arrhythmias, asthma, cardiovascular shock, Cushing's syndrome, hypertension, hypoglycemia, myocardial infarction, migraine, and pheochromocytoma, and myopathies including cardiomyopathy, encephalopathy, epilepsy, Kearns-Sayre syndrome, lactic acidosis, myoclonic disorder, and ophthalmoplegia.

25 In another embodiment, a vector capable of expressing CYSKP or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of CYSKP including, but not limited to, those described above.

In a further embodiment, a composition comprising a substantially purified CYSKP in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a 30 disorder associated with decreased expression or activity of CYSKP including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of CYSKP may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of CYSKP including, but not limited to, those listed above.

In a further embodiment, an antagonist of CYSKP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of CYSKP. Examples of such disorders include, but are not limited to, those cell proliferative, autoimmune/inflammatory, vesicle trafficking, neurological, cell motility, reproductive, and muscle disorders described above. In one aspect, an antibody which specifically binds CYSKP may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express CYSKP.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding CYSKP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of CYSKP including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of CYSKP may be produced using methods which are generally known in the art. In particular, purified CYSKP may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind CYSKP. Antibodies to CYSKP may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with CYSKP or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to CYSKP have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or

fragments are identical to a portion of the amino acid sequence of the natural protein. Short stretches of CYSKP amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to CYSKP may be prepared using any technique which provides for the  
5 production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) Nature 256:495-497; Kozbor, D. et al. (1985) J. Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) Proc. Natl. Acad. Sci. USA 80:2026-2030; and Cole, S.P. et al. (1984) Mol. Cell Biol. 62:109-120.)

10 In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) Proc. Natl. Acad. Sci. USA 81:6851-6855; Neuberger, M.S. et al. (1984) Nature 312:604-608; and Takeda, S. et al. (1985) Nature 314:452-454.) Alternatively, techniques described for the production of single  
15 chain antibodies may be adapted, using methods known in the art, to produce CYSKP-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) Proc. Natl. Acad. Sci. USA 88:10134-10137.)

Antibodies may also be produced by inducing in vivo production in the lymphocyte population  
20 or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) Proc. Natl. Acad. Sci. USA 86:3833-3837; Winter, G. et al. (1991) Nature 349:293-299.)

Antibody fragments which contain specific binding sites for CYSKP may also be generated. For example, such fragments include, but are not limited to, F(ab')<sub>2</sub> fragments produced by pepsin  
25 digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')<sub>2</sub> fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) Science 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired  
30 specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between CYSKP and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering CYSKP epitopes is generally used, but a competitive binding assay may also be

employed (Pound, supra).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for CYSKP. Affinity is expressed as an association constant,  $K_a$ , which is defined as the molar concentration of CYSKP-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The  $K_a$  determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple CYSKP epitopes, represents the average affinity, or avidity, of the antibodies for CYSKP. The  $K_a$  determined for a preparation of monoclonal antibodies, which are monospecific for a particular CYSKP epitope, represents a true measure of affinity. High-affinity antibody preparations with  $K_a$  ranging from about  $10^9$  to  $10^{12}$  L/mole are preferred for use in immunoassays in which the CYSKP-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with  $K_a$  ranging from about  $10^6$  to  $10^7$  L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of CYSKP, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington DC; Liddell, J.E. and A. Cryer (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of CYSKP-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al. supra.)

In another embodiment of the invention, the polynucleotides encoding CYSKP, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, modifications of gene expression can be achieved by designing complementary sequences or antisense molecules (DNA, RNA, PNA, or modified oligonucleotides) to the coding or regulatory regions of the gene encoding CYSKP. Such technology is well known in the art, and antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding CYSKP. (See, e.g., Agrawal, S., ed. (1996) Antisense Therapeutics, Humana Press Inc., Totawa NJ.)

In therapeutic use, any gene delivery system suitable for introduction of the antisense sequences into appropriate target cells can be used. Antisense sequences can be delivered intracellularly in the form of an expression plasmid which, upon transcription, produces a sequence complementary to at least a portion of the cellular sequence encoding the target protein. (See, e.g., Slater, J.E. et al. (1998) *J. Allergy Clin. Immunol.* 102(3):469-475; and Scanlon, K.J. et al. (1995)

9(13):1288-1296.) Antisense sequences can also be introduced intracellularly through the use of viral vectors, such as retrovirus and adeno-associated virus vectors. (See, e.g., Miller, A.D. (1990) *Blood* 76:271; Ausubel, supra; Uckert, W. and W. Walther (1994) *Pharmacol. Ther.* 63(3):323-347.) Other gene delivery mechanisms include liposome-derived systems, artificial viral envelopes, and other systems known in the art. (See, e.g., Rossi, J.J. (1995) *Br. Med. Bull.* 51(1):217-225; Boado, R.J. et al. (1998) *J. Pharm. Sci.* 87(11):1308-1315; and Morris, M.C. et al. (1997) *Nucleic Acids Res.* 25(14):2730-2736.)

In another embodiment of the invention, polynucleotides encoding CYSKP may be used for somatic or germline gene therapy. Gene therapy may be performed to (i) correct a genetic deficiency (e.g., in the cases of severe combined immunodeficiency (SCID)-X1 disease characterized by X-linked inheritance (Cavazzana-Calvo, M. et al. (2000) *Science* 288:669-672), severe combined immunodeficiency syndrome associated with an inherited adenosine deaminase (ADA) deficiency (Blaese, R.M. et al. (1995) *Science* 270:475-480; Bordignon, C. et al. (1995) *Science* 270:470-475), cystic fibrosis (Zabner, J. et al. (1993) *Cell* 75:207-216; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:643-666; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:667-703), thalassemias, familial hypercholesterolemia, and hemophilia resulting from Factor VIII or Factor IX deficiencies (Crystal, R.G. (1995) *Science* 270:404-410; Verma, I.M. and N. Somia (1997) *Nature* 389:239-242)), (ii) express a conditionally lethal gene product (e.g., in the case of cancers which result from unregulated cell proliferation), or (iii) express a protein which affords protection against intracellular parasites (e.g., against human retroviruses, such as human immunodeficiency virus (HIV) (Baltimore, D. (1988) *Nature* 335:395-396; Poeschla, E. et al. (1996) *Proc. Natl. Acad. Sci. USA.* 93:11395-11399), hepatitis B or C virus (HBV, HCV); fungal parasites, such as Candida albicans and Paracoccidioides brasiliensis; and protozoan parasites such as Plasmodium falciparum and Trypanosoma cruzi). In the case where a genetic deficiency in CYSKP expression or regulation causes disease, the expression of CYSKP from an appropriate population of transduced cells may alleviate the clinical manifestations caused by the genetic deficiency.

In a further embodiment of the invention, diseases or disorders caused by deficiencies in CYSKP are treated by constructing mammalian expression vectors encoding CYSKP and introducing these vectors by mechanical means into CYSKP-deficient cells. Mechanical transfer technologies for use with cells in vivo or ex vitro include (i) direct DNA microinjection into individual cells, (ii) ballistic gold particle delivery, (iii) liposome-mediated transfection, (iv) receptor-mediated gene transfer, and (v) the use of DNA transposons (Morgan, R.A. and W.F. Anderson (1993) *Annu. Rev. Biochem.* 62:191-217; Ivics, Z. (1997) *Cell* 91:501-510; Boulay, J-L. and H. Récipon (1998) *Curr. Opin. Biotechnol.* 9:445-450).

Expression vectors that may be effective for the expression of CYSKP include, but are not limited to, the PCDNA 3.1, EPITAG, PRCCMV2, PREP, PVAX vectors (Invitrogen, Carlsbad CA), PCMV-SCRIPT, PCMV-TAG, PEGSH/PERV (Stratagene, La Jolla CA), and PTET-OFF, PTET-ON, PTRE2, PTRE2-LUC, PTK-HYG (Clontech, Palo Alto CA). CYSKP may be expressed

5 using (i) a constitutively active promoter, (e.g., from cytomegalovirus (CMV), Rous sarcoma virus (RSV), SV40 virus, thymidine kinase (TK), or  $\beta$ -actin genes), (ii) an inducible promoter (e.g., the tetracycline-regulated promoter (Gossen, M. and H. Bujard (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551; Gossen, M. et al. (1995) *Science* 268:1766-1769; Rossi, F.M.V. and H.M. Blau (1998) *Curr. Opin. Biotechnol.* 9:451-456), commercially available in the T-REX plasmid (Invitrogen)); the

10 ecdysone-inducible promoter (available in the plasmids PVGRXR and PIND; Invitrogen); the FK506/rapamycin inducible promoter; or the RU486/mifepristone inducible promoter (Rossi, F.M.V. and Blau, H.M. *supra*)), or (iii) a tissue-specific promoter or the native promoter of the endogenous gene encoding CYSKP from a normal individual.

Commercially available liposome transformation kits (e.g., the PERFECT LIPID

15 TRANSFECTION KIT, available from Invitrogen) allow one with ordinary skill in the art to deliver polynucleotides to target cells in culture and require minimal effort to optimize experimental parameters. In the alternative, transformation is performed using the calcium phosphate method (Graham, F.L. and A.J. Eb (1973) *Virology* 52:456-467), or by electroporation (Neumann, E. et al. (1982) *EMBO J.* 1:841-845). The introduction of DNA to primary cells requires modification of these

20 standardized mammalian transfection protocols.

In another embodiment of the invention, diseases or disorders caused by genetic defects with respect to CYSKP expression are treated by constructing a retrovirus vector consisting of (i) the polynucleotide encoding CYSKP under the control of an independent promoter or the retrovirus long terminal repeat (LTR) promoter, (ii) appropriate RNA packaging signals, and (iii) a Rev-responsive

25 element (RRE) along with additional retrovirus *cis*-acting RNA sequences and coding sequences required for efficient vector propagation. Retrovirus vectors (e.g., PFB and PFBNEO) are commercially available (Stratagene) and are based on published data (Riviere, I. et al. (1995) *Proc. Natl. Acad. Sci. USA* 92:6733-6737), incorporated by reference herein. The vector is propagated in an appropriate vector producing cell line (VPCL) that expresses an envelope gene with a tropism for

30 receptors on the target cells or a promiscuous envelope protein such as VSVg (Armentano, D. et al. (1987) *J. Virol.* 61:1647-1650; Bender, M.A. et al. (1987) *J. Virol.* 61:1639-1646; Adam, M.A. and A.D. Miller (1988) *J. Virol.* 62:3802-3806; Dull, T. et al. (1998) *J. Virol.* 72:8463-8471; Zufferey, R. et al. (1998) *J. Virol.* 72:9873-9880). U.S. Patent Number 5,910,434 to Rigg ("Method for obtaining retrovirus packaging cell lines producing high transducing efficiency retroviral supernatant") discloses a

method for obtaining retrovirus packaging cell lines and is hereby incorporated by reference.

Propagation of retrovirus vectors, transduction of a population of cells (e.g., CD4<sup>+</sup> T-cells), and the return of transduced cells to a patient are procedures well known to persons skilled in the art of gene therapy and have been well documented (Ranga, U. et al. (1997) J. Virol. 71:7020-7029; Bauer, G. et al. (1997) Blood 89:2259-2267; Bonyhadi, M.L. (1997) J. Virol. 71:4707-4716; Ranga, U. et al. (1998) Proc. Natl. Acad. Sci. USA 95:1201-1206; Su, L. (1997) Blood 89:2283-2290).

In the alternative, an adenovirus-based gene therapy delivery system is used to deliver polynucleotides encoding CYSKP to cells which have one or more genetic abnormalities with respect to the expression of CYSKP. The construction and packaging of adenovirus-based vectors are well known to those with ordinary skill in the art. Replication defective adenovirus vectors have proven to be versatile for importing genes encoding immunoregulatory proteins into intact islets in the pancreas (Csete, M.E. et al. (1995) Transplantation 27:263-268). Potentially useful adenoviral vectors are described in U.S. Patent Number 5,707,618 to Armentano ("Adenovirus vectors for gene therapy"), hereby incorporated by reference. For adenoviral vectors, see also Antinozzi, P.A. et al. (1999) Annu. Rev. Nutr. 19:511-544 and Verma, I.M. and N. Somia (1997) Nature 18:389:239-242, both incorporated by reference herein.

In another alternative, a herpes-based, gene therapy delivery system is used to deliver polynucleotides encoding CYSKP to target cells which have one or more genetic abnormalities with respect to the expression of CYSKP. The use of herpes simplex virus (HSV)-based vectors may be especially valuable for introducing CYSKP to cells of the central nervous system, for which HSV has a tropism. The construction and packaging of herpes-based vectors are well known to those with ordinary skill in the art. A replication-competent herpes simplex virus (HSV) type 1-based vector has been used to deliver a reporter gene to the eyes of primates (Liu, X. et al. (1999) Exp. Eye Res. 169:385-395). The construction of a HSV-1 virus vector has also been disclosed in detail in U.S. Patent Number 5,804,413 to DeLuca ("Herpes simplex virus strains for gene transfer"), which is hereby incorporated by reference. U.S. Patent Number 5,804,413 teaches the use of recombinant HSV d92 which consists of a genome containing at least one exogenous gene to be transferred to a cell under the control of the appropriate promoter for purposes including human gene therapy. Also taught by this patent are the construction and use of recombinant HSV strains deleted for ICP4, ICP27 and ICP22. For HSV vectors, see also Goins, W.F. et al. (1999) J. Virol. 73:519-532 and Xu, H. et al. (1994) Dev. Biol. 163:152-161, hereby incorporated by reference. The manipulation of cloned herpesvirus sequences, the generation of recombinant virus following the transfection of multiple plasmids containing different segments of the large herpesvirus genomes, the growth and propagation of herpesvirus, and the infection of cells with herpesvirus are techniques well known to those of ordinary



skill in the art.

In another alternative, an alphavirus (positive, single-stranded RNA virus) vector is used to deliver polynucleotides encoding CYSKP to target cells. The biology of the prototypic alphavirus, Semliki Forest Virus (SFV), has been studied extensively and gene transfer vectors have been based on the SFV genome (Garoff, H. and K.-J. Li (1998) Curr. Opin. Biotechnol. 9:464-469). During alphavirus RNA replication, a subgenomic RNA is generated that normally encodes the viral capsid proteins. This subgenomic RNA replicates to higher levels than the full length genomic RNA, resulting in the overproduction of capsid proteins relative to the viral proteins with enzymatic activity (e.g., protease and polymerase). Similarly, inserting the coding sequence for CYSKP into the alphavirus genome in place of the capsid-coding region results in the production of a large number of CYSKP-coding RNAs and the synthesis of high levels of CYSKP in vector transduced cells. While alphavirus infection is typically associated with cell lysis within a few days, the ability to establish a persistent infection in hamster normal kidney cells (BHK-21) with a variant of Sindbis virus (SIN) indicates that the lytic replication of alphaviruses can be altered to suit the needs of the gene therapy application (Dryga, S.A. et al. (1997) Virology 228:74-83). The wide host range of alphaviruses will allow the introduction of CYSKP into a variety of cell types. The specific transduction of a subset of cells in a population may require the sorting of cells prior to transduction. The methods of manipulating infectious cDNA clones of alphaviruses, performing alphavirus cDNA and RNA transfections, and performing alphavirus infections, are well known to those with ordinary skill in the art.

Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may also be employed to inhibit gene expression. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding CYSKP.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by

scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis.

Alternatively, RNA molecules may be generated by *in vitro* and *in vivo* transcription of DNA sequences encoding CYSKP. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

An additional embodiment of the invention encompasses a method for screening for a compound which is effective in altering expression of a polynucleotide encoding CYSKP.

Compounds which may be effective in altering expression of a specific polynucleotide may include, but are not limited to, oligonucleotides, antisense oligonucleotides, triple helix-forming oligonucleotides, transcription factors and other polypeptide transcriptional regulators, and non-macromolecular chemical entities which are capable of interacting with specific polynucleotide sequences. Effective compounds may alter polynucleotide expression by acting as either inhibitors or promoters of polynucleotide expression. Thus, in the treatment of disorders associated with increased CYSKP expression or activity, a compound which specifically inhibits expression of the polynucleotide encoding CYSKP may be therapeutically useful, and in the treatment of disorders associated with decreased CYSKP expression or activity, a compound which specifically promotes expression of the polynucleotide encoding CYSKP may be therapeutically useful.

At least one, and up to a plurality, of test compounds may be screened for effectiveness in

altering expression of a specific polynucleotide. A test compound may be obtained by any method commonly known in the art, including chemical modification of a compound known to be effective in altering polynucleotide expression; selection from an existing, commercially-available or proprietary library of naturally-occurring or non-natural chemical compounds; rational design of a compound  
5 based on chemical and/or structural properties of the target polynucleotide; and selection from a library of chemical compounds created combinatorially or randomly. A sample comprising a polynucleotide encoding CYSKP is exposed to at least one test compound thus obtained. The sample may comprise, for example, an intact or permeabilized cell, or an in vitro cell-free or reconstituted biochemical system. Alterations in the expression of a polynucleotide encoding CYSKP are assayed  
10 by any method commonly known in the art. Typically, the expression of a specific nucleotide is detected by hybridization with a probe having a nucleotide sequence complementary to the sequence of the polynucleotide encoding CYSKP. The amount of hybridization may be quantified, thus forming the basis for a comparison of the expression of the polynucleotide both with and without exposure to one or more test compounds. Detection of a change in the expression of a polynucleotide  
15 exposed to a test compound indicates that the test compound is effective in altering the expression of the polynucleotide. A screen for a compound effective in altering expression of a specific polynucleotide can be carried out, for example, using a Schizosaccharomyces pombe gene expression system (Atkins, D. et al. (1999) U.S. Patent No. 5,932,435; Arndt, G.M. et al. (2000) Nucleic Acids Res. 28:E15) or a human cell line such as HeLa cell (Clarke, M.L. et al. (2000) Biochem. Biophys. Res. Commun. 268:8-13). A particular embodiment of the present invention involves screening a  
20 combinatorial library of oligonucleotides (such as deoxyribonucleotides, ribonucleotides, peptide nucleic acids, and modified oligonucleotides) for antisense activity against a specific polynucleotide sequence (Bruice, T.W. et al. (1997) U.S. Patent No. 5,686,242; Bruice, T.W. et al. (2000) U.S. Patent No. 6,022,691).

25 Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat.  
30 Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a composition which  
35 generally comprises an active ingredient formulated with a pharmaceutically acceptable excipient.

Excipients may include, for example, sugars, starches, celluloses, gums, and proteins. Various formulations are commonly known and are thoroughly discussed in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA). Such compositions may consist of CYSKP, antibodies to CYSKP, and mimetics, agonists, antagonists, or inhibitors of CYSKP.

5           The compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, pulmonary, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

Compositions for pulmonary administration may be prepared in liquid or dry powder form.

10       These compositions are generally aerosolized immediately prior to inhalation by the patient. In the case of small molecules (e.g. traditional low molecular weight organic drugs), aerosol delivery of fast-acting formulations is well-known in the art. In the case of macromolecules (e.g. larger peptides and proteins), recent developments in the field of pulmonary delivery via the alveolar region of the lung have enabled the practical delivery of drugs such as insulin to blood circulation (see, e.g., Patton, J.S. et al., U.S. Patent No. 5,997,848). Pulmonary delivery has the advantage of administration without needle  
15       injection, and obviates the need for potentially toxic penetration enhancers.

Compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

20       Specialized forms of compositions may be prepared for direct intracellular delivery of macromolecules comprising CYSKP or fragments thereof. For example, liposome preparations containing a cell-impermeable macromolecule may promote cell fusion and intracellular delivery of the macromolecule. Alternatively, CYSKP or a fragment thereof may be joined to a short cationic N-terminal portion from the HIV Tat-1 protein. Fusion proteins thus generated have been found to  
25       transduce into the cells of all tissues, including the brain, in a mouse model system (Schwarze, S.R. et al. (1999) Science 285:1569-1572).

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, monkeys, or pigs. An animal model may also be used to determine the appropriate concentration range and route  
30       of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example CYSKP or fragments thereof, antibodies of CYSKP, and agonists, antagonists or inhibitors of CYSKP, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by

standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the  $ED_{50}$  (the dose therapeutically effective in 50% of the population) or  $LD_{50}$  (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the  $LD_{50}/ED_{50}$  ratio. Compositions which exhibit large  
5 therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the  $ED_{50}$  with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

10 The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy.  
15 Long-acting compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1  $\mu$ g to 100,000  $\mu$ g, up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art.

20 Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

## DIAGNOSTICS

In another embodiment, antibodies which specifically bind CYSKP may be used for the  
25 diagnosis of disorders characterized by expression of CYSKP, or in assays to monitor patients being treated with CYSKP or agonists, antagonists, or inhibitors of CYSKP. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for CYSKP include methods which utilize the antibody and a label to detect CYSKP in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and  
30 may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring CYSKP, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of CYSKP expression. Normal or standard values for CYSKP expression are established by combining body fluids or cell extracts

taken from normal mammalian subjects, for example, human subjects, with antibodies to CYSKP under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of CYSKP expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values.

5 Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding CYSKP may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of CYSKP may be correlated with  
10 disease. The diagnostic assay may be used to determine absence, presence, and excess expression of CYSKP, and to monitor regulation of CYSKP levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding CYSKP or closely related molecules may be used to identify nucleic acid sequences which encode CYSKP. The specificity of the probe, whether it is made  
15 from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding CYSKP, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and may have at least 50%  
20 sequence identity to any of the CYSKP encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:35-68 or from genomic sequences including promoters, enhancers, and introns of the CYSKP gene.

Means for producing specific hybridization probes for DNAs encoding CYSKP include the cloning of polynucleotide sequences encoding CYSKP or CYSKP derivatives into vectors for the  
25 production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as <sup>32</sup>P or <sup>35</sup>S, or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

30 Polynucleotide sequences encoding CYSKP may be used for the diagnosis of disorders associated with expression of CYSKP. Examples of such disorders include, but are not limited to, a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including

adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; an autoimmune/inflammatory disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with

5 lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus,

10 systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a vesicle trafficking disorder such as cystic fibrosis, glucose-galactose malabsorption syndrome, hypercholesterolemia, diabetes mellitus, diabetes insipidus, hyper- and hypoglycemia, Grave's disease, goiter, Cushing's disease, and Addison's disease,

20 gastrointestinal disorders including ulcerative colitis, gastric and duodenal ulcers, other conditions associated with abnormal vesicle trafficking, including acquired immunodeficiency syndrome (AIDS), allergies including hay fever, asthma, and urticaria (hives), autoimmune hemolytic anemia, proliferative glomerulonephritis, inflammatory bowel disease, multiple sclerosis, myasthenia gravis, rheumatoid and osteoarthritis, scleroderma, Chediak-Higashi and Sjogren's syndromes, systemic lupus erythematosus,

25 toxic shock syndrome, traumatic tissue damage, and viral, bacterial, fungal, helminthic, and protozoal infections; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and

30 other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal

syndrome, mental retardation and other developmental disorders of the central nervous system including Down syndrome, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; a cell motility disorder such as ankylosing spondylitis, Chediak-Higashi syndrome, Duchenne and Becker muscular dystrophy, intrahepatic cholestasis, myocardial hyperplasia, cardiomyopathy, early onset peridontitis, cancers such as adenocarcinoma, ovarian carcinoma, and chronic myelogenous leukemia, and bacterial and helminthic infections; a reproductive disorder such as a disorder of prolactin production, infertility, including tubal disease, ovulatory defects, endometriosis, a disruption of the estrous cycle, a disruption of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation syndrome, an endometrial or ovarian tumor, a uterine fibroid, autoimmune disorders, ectopic pregnancy, teratogenesis, cancer of the breast, fibrocystic breast disease, galactorrhea, a disruption of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the male breast, gynecomastia, hypergonadotropic and hypogonadotropic hypogonadism, pseudohermaphroditism, azoospermia, premature ovarian failure, acrosin deficiency, delayed puberty, retrograde ejaculation and anejaculation, haemangioblastomas, cystsphaeochromocytomas, paraganglioma, cystadenomas of the epididymis, and endolymphatic sac tumours; and a muscle disorder such as myocarditis, Duchenne's muscular dystrophy, Becker's muscular dystrophy, myotonic dystrophy, central core disease, nemaline myopathy, centronuclear myopathy, lipid myopathy, mitochondrial myopathy, infectious myositis, polymyositis, dermatomyositis, inclusion body myositis, thyrotoxic myopathy, and ethanol myopathy, angina, anaphylactic shock, arrhythmias, asthma, cardiovascular shock, Cushing's syndrome, hypertension, hypoglycemia, myocardial infarction, migraine, and pheochromocytoma, and myopathies including cardiomyopathy, encephalopathy, epilepsy, Kearns-Sayre syndrome, lactic acidosis, myoclonic disorder, and ophthalmoplegia. The polynucleotide sequences encoding CYSKP may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered CYSKP expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding CYSKP may be useful in assays that



detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding CYSKP may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding CYSKP in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of CYSKP, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding CYSKP, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding CYSKP may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide encoding CYSKP, or a fragment of a polynucleotide complementary to the polynucleotide encoding CYSKP, and will be employed under optimized conditions for identification of a specific gene or condition.

Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

In a particular aspect, oligonucleotide primers derived from the polynucleotide sequences encoding CYSKP may be used to detect single nucleotide polymorphisms (SNPs). SNPs are substitutions, insertions and deletions that are a frequent cause of inherited or acquired genetic disease in humans. Methods of SNP detection include, but are not limited to, single-stranded conformation polymorphism (SSCP) and fluorescent SSCP (fSSCP) methods. In SSCP, oligonucleotide primers derived from the polynucleotide sequences encoding CYSKP are used to amplify DNA using the polymerase chain reaction (PCR). The DNA may be derived, for example, from diseased or normal tissue, biopsy samples, bodily fluids, and the like. SNPs in the DNA cause differences in the secondary and tertiary structures of PCR products in single-stranded form, and these differences are detectable using gel electrophoresis in non-denaturing gels. In fSSCP, the oligonucleotide primers are fluorescently labeled, which allows detection of the amplimers in high-throughput equipment such as DNA sequencing machines. Additionally, sequence database analysis methods, termed *in silico* SNP (isSNP), are capable of identifying polymorphisms by comparing the sequence of individual overlapping DNA fragments which assemble into a common consensus sequence. These computer-based methods filter out sequence variations due to laboratory preparation of DNA and sequencing errors using statistical models and automated analyses of DNA sequence chromatograms. In the alternative, SNPs may be detected and characterized by mass spectrometry using, for example, the high throughput MASSARRAY system (Sequenom, Inc., San Diego CA).

Methods which may also be used to quantify the expression of CYSKP include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) *J. Immunol. Methods* 159:235-244; Duplaa, C. et al. (1993) *Anal. Biochem.* 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer or polynucleotide of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as elements on a microarray. The microarray can be used in transcript imaging techniques which monitor the relative expression levels of large numbers of genes simultaneously as described below. The microarray may also be used to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, to monitor progression/regression of disease as a function of gene expression, and to develop and monitor the

activities of therapeutic agents in the treatment of disease. In particular, this information may be used to develop a pharmacogenomic profile of a patient in order to select the most appropriate and effective treatment regimen for that patient. For example, therapeutic agents which are highly effective and display the fewest side effects may be selected for a patient based on his/her pharmacogenomic profile.

5 In another embodiment, CYSKP, fragments of CYSKP, or antibodies specific for CYSKP may be used as elements on a microarray. The microarray may be used to monitor or measure protein-protein interactions, drug-target interactions, and gene expression profiles, as described above.

A particular embodiment relates to the use of the polynucleotides of the present invention to generate a transcript image of a tissue or cell type. A transcript image represents the global pattern of  
10 gene expression by a particular tissue or cell type. Global gene expression patterns are analyzed by quantifying the number of expressed genes and their relative abundance under given conditions and at a given time. (See Seilhamer et al., "Comparative Gene Transcript Analysis," U.S. Patent Number 5,840,484, expressly incorporated by reference herein.) Thus a transcript image may be generated by hybridizing the polynucleotides of the present invention or their complements to the totality of  
15 transcripts or reverse transcripts of a particular tissue or cell type. In one embodiment, the hybridization takes place in high-throughput format, wherein the polynucleotides of the present invention or their complements comprise a subset of a plurality of elements on a microarray. The resultant transcript image would provide a profile of gene activity.

Transcript images may be generated using transcripts isolated from tissues, cell lines, biopsies,  
20 or other biological samples. The transcript image may thus reflect gene expression in vivo, as in the case of a tissue or biopsy sample, or in vitro, as in the case of a cell line.

Transcript images which profile the expression of the polynucleotides of the present invention may also be used in conjunction with in vitro model systems and preclinical evaluation of pharmaceuticals, as well as toxicological testing of industrial and naturally-occurring environmental  
25 compounds. All compounds induce characteristic gene expression patterns, frequently termed molecular fingerprints or toxicant signatures, which are indicative of mechanisms of action and toxicity (Nuwaysir, E.F. et al. (1999) Mol. Carcinog. 24:153-159; Steiner, S. and N.L. Anderson (2000) Toxicol. Lett. 112-113:467-471, expressly incorporated by reference herein). If a test compound has a signature similar to that of a compound with known toxicity, it is likely to share those toxic properties.  
30 These fingerprints or signatures are most useful and refined when they contain expression information from a large number of genes and gene families. Ideally, a genome-wide measurement of expression provides the highest quality signature. Even genes whose expression is not altered by any tested compounds are important as well, as the levels of expression of these genes are used to normalize the rest of the expression data. The normalization procedure is useful for comparison of expression data

after treatment with different compounds. While the assignment of gene function to elements of a toxicant signature aids in interpretation of toxicity mechanisms, knowledge of gene function is not necessary for the statistical matching of signatures which leads to prediction of toxicity. (See, for example, Press Release 00-02 from the National Institute of Environmental Health Sciences, released  
5 February 29, 2000, available at <http://www.niehs.nih.gov/oc/news/toxchip.htm>.) Therefore, it is important and desirable in toxicological screening using toxicant signatures to include all expressed gene sequences.

In one embodiment, the toxicity of a test compound is assessed by treating a biological sample containing nucleic acids with the test compound. Nucleic acids that are expressed in the treated  
10 biological sample are hybridized with one or more probes specific to the polynucleotides of the present invention, so that transcript levels corresponding to the polynucleotides of the present invention may be quantified. The transcript levels in the treated biological sample are compared with levels in an untreated biological sample. Differences in the transcript levels between the two samples are indicative of a toxic response caused by the test compound in the treated sample.

15 Another particular embodiment relates to the use of the polypeptide sequences of the present invention to analyze the proteome of a tissue or cell type. The term proteome refers to the global pattern of protein expression in a particular tissue or cell type. Each protein component of a proteome can be subjected individually to further analysis. Proteome expression patterns, or profiles, are analyzed by quantifying the number of expressed proteins and their relative abundance under given  
20 conditions and at a given time. A profile of a cell's proteome may thus be generated by separating and analyzing the polypeptides of a particular tissue or cell type. In one embodiment, the separation is achieved using two-dimensional gel electrophoresis, in which proteins from a sample are separated by isoelectric focusing in the first dimension, and then according to molecular weight by sodium dodecyl sulfate slab gel electrophoresis in the second dimension (Steiner and Anderson, *supra*). The proteins are  
25 visualized in the gel as discrete and uniquely positioned spots, typically by staining the gel with an agent such as Coomassie Blue or silver or fluorescent stains. The optical density of each protein spot is generally proportional to the level of the protein in the sample. The optical densities of equivalently positioned protein spots from different samples, for example, from biological samples either treated or untreated with a test compound or therapeutic agent, are compared to identify any changes in protein  
30 spot density related to the treatment. The proteins in the spots are partially sequenced using, for example, standard methods employing chemical or enzymatic cleavage followed by mass spectrometry. The identity of the protein in a spot may be determined by comparing its partial sequence, preferably of at least 5 contiguous amino acid residues, to the polypeptide sequences of the present invention. In some cases, further sequence data may be obtained for definitive protein identification.

A proteomic profile may also be generated using antibodies specific for CYSKP to quantify the levels of CYSKP expression. In one embodiment, the antibodies are used as elements on a microarray, and protein expression levels are quantified by exposing the microarray to the sample and detecting the levels of protein bound to each array element (Lueking, A. et al. (1999) *Anal. Biochem.* 270:103-111; 5 Mendoze, L.G. et al. (1999) *Biotechniques* 27:778-788). Detection may be performed by a variety of methods known in the art, for example, by reacting the proteins in the sample with a thiol- or amino-reactive fluorescent compound and detecting the amount of fluorescence bound at each array element.

Toxicant signatures at the proteome level are also useful for toxicological screening, and should be analyzed in parallel with toxicant signatures at the transcript level. There is a poor correlation 10 between transcript and protein abundances for some proteins in some tissues (Anderson, N.L. and J. Seilhamer (1997) *Electrophoresis* 18:533-537), so proteome toxicant signatures may be useful in the analysis of compounds which do not significantly affect the transcript image, but which alter the proteomic profile. In addition, the analysis of transcripts in body fluids is difficult, due to rapid degradation of mRNA, so proteomic profiling may be more reliable and informative in such cases.

15 In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins that are expressed in the treated biological sample are separated so that the amount of each protein can be quantified. The amount of each protein is compared to the amount of the corresponding protein in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound 20 in the treated sample. Individual proteins are identified by sequencing the amino acid residues of the individual proteins and comparing these partial sequences to the polypeptides of the present invention.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins from the biological sample are incubated with antibodies specific to the polypeptides of the present invention. The amount of protein recognized 25 by the antibodies is quantified. The amount of protein in the treated biological sample is compared with the amount in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) *Proc. Natl. Acad. Sci. USA* 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. 30 (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.) Various types of microarrays are well known and thoroughly described in DNA Microarrays: A Practical Approach, M. Schena, ed. (1999) Oxford University Press, London, hereby expressly incorporated by reference.

In another embodiment of the invention, nucleic acid sequences encoding CYSKP may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. Either coding or noncoding sequences may be used, and in some instances, noncoding sequences may be preferable over coding sequences. For example, conservation of a coding sequence among members of a multi-gene family may potentially cause undesired cross hybridization during chromosomal mapping. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.) Once mapped, the nucleic acid sequences of the invention may be used to develop genetic linkage maps, for example, which correlate the inheritance of a disease state with the inheritance of a particular chromosome region or restriction fragment length polymorphism (RFLP). (See, for example, Lander, E.S. and D. Botstein (1986) Proc. Natl. Acad. Sci. USA 83:7353-7357.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding CYSKP on a physical map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder and thus may further positional cloning efforts.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the exact chromosomal locus is not known. This information is valuable to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the gene or genes responsible for a disease or syndrome have been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the instant invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, CYSKP, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a

solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between CYSKP and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with CYSKP, or fragments thereof, and washed. Bound CYSKP is then detected by methods well known in the art. Purified CYSKP can also be coated directly onto plates for use in the aforementioned drug screening techniques.

Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding CYSKP specifically compete with a test compound for binding CYSKP. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with CYSKP.

In additional embodiments, the nucleotide sequences which encode CYSKP may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, in particular U.S. Ser. No. 60/201,960, U.S. Ser. No. 60/202,729, U.S. Ser. No. 60/209,705, U.S. Ser. No. 60/210,149, and U.S. Ser. No. 60/213,215, are hereby expressly incorporated by reference.

## EXAMPLES

### I. Construction of cDNA Libraries

Incyte cDNAs were derived from cDNA libraries described in the LIFESEQ GOLD database (Incyte Genomics, Palo Alto CA) and shown in Table 4, column 5. Some tissues were homogenized

and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A)+ RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, supra, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUESCRIPT plasmid (Stratagene), PSPORT1 plasmid (Life Technologies), PCDNA2.1 plasmid (Invitrogen, Carlsbad CA), PBK-CMV plasmid (Stratagene), or pINCY (Incyte Genomics, Palo Alto CA), or derivatives thereof. Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 $\alpha$ , DH10B, or ElectroMAX DH10B from Life Technologies.

## II. Isolation of cDNA Clones

Plasmids obtained as described in Example I were recovered from host cells by in vivo excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.



Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) *Anal. Biochem.* 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using  
5 PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

### III. Sequencing and Analysis

Incyte cDNA recovered in plasmids as described in Example II were sequenced as follows. Sequencing reactions were processed using standard methods or high-throughput instrumentation  
10 such as the ABI CATALYST 800 (Applied Biosystems) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems).  
15 Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (Applied Biosystems) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997,  
20 supra, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example VIII.

The polynucleotide sequences derived from Incyte cDNAs were validated by removing vector, linker, and poly(A) sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The Incyte cDNA  
25 sequences or translations thereof were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and hidden Markov model (HMM)-based protein family databases such as PFAM. (HMM is a probabilistic approach which analyzes consensus primary structures of gene families. See, for example, Eddy, S.R. (1996) *Curr. Opin. Struct. Biol.* 6:361-365.) The queries were  
30 performed using programs based on BLAST, FASTA, BLIMPS, and HMMER. The Incyte cDNA sequences were assembled to produce full length polynucleotide sequences. Alternatively, GenBank cDNAs, GenBank ESTs, stitched sequences, stretched sequences, or Genscan-predicted coding sequences (see Examples IV and V) were used to extend Incyte cDNA assemblages to full length. Assembly was performed using programs based on Phred, Phrap, and Consed, and cDNA assemblages

were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length polypeptide sequences. Alternatively, a polypeptide of the invention may begin at any of the methionine residues of the full length translated polypeptide. Full length polypeptide sequences were subsequently  
5 analyzed by querying against databases such as the GenBank protein databases (genpept), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and hidden Markov model (HMM)-based protein family databases such as PFAM. Full length polynucleotide sequences are also analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments are  
10 generated using default parameters specified by the CLUSTAL algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

Table 7 summarizes the tools, programs, and algorithms used for the analysis and assembly of Incyte cDNA and full length sequences and provides applicable descriptions, references, and threshold  
15 parameters. The first column of Table 7 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score or the lower the probability value, the greater the identity  
20 between two sequences).

The programs described above for the assembly and analysis of full length polynucleotide and polypeptide sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:35-68. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies are described in Table 4, column 4.

#### 25 **IV. Identification and Editing of Coding Sequences from Genomic DNA**

Putative cytoskeleton-associated proteins were initially identified by running the Genscan gene identification program against public genomic sequence databases (e.g., gbpri and gbhtg). Genscan is a general-purpose gene identification program which analyzes genomic DNA sequences from a variety of organisms (See Burge, C. and S. Karlin (1997) J. Mol. Biol. 268:78-94, and Burge, C. and S. Karlin  
30 (1998) Curr. Opin. Struct. Biol. 8:346-354). The program concatenates predicted exons to form an assembled cDNA sequence extending from a methionine to a stop codon. The output of Genscan is a FASTA database of polynucleotide and polypeptide sequences. The maximum range of sequence for Genscan to analyze at once was set to 30 kb. To determine which of these Genscan predicted cDNA sequences encode cytoskeleton-associated proteins, the encoded polypeptides were analyzed by querying

against PFAM models for cytoskeleton-associated proteins. Potential cytoskeleton-associated proteins were also identified by homology to Incyte cDNA sequences that had been annotated as cytoskeleton-associated proteins. These selected Genscan-predicted sequences were then compared by BLAST analysis to the genpept and gbpri public databases. Where necessary, the Genscan-predicted sequences were then edited by comparison to the top BLAST hit from genpept to correct errors in the sequence predicted by Genscan, such as extra or omitted exons. BLAST analysis was also used to find any Incyte cDNA or public cDNA coverage of the Genscan-predicted sequences, thus providing evidence for transcription. When Incyte cDNA coverage was available, this information was used to correct or confirm the Genscan predicted sequence. Full length polynucleotide sequences were obtained by assembling Genscan-predicted coding sequences with Incyte cDNA sequences and/or public cDNA sequences using the assembly process described in Example III. Alternatively, full length polynucleotide sequences were derived entirely from edited or unedited Genscan-predicted coding sequences.

#### **V. Assembly of Genomic Sequence Data with cDNA Sequence Data**

##### **"Stitched" Sequences**

Partial cDNA sequences were extended with exons predicted by the Genscan gene identification program described in Example IV. Partial cDNAs assembled as described in Example III were mapped to genomic DNA and parsed into clusters containing related cDNAs and Genscan exon predictions from one or more genomic sequences. Each cluster was analyzed using an algorithm based on graph theory and dynamic programming to integrate cDNA and genomic information, generating possible splice variants that were subsequently confirmed, edited, or extended to create a full length sequence. Sequence intervals in which the entire length of the interval was present on more than one sequence in the cluster were identified, and intervals thus identified were considered to be equivalent by transitivity. For example, if an interval was present on a cDNA and two genomic sequences, then all three intervals were considered to be equivalent. This process allows unrelated but consecutive genomic sequences to be brought together, bridged by cDNA sequence. Intervals thus identified were then "stitched" together by the stitching algorithm in the order that they appear along their parent sequences to generate the longest possible sequence, as well as sequence variants. Linkages between intervals which proceed along one type of parent sequence (cDNA to cDNA or genomic sequence to genomic sequence) were given preference over linkages which change parent type (cDNA to genomic sequence). The resultant stitched sequences were translated and compared by BLAST analysis to the genpept and gbpri public databases. Incorrect exons predicted by Genscan were corrected by comparison to the top BLAST hit from genpept. Sequences were further extended with additional cDNA sequences, or by inspection of genomic DNA, when necessary.

**“Stretched” Sequences**

Partial DNA sequences were extended to full length with an algorithm based on BLAST analysis. First, partial cDNAs assembled as described in Example III were queried against public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases using the BLAST program. The nearest GenBank protein homolog was then compared by BLAST analysis to either Incyte cDNA sequences or GenScan exon predicted sequences described in Example IV. A chimeric protein was generated by using the resultant high-scoring segment pairs (HSPs) to map the translated sequences onto the GenBank protein homolog. Insertions or deletions may occur in the chimeric protein with respect to the original GenBank protein homolog. The GenBank protein homolog, the chimeric protein, or both were used as probes to search for homologous genomic sequences from the public human genome databases. Partial DNA sequences were therefore “stretched” or extended by the addition of homologous genomic sequences. The resultant stretched sequences were examined to determine whether it contained a complete gene.

**VI. Chromosomal Mapping of CYSKP Encoding Polynucleotides**

The sequences which were used to assemble SEQ ID NO:35-68 were compared with sequences from the Incyte LIFESEQ database and public domain databases using BLAST and other implementations of the Smith-Waterman algorithm. Sequences from these databases that matched SEQ ID NO:35-68 were assembled into clusters of contiguous and overlapping sequences using assembly algorithms such as Phrap (Table 7). Radiation hybrid and genetic mapping data available from public resources such as the Stanford Human Genome Center (SHGC), Whitehead Institute for Genome Research (WIGR), and Généthon were used to determine if any of the clustered sequences had been previously mapped. Inclusion of a mapped sequence in a cluster resulted in the assignment of all sequences of that cluster, including its particular SEQ ID NO., to that map location.

Map locations are represented by ranges, or intervals, of human chromosomes. The map position of an interval, in centiMorgans, is measured relative to the terminus of the chromosome’s p-arm. (The centiMorgan (cM) is a unit of measurement based on recombination frequencies between chromosomal markers. On average, 1 cM is roughly equivalent to 1 megabase (Mb) of DNA in humans, although this can vary widely due to hot and cold spots of recombination.) The cM distances are based on genetic markers mapped by Généthon which provide boundaries for radiation hybrid markers whose sequences were included in each of the clusters. Human genome maps and other resources available to the public, such as the NCBI “GeneMap’99” World Wide Web site (<http://www.ncbi.nlm.nih.gov/genemap/>), can be employed to determine if previously identified disease genes map within or in proximity to the intervals indicated above.

In this manner, SEQ ID NO:44 was mapped to chromosome 17 within the interval from

62.90 to 64.20 centiMorgans, SEQ ID NO:49 was mapped to chromosome 14 within the interval from 73.70 to 76.40 centiMorgans, SEQ ID NO:50 was mapped to chromosome 8 within the interval from 25.80 to 40.30 centiMorgans, SEQ ID NO:54 was mapped to chromosome 1 within the interval from 117.6 to 132.4 centiMorgans, SEQ ID NO:64 was mapped to chromosome 4 within the interval from 56.7 to 60.5 centiMorgans, and SEQ ID NO:65 was mapped to chromosome 5 within the interval from 141.40 to 142.60 centiMorgans.

## VII. Analysis of Polynucleotide Expression

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, supra, ch. 7; Ausubel (1995) supra, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in cDNA databases such as GenBank or LIFESEQ (Incyte Genomics). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\text{BLAST Score} \times \text{Percent Identity}}{5 \times \text{minimum \{length(Seq. 1), length(Seq. 2)\}}}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. The product score is a normalized value between 0 and 100, and is calculated as follows: the BLAST score is multiplied by the percent nucleotide identity and the product is divided by (5 times the length of the shorter of the two sequences). The BLAST score is calculated by assigning a score of +5 for every base that matches in a high-scoring segment pair (HSP), and -4 for every mismatch. Two sequences may share more than one HSP (separated by gaps). If there is more than one HSP, then the pair with the highest BLAST score is used to calculate the product score. The product score represents a balance between fractional overlap and quality in a BLAST alignment. For example, a product score of 100 is produced only for 100% identity over the entire length of the shorter of the two sequences being compared. A product score of 70 is produced either by 100% identity and 70% overlap at one end, or by 88% identity and 100% overlap at the other. A product score of 50 is produced either by 100% identity and 50% overlap at one end, or 79% identity and 100% overlap.

Alternatively, polynucleotide sequences encoding CYSKP are analyzed with respect to the tissue sources from which they were derived. For example, some full length sequences are assembled,

at least in part, with overlapping Incyte cDNA sequences (see Example III). Each cDNA sequence is derived from a cDNA library constructed from a human tissue. Each human tissue is classified into one of the following organ/tissue categories: cardiovascular system; connective tissue; digestive system; embryonic structures; endocrine system; exocrine glands; genitalia, female; genitalia, male; germ cells; hemic and immune system; liver; musculoskeletal system; nervous system; pancreas; respiratory system; sense organs; skin; stomatognathic system; unclassified/mixed; or urinary tract. The number of libraries in each category is counted and divided by the total number of libraries across all categories. Similarly, each human tissue is classified into one of the following disease/condition categories: cancer, cell line, developmental, inflammation, neurological, trauma, cardiovascular, pooled, and other, and the number of libraries in each category is counted and divided by the total number of libraries across all categories. The resulting percentages reflect the tissue- and disease-specific expression of cDNA encoding CYSKP. cDNA sequences and cDNA library/tissue information are found in the LIFESEQ GOLD database (Incyte Genomics, Palo Alto CA).

#### **VIII. Extension of CYSKP Encoding Polynucleotides**

Full length polynucleotide sequences were also produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer was synthesized to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing  $Mg^{2+}$ ,  $(NH_4)_2SO_4$ , and 2-mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100  $\mu$ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5  $\mu$ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II  
5 (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5  $\mu$ l to 10  $\mu$ l aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and  
10 sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs,  
15 and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, and individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1:  
20 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC  
25 DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems).

In like manner, full length polynucleotide sequences are verified using the above procedure or are used to obtain 5' regulatory sequences using the above procedure along with oligonucleotides designed for such extension, and an appropriate genomic library.

#### 30 IX. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:35-68 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National

Biosciences) and labeled by combining 50 pmol of each oligomer, 250  $\mu$ Ci of [ $\gamma$ - $^{32}$ P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing  $10^7$  counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and compared.

#### **X. Microarrays**

The linkage or synthesis of array elements upon a microarray can be achieved utilizing photolithography, piezoelectric printing (ink-jet printing, See, e.g., Baldeschweiler, supra), mechanical microspotting technologies, and derivatives thereof. The substrate in each of the aforementioned technologies should be uniform and solid with a non-porous surface (Skena (1999), supra). Suggested substrates include silicon, silica, glass slides, glass chips, and silicon wafers. Alternatively, a procedure analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced using available methods and machines well known to those of ordinary skill in the art and may contain any appropriate number of elements. (See, e.g., Skena, M. et al. (1995) Science 270:467-470; Shalon, D. et al. (1996) Genome Res. 6:639-645; Marshall, A. and J. Hodgson (1998) Nat. Biotechnol. 16:27-31.)

Full length cDNAs, Expressed Sequence Tags (ESTs), or fragments or oligomers thereof may comprise the elements of the microarray. Fragments or oligomers suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). The array elements are hybridized with polynucleotides in a biological sample. The polynucleotides in the biological sample are conjugated to a fluorescent label or other molecular tag for ease of detection. After hybridization, nonhybridized nucleotides from the biological sample are removed, and a fluorescence scanner is used to detect hybridization at each array element. Alternatively, laser desorption and mass spectrometry may be used for detection of hybridization. The degree of complementarity and the relative abundance of each polynucleotide which hybridizes to an element on



the microarray may be assessed. In one embodiment, microarray preparation and usage is described in detail below.

#### **Tissue or Cell Sample Preparation**

Total RNA is isolated from tissue samples using the guanidinium thiocyanate method and poly(A)<sup>+</sup> RNA is purified using the oligo-(dT) cellulose method. Each poly(A)<sup>+</sup> RNA sample is  
5 reverse transcribed using MMLV reverse-transcriptase, 0.05 pg/μl oligo-(dT) primer (21mer), 1X first strand buffer, 0.03 units/μl RNase inhibitor, 500 μM dATP, 500 μM dGTP, 500 μM dTTP, 40 μM dCTP, 40 μM dCTP-Cy3 (BDS) or dCTP-Cy5 (Amersham Pharmacia Biotech). The reverse transcription reaction is performed in a 25 ml volume containing 200 ng poly(A)<sup>+</sup> RNA with  
10 GEMBRIGHT kits (Incyte). Specific control poly(A)<sup>+</sup> RNAs are synthesized by in vitro transcription from non-coding yeast genomic DNA. After incubation at 37° C for 2 hr, each reaction sample (one with Cy3 and another with Cy5 labeling) is treated with 2.5 ml of 0.5M sodium hydroxide and incubated for 20 minutes at 85° C to stop the reaction and degrade the RNA. Samples are purified using two successive CHROMA SPIN 30 gel filtration spin columns (CLONTECH Laboratories, Inc.  
15 (CLONTECH), Palo Alto CA) and after combining, both reaction samples are ethanol precipitated using 1 ml of glycogen (1 mg/ml), 60 ml sodium acetate, and 300 ml of 100% ethanol. The sample is then dried to completion using a SpeedVAC (Savant Instruments Inc., Holbrook NY) and resuspended in 14 μl 5X SSC/0.2% SDS.

#### **Microarray Preparation**

Sequences of the present invention are used to generate array elements. Each array element is  
20 amplified from bacterial cells containing vectors with cloned cDNA inserts. PCR amplification uses primers complementary to the vector sequences flanking the cDNA insert. Array elements are amplified in thirty cycles of PCR from an initial quantity of 1-2 ng to a final quantity greater than 5 μg. Amplified array elements are then purified using SEPHACRYL-400 (Amersham Pharmacia  
25 Biotech).

Purified array elements are immobilized on polymer-coated glass slides. Glass microscope slides (Corning) are cleaned by ultrasound in 0.1% SDS and acetone, with extensive distilled water washes between and after treatments. Glass slides are etched in 4% hydrofluoric acid (VWR Scientific Products Corporation (VWR), West Chester PA), washed extensively in distilled water, and  
30 coated with 0.05% aminopropyl silane (Sigma) in 95% ethanol. Coated slides are cured in a 110°C oven.

Array elements are applied to the coated glass substrate using a procedure described in US Patent No. 5,807,522, incorporated herein by reference. 1 μl of the array element DNA, at an average concentration of 100 ng/μl, is loaded into the open capillary printing element by a high-speed robotic  
35 apparatus. The apparatus then deposits about 5 nl of array element sample per slide.

Microarrays are UV-crosslinked using a STRATALINKER UV-crosslinker (Stratagene). Microarrays are washed at room temperature once in 0.2% SDS and three times in distilled water. Non-specific binding sites are blocked by incubation of microarrays in 0.2% casein in phosphate buffered saline (PBS) (Tropix, Inc., Bedford MA) for 30 minutes at 60° C followed by washes in  
5 0.2% SDS and distilled water as before.

#### **Hybridization**

Hybridization reactions contain 9 µl of sample mixture consisting of 0.2 µg each of Cy3 and Cy5 labeled cDNA synthesis products in 5X SSC, 0.2% SDS hybridization buffer. The sample mixture is heated to 65° C for 5 minutes and is aliquoted onto the microarray surface and covered with  
10 an 1.8 cm<sup>2</sup> coverslip. The arrays are transferred to a waterproof chamber having a cavity just slightly larger than a microscope slide. The chamber is kept at 100% humidity internally by the addition of 140 µl of 5X SSC in a corner of the chamber. The chamber containing the arrays is incubated for about 6.5 hours at 60° C. The arrays are washed for 10 min at 45° C in a first wash buffer (1X SSC, 0.1% SDS), three times for 10 minutes each at 45° C in a second wash buffer (0.1X SSC), and dried.

#### **Detection**

Reporter-labeled hybridization complexes are detected with a microscope equipped with an Innova 70 mixed gas 10 W laser (Coherent, Inc., Santa Clara CA) capable of generating spectral lines at 488 nm for excitation of Cy3 and at 632 nm for excitation of Cy5. The excitation laser light is focused on the array using a 20X microscope objective (Nikon, Inc., Melville NY). The slide  
20 containing the array is placed on a computer-controlled X-Y stage on the microscope and raster-scanned past the objective. The 1.8 cm x 1.8 cm array used in the present example is scanned with a resolution of 20 micrometers.

In two separate scans, a mixed gas multiline laser excites the two fluorophores sequentially. Emitted light is split, based on wavelength, into two photomultiplier tube detectors (PMT R1477,  
25 Hamamatsu Photonics Systems, Bridgewater NJ) corresponding to the two fluorophores. Appropriate filters positioned between the array and the photomultiplier tubes are used to filter the signals. The emission maxima of the fluorophores used are 565 nm for Cy3 and 650 nm for Cy5. Each array is typically scanned twice, one scan per fluorophore using the appropriate filters at the laser source, although the apparatus is capable of recording the spectra from both fluorophores simultaneously.

The sensitivity of the scans is typically calibrated using the signal intensity generated by a cDNA control species added to the sample mixture at a known concentration. A specific location on the array contains a complementary DNA sequence, allowing the intensity of the signal at that location to be correlated with a weight ratio of hybridizing species of 1:100,000. When two samples from different sources (e.g., representing test and control cells), each labeled with a different  
35 fluorophore, are hybridized to a single array for the purpose of identifying genes that are differentially

expressed, the calibration is done by labeling samples of the calibrating cDNA with the two fluorophores and adding identical amounts of each to the hybridization mixture.

The output of the photomultiplier tube is digitized using a 12-bit RTI-835H analog-to-digital (A/D) conversion board (Analog Devices, Inc., Norwood MA) installed in an IBM-compatible PC  
5 computer. The digitized data are displayed as an image where the signal intensity is mapped using a linear 20-color transformation to a pseudocolor scale ranging from blue (low signal) to red (high signal). The data is also analyzed quantitatively. Where two different fluorophores are excited and measured simultaneously, the data are first corrected for optical crosstalk (due to overlapping emission spectra) between the fluorophores using each fluorophore's emission spectrum.

10 A grid is superimposed over the fluorescence signal image such that the signal from each spot is centered in each element of the grid. The fluorescence signal within each element is then integrated to obtain a numerical value corresponding to the average intensity of the signal. The software used for signal analysis is the GEMTOOLS gene expression analysis program (Incyte).

#### **XI. Complementary Polynucleotides**

15 Sequences complementary to the CYSKP-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring CYSKP. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of CYSKP. To inhibit transcription, a  
20 complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the CYSKP-encoding transcript.

#### **XII. Expression of CYSKP**

Expression and purification of CYSKP is achieved using bacterial or virus-based expression  
25 systems. For expression of CYSKP in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac (tac)* hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3).

30 Antibiotic resistant bacteria express CYSKP upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of CYSKP in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding CYSKP by either homologous recombination or bacterial-mediated

transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, CYSKP is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from CYSKP at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified CYSKP obtained by these methods can be used directly in the assays shown in Examples XVI and XVII, where applicable.

### **XIII. Functional Assays**

CYSKP function is assessed by expressing the sequences encoding CYSKP at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include PCMV SPORT (Life Technologies) and PCR3.1 (Invitrogen, Carlsbad CA), both of which contain the cytomegalovirus promoter. 5-10  $\mu$ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2  $\mu$ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in

cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of CYSKP on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding CYSKP and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding CYSKP and other genes of interest can be analyzed by northern analysis or microarray techniques.

#### 15 **XIV. Production of CYSKP Specific Antibodies**

CYSKP substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the CYSKP amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, supra, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Applied Biosystems) using Fmoc chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, supra.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-CYSKP activity by, for example, binding the peptide or CYSKP to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

#### 30 **XV. Purification of Naturally Occurring CYSKP Using Specific Antibodies**

Naturally occurring or recombinant CYSKP is substantially purified by immunoaffinity chromatography using antibodies specific for CYSKP. An immunoaffinity column is constructed by covalently coupling anti-CYSKP antibody to an activated chromatographic resin, such as

CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing CYSKP are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of CYSKP (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/CYSKP binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and CYSKP is collected.

#### **XVI. Identification of Molecules Which Interact with CYSKP**

CYSKP, or biologically active fragments thereof, are labeled with <sup>125</sup>I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) Biochem. J. 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled CYSKP, washed, and any wells with labeled CYSKP complex are assayed. Data obtained using different concentrations of CYSKP are used to calculate values for the number, affinity, and association of CYSKP with the candidate molecules.

Alternatively, molecules interacting with CYSKP are analyzed using the yeast two-hybrid system as described in Fields, S. and O. Song (1989) Nature 340:245-246, or using commercially available kits based on the two-hybrid system, such as the MATCHMAKER system (Clontech).

CYSKP may also be used in the PATHCALLING process (CuraGen Corp., New Haven CT) which employs the yeast two-hybrid system in a high-throughput manner to determine all interactions between the proteins encoded by two large libraries of genes (Nandabalan, K. et al. (2000) U.S. Patent No. 6,057,101).

#### **XVII. Demonstration of CYSKP Activity**

A microtubule motility assay for CYSKP measures motor protein activity. In this assay, recombinant CYSKP is immobilized onto a glass slide or similar substrate. Taxol-stabilized bovine brain microtubules (commercially available) in a solution containing ATP and cytosolic extract are perfused onto the slide. Movement of microtubules as driven by CYSKP motor activity can be visualized and quantified using video-enhanced light microscopy and image analysis techniques. CYSKP activity is directly proportional to the frequency and velocity of microtubule movement.

Alternatively, an assay for CYSKP measures the formation of protein filaments in vitro. A solution of CYSKP at a concentration greater than the "critical concentration" for polymer assembly is applied to carbon-coated grids. Appropriate nucleation sites may be supplied in the solution. The grids are negative stained with 0.7% (w/v) aqueous uranyl acetate and examined by electron microscopy. The appearance of filaments of approximately 25 nm (microtubules), 8 nm (actin), or 10 nm (intermediate filaments) is a demonstration of protein activity.

Alternatively, an assay for CYSKP measures the binding affinity of CYSKP for actin as described by Hammell, R.L. and Hitchcock-DeGregori, S.E. (1997, J. Biol. Chem. 272:22409-22416). CYSKP and actin are prepared from in vitro recombinant cDNA expression systems and the N-terminus of CYSKP is acetylated using methods well known in the art. Binding of N-terminal acetyl-  
5 CYSKP to actin is measured by cosedimentation at 25 °C in a Beckman model TL-100 centrifuge as described. The bound and free CYSKP are determined by quantitative densitometry of SDS-polyacrylamide gels stained with Coomassie Blue. Apparent binding constants ( $K_{app}$ ) and Hill coefficients (H) are determined by using methods well known in the art to fit the data to the equation as described by Hammell and Hitchcock-DeGregori (1997, supra). The CYSKP:actin ratio, determined  
10 using densitometry, is normalized. Hammell and Hitchcock-DeGregori (1997, supra) have shown that saturation of binding corresponds to a CYSKP:actin molar ratio of 0.14, a stoichiometry of 1 CYSKP:7 actin. The binding of CYSKP to actin is proportional to the CYSKP activity.

Alternatively, CYSKP activity is measured as ability to bind to microtubules. Microtubules are purified from adult rat brain by reversible assembly (Vallee, R. B. (1982) Methods Enzymol.  
15 134:89-104) or the taxol method (Vallee, R. B. (1982) J. Cell Biol. 92:435-442) using PEM buffer (100 mM PIPES, pH 6.6, 1mM EGTA, 1mM MgSO<sub>4</sub>). To separate the MAPs from tubulin, the pellets from twice-cycled microtubules are resuspended in PEM buffer and applied to a 0.1 M MgSO<sub>4</sub>-saturated phosphocellulose column as described by Sloboda, R. D. and Rosenbaum, J. L. ((1982) Methods Enzymol. 85:409-416). The fractions containing protein are applied to a second  
20 phosphocellulose column. In a total volume of 100 ml, 20 ml of CYSKP (250 mg/ml) is added to 80 ml of whole microtubules (450 mg/ml) or tubulin (300 mg/ml) and incubated at 37 °C for 10 minutes in the presence of 1 mM GTP and 50 mM taxol. The suspension is centrifuged, the supernatant is removed, and the microtubule pellet is resuspended to the original reaction volume in PEM buffer. To assess the partitioning of CYSKP between the supernatant and pellet fractions, equal amounts of supernatant and  
25 resuspended pellet are placed in SDS sample buffer and assayed on a 5-20% gradient SDS polyacrylamide gel stained with Coomassie Brilliant Blue. The amount of CYSKP in the pellet fraction is proportional to the binding of CYSKP to microtubules.

Alternatively, CYSKP activity is associated with its ability to form protein-protein complexes and is measured by its ability to regulate growth characteristics of NIH3T3 mouse fibroblast cells. A  
30 cDNA encoding CYSKP is subcloned into an appropriate eukaryotic expression vector. This vector is transfected into NIH3T3 cells using methods known in the art. Transfected cells are compared with non-transfected cells for the following quantifiable properties: growth in culture to high density, reduced attachment of cells to the substrate, altered cell morphology, and ability to induce tumors when injected into immunodeficient mice. The activity of CYSKP is proportional to the extent of increased growth or

frequency of altered cell morphology in NIH3T3 cells transfected with CYSKP.

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention.

- 5 Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.



Table 1

Incyte Project ID	Polypeptide SEQ ID NO:	Incyte Polypeptide ID	Poly-nucleotide SEQ ID NO:	Incyte Polynucleotide ID
1889577	1	1889577CD1	35	1889577CB1
2427982	2	2427982CD1	36	2427982CB1
2470833	3	2470833CD1	37	2470833CB1
2080579	4	2080579CD1	38	2080579CB1
2156553	5	2156553CD1	39	2156553CB1
2182855	6	2182855CD1	40	2182855CB1
2242106	7	2242106CD1	41	2242106CB1
2726877	8	2726877CD1	42	2726877CB1
2738233	9	2738233CD1	43	2738233CB1
1833116	10	1833116CD1	44	1833116CB1
001799	11	001799CD1	45	001799CB1
119814	12	119814CD1	46	119814CB1
1295420	13	1295420CD1	47	1295420CB1
1309364	14	1309364CD1	48	1309364CB1
1315267	15	1315267CD1	49	1315267CB1
1403289	16	1403289CD1	50	1403289CB1
1607607	17	1607607CD1	51	1607607CB1
1660025	18	1660025CD1	52	1660025CB1
1796836	19	1796836CD1	53	1796836CB1
2880670	20	2880670CD1	54	2880670CB1
2913976	21	2913976CD1	55	2913976CB1
3092084	22	3092084CD1	56	3092084CB1
3882482	23	3882482CD1	57	3882482CB1
4933451	24	4933451CD1	58	4933451CB1
5043904	25	5043904CD1	59	5043904CB1
5202390	26	5202390CD1	60	5202390CB1
5526375	27	5526375CD1	61	5526375CB1
5677408	28	5677408CD1	62	5677408CB1
5982278	29	5982278CD1	63	5982278CB1
6437362	30	6437362CD1	64	6437362CB1
4173970	31	4173970CD1	65	4173970CB1
2772751	32	2772751CD1	66	2772751CB1
2793768	33	2793768CD1	67	2793768CB1
3035248	34	3035248CD1	68	3035248CB1

Table 2

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO:	Probability score	GenBank Homolog
1	1889577CD1	g3347848	0.00E+00	kinesin light chain 2 [Mus musculus]
2	2427982CD1	g2760161	3.00E-64	outer arm dynein light chain 2 [Anthocidaris crassispina]
3	2470833CD1	g11094032	1.00E-147	[Mus musculus] (AF312712) gamma-parvin
4	2080579CD1	g11036542	0	[Homo sapiens] (AF237772) gamma-parvin
5	2156553CD1	g6141549	2.50E-101	JNK/SAPK-associated protein-1 (JIP-1) scaffold protein [Mus musculus] (Meyer, D. et al. (1999) J. Biol. Chem. 574:35113-35118)
6	2182855CD1	g5419859	2.00E-170	hypothetical protein similar to tubulin- tyrosine ligase [Homo sapiens] (Lafanechere, L. et al. (1998) J. Cell Sci. 11:171-181)
7	2242106CD1	g2276319	0	axonemal dynein heavy chain [Homo sapiens]
8	2726877CD1	g3834443	2.00E-13	[Drosophila melanogaster] cytoplasmic dynein intermediate chain isoform DIC5b
9	2738233CD1	g18156	1.20E-10	70kD dynein intermediate chain [Chlamydomonas reinhardtii]
10	1833116CD1	g4778	1.30E-12	Uso1 protein [Saccharomyces cerevisiae] (Nakajima, H. et al. (1991) J. Cell Biol. 113:245-260)
11	1799CD1	g4185884	7.70E-33	Groovin (Kakapo) [Drosophila melanogaster] (Strumpf, D. and T. Volk (1998) J. Cell Biol. 143:1259-1270)
12	119814CD1	g10880797	0	[Mus musculus] Syne-1A
13	1295420CD1	g12082089	0	[Homo sapiens] hARPX
14	1309364CD1	g12082091	0	[Gallus gallus] gARPX
		g3283070	1.70E-07	p80 katanin [Xenopus laevis] (McNally, F.J., Thomas, S. (1998) Mol. Biol. Cell 9:1847- 1861)
		g3005599	5.00E-09	[Homo sapiens] (AF052432) katanin p80 subunit
		g3243131	4.40E-18	titin [Drosophila melanogaster] (Machado, C. et al. (1998) J. Cell Biol. 141:321-333)
		g5870837	1.00E-113	[Homo sapiens] titin-like protein
		g180622	5.60E-37	cytoplasmic linker protein-170 alpha-2 [Homo sapiens] (Pierre, P. et al. (1992) Cell 70:887-900)
		g12667401	0	[Homo sapiens] NUF2R
		g12667403	0	[Mus musculus] NUF2R

Table 2 (cont.)

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO:	Probability score	GenBank Homolog
15	1315267CD1	g53996	8.00E-74	Tcp-10 (transmission control protein) [Mus musculus] (Davies, P. et al. (1991) Mamm. Genome 1:235-241)
16	1403289CD1	g5733814	4.60E-196	angiotensin II AT2 receptor-interacting protein (Bedecs, K. et al. (1997) Biochem. J. 325:449-454)
17	1607607CD1	g3158498	1.60E-19	Contains similarity to Pfam domain: PF00628 (PHD finger) (Aasland, R. et al. (1995) Trends Biochem. Sci. 20:56-59)
18	1660025CD1	g3253105	9.80E-20	[Caenorhabditis elegans] strong similarity to the SNF2/RAD54 family of helicases (Eisen, J. et al. (1995) Nucleic Acids Res. 23:2715-2723)
19	1796836CD1	g414111	7.20E-14	class II INCENP protein (inner centromere protein) [Gallus gallus] (Mackay, A. et al. (1993) J. Cell Biol. 123:373-385)
20	2880670CD1	g1813638	6.90E-16	PF20 [Chlamydomonas reinhardtii] (Smith, E. and P. Lefebvre (1997) Mol. Biol. Cell 8:455-467)
21	2913976CD1	g63898	3.10E-56	Zyxin [Gallus gallus] (Sadler, I. et al. (1992) J. Cell Biol. 119:1573-1587)
22	3092084CD1	g1154645	2.30E-10	head-elevated expression in 0.9 kb [Drosophila melanogaster] (Yang, M.Y. et al. (2000) Genetics 154:285-297)
23	3882482CD1	g5825592	7.60E-171	katanin p60 [Xenopus laevis]
24	4933451CD1	g684936	3.20E-30	peptide with resemblance to the actin family [Homo sapiens]
25	5043904CD1	g2832237	2.10E-06	cep250 centrosome associated protein [Homo sapiens] (Mack, G.J. et al. (1998) Arthritis Rheum. 41:551-558)
26	5202390CD1	g6572155	2.90E-21	[Homo sapiens] dJ1014D13.2 (novel protein similar to ACTN3 (actinin, alpha 3))
27	5526375CD1	g2443272	2.80E-77	motor domain of KIF12 [Mus musculus] (Nakagawa, T. et al. (1997) Proc. Natl. Acad. Sci. USA 94:9654-9659)
28	5677408CD1	g6651427	2.20E-05	dynein light intermediate chain 1 (LIC-2) [Rattus norvegicus] (Hughes, S.M. et al. (1995) J. Cell Sci. 108:17-24)
29	5982278CD1	g6006743	0	mitotic kinesin-like protein 1 [Danio rerio]
		g6723675	0	[Homo sapiens] mitotic kinase-like protein-1

Table 2 (cont.)

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO:	Probability score	GenBank Homolog
30	6437362CD1	g4929268	2.00E-38	LOMP protein (LIM and PDZ domain protein) [Homo sapiens]
31	4173970CD1	g3879121	1.10E-146	predicted using Genefinder; Similarity to Mouse ankyrin [Caenorhabditis elegans]
32	2772751CD1	g4545313	7.00E-103	[Mus musculus] prominin-like protein (Corbeil, D. et al. (2000) J. Biol. Chem. 275:5512-5520)
33	2793768CD1	g485107	2.70E-85	weakly similar to ANK repeat region of Fowlpox virus BamHI-orf7
34	3035248CD1	g1333846	8.20E-45	intermediate filament associated protein [Cricetulus griseus] (Skalli, O. et al. (1994) J. Cell Biol. 125:159-170)

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
1	1889577CD1	622	T30 S90 T451 S499 S507 S539 T568 S615 Y345 Y431 S428 S557 S581 S619 S13 S151 T163 S232 T470 S507 S519 S521 T568 S589 S610	N449 N587	Kinesin light chain repeat BL01160:V88-S141, G191-P237, D238-A266, L267-C305, A308-R348, R349-C375, Q379-E420, E433-K480, L12-P50 KINESIN LIGHT CHAIN SIGNATURE PR00381:A97-A114, G191-S210, R213-T231, H278-R295, D322-E342, R357-K378 KINESIN LIGHT CHAIN PROTEIN KLC MOTOR MICROTUBULES COILED COIL REPEAT PD012762:L12-Q174 KINESIN LIGHT CHAIN REPEAT DM01439 A41539 1-234:M1-L220 signal_cleavage:M1-P50 Kinesin light chain repeat kinesin2:Q223-N264, D265-K306 Kinesin light chain repeat Kinesin_Light:Q223-N264 D265-K306 Leucine Rich Repeat LRR:N49-K70, N 71-G92, T94-K115, K116-P140 COSMID C06A8 PROTEIN T09A5.9 CHROMOSOME III LEUCINEREPEAT REPEAT PD035408:S54-T179 Leucine Rich Repeat signature PR00019:L69-I82	BLIMPS-BLOCKS
2	2427982CD1	190	T6 T94 T6 T167		Calponin homology (CH) domain CH:N210-T317	HMMER-PFAM
3	2470833CD1	331	S37 S67 T188 S267 S293 T36 S37 S101 S169 S176 T188 T305 T317	N55 N114 N274	PROTEIN CHROMOSOME TUBULIN TYROSINE LIGASE TTL C55A6.2 ZK1128.6 III PD008766:G63-V285	BLAST-PRODOM
4	2080579CD1	239	T92 S148 T174 S191 S26			MOTIFS
5	2156553CD1	488	S237 T370 T402 T121 T226 S428	N167 N168		MOTIFS

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
6	2182855CD1	1190	S26 S197 S207 S312 T354 S356 S492 S509 S519 S593 S686 T902 S920 S976 T992 Y811 Y955 T45 T98 S149 T163 T233 T350 S406 T446 S468 S524 S896 S976	N20 N23 N156 N308 N433 N548 N635 N777	PROTEIN DYNEIN CHAIN MOTOR MICROTUBULES ATPBINDING HEPTAD REPEAT PATTERN HEAVY PD003982:S920-V1190 do DYNEIN; HEAVY; CILIARY; CYTOSOLIC DM04585 P39057 2948-4465:L14-V1190 G_Beta_Repeats:L130-N144	BLAST-PRODOM BLAST-DOMO MOTIFS
7	2242106CD1	270	S15 S56 S168 S189 S203 T240 S77 T213		signal_cleavage:M1-T25 transmem_domain:V29-L53 WD domain, G-beta repeat WD40:A116-S155, T207-Q245	SPScan HMMER HMMER-PFAM
8	2726877CD1	647	S38 T173 T184 S322 S442 T483 T503 S510 S589 T14 S122 T134 T189 T408 S447 S461 S472 S510 S579 S593	N99 N120 N316 N480 N508 N644	PROTEIN COILED COIL CHAIN MYOSIN REPEAT HEAVY ATPBINDING FILAMENT HEPTAD PD000002:Q409-E619	BLAST-PRODOM
9	2738233CD1	1086	T386 S12 S32 T86 S142 T251 S298 T343 T404 T414 T421 S427 S512 S559 S594 S618 T651 S671 T748 T799 S825 S870 S900 S954 S962 S963 Y146 S58 S75 S185 T286 S307 S366 T404 S512 S556 S559 S658 S675 S977 T987	N182 N359 N545	Spectrin repeat:R2-E66, N69-E171, V174-E285, R288-H394, G397-R501, T699-Q726, Q729-D836 Spectrin repeat proteins PF00435:W155-K170	HMMER-PFAM BLIMPS-PFAM
10	1833116CD1	396	S104 S108 T341 T343 S367 T378 S388 T34 S163 S189 T243 S258	N21 N101	P59-K245, Q266-D395 Actin T3-L37, Q73-A127, R137-R191, I289-T343, D346-D395 Actins V333-K393 Actins signatures L4-I289 ACTINS AND ACTIN-RELATED PROTEINS Q266-K393 ACTIN	HMMER PFAM BLIMPS_BLOCKS PROFILESCAN BLAST DOMO BLAST PRODOM

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
11	001799CD1	304	Y121 Y183 T33 S43 T58 S137 S254 S30 S89 S176 S131 S229 S255	N31 N52 N124	W172-D189 Aldehyde dehydrogenases motif V147-W249 P_value 5.9e-07 KATANIN P80 centrosome-binding subunit	BLIMPS_BLOCKS BLAST_PRODUM
12	119814CD1	201	T139 Y24 S4 S60 S68 T96 T106 T144	N148	M1-A23 signal cleavage G66-T125, S10-A28 Immunoglobulin domain	SPSCAN HMMER_PFAM
13	1295420CD1	547	T399T2 T58 T77 T192 T260 S491 T518 T154 T309 T374 T377 T386 T454 T515 T518	N190	G314-F345 G436-F467 Cap_Gly G314-S356, G436-P478 CAP-Gly domain T117-R158, T160-S191, N197-R229 Ank repeat G321-F345 CAP-Gly domain proteins G436-F476 MICROTUBULES CYTOSKELETON COILED COIL E417-P492, L294-K362 CAP(cytoskeleton-associated protein)-GLY DOMAIN Q177-K418 COILED COIL MYOSIN REPEAT	MOTIFS HMMER_PFAM HMMER_PFAM BLIMPS_BLOCKS BLAST_PRODUM
14	1309364CD1	464	Y369 Y445 S118 T117 S217 T220 S232 S239 S340 T24 T32 T90 T137 S147 S232 T372 T428	N30 N215		BLAST_DOMO BLAST_PRODUM
15	1315267CD1	569	S3 T68 S85 S103 T229 S306 S356 T408 T482 T535 S551 T246 S20 T29 T31 T146 S167 T217 S292 T318 S385 T450 T477	N46 N121 N155 N304 N406	I465-E475 Muscarinic M4 receptor signature H386-A524 TCOMPLEX Male germ-cell specific protein	BLIMPS_PFAM BLAST_PRODUM
16	1403289CD1	436	S4 T17 S111 T167 T212 S222 S255 S308 S407 S421 S434 T35 S390 T58 T76 T97 T139 T153 T187 S213 S220 S235 S249 S270 Y74	N80 N336	L254-L275 L306-L327 Leucine Zipper K196-E395 COILED COIL MYOSIN REPEAT Q107-E395 TRICHOHYALIN (hair root sheath protein)	MOTIFS BLAST_PRODUM BLAST_DOMO

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
17	1607607CD1	363	S106 T206 T275 S288 S324 T331 T341 S51 T63 T162 S212 T336	N229 N307	L6-L27 L55-L76 Leucine Zipper L6-E16 Prepro orexin signature S3-E199 F33E11.3 PROTEIN similar to PHD finger K159-K165 Regulator of G protein signaling domain Q92-R103 5-hydroxytryptamine 2C receptor signature E60-P247 TOPOISOMERASE I DNA ISOMERASE REPEAT A17-S246 CYLICIN II sperm head cytoskeletal protein	MOTIFS BLIMPS_PRINTS BLAST_PRODUM BLIMPS_PPFAM BLIMPS_PRINTS BLAST_PRODUM BLAST_DOMO
18	1660025CD1	247	S366 S45 S69 T96 S139 S148 S161 S183 S238 T264 T392 S416 Y399 S224 T264 T369 S381		Q133-K383 COILED COIL MYOSIN REPEAT Q135-Q412 TRICHOHYALIN (hair root sheath protein)	BLAST_PRODUM BLAST_DOMO
19	1796836CD1	441	S366 S45 S69 T96 S139 S148 S161 S183 S238 T264 T392 S416 Y399 S224 T369 S381	N113 N128	WDREPEAT PROTEIN PF20 REPEAT WD FLAGELLA PD134845: E51-A178 LIM domains: C22-E80; C82-A139; C142-A208 LIM domain BL00478: Y43-L57 LIM domain signatures: E3-Y75; Y63-R206; M1-K137 LIM METAL-BINDING REPEAT DM00055 Q04584 464-533: F134-H203 LIM domain motifs: C22-L57; C82-I115; C142-L181	BLAST_PRODUM HMMER-PFAM BLIMPS-BLOCKS PROFILESSCAN BLAST-DOMO MOTIFS
20	2880670CD1	183	T48 S53 S68 T88	N66		BLAST_PRODUM
21	2913976CD1	212	S124 S143 T49 T52 Y75 Y172			BLAST_PRODUM
22	3092084CD1	227	T11 S79 S56 S58 S113 S221 S6 T7 S46 T201 Y141	N19	CALDESMON DM06224 P12957 1-755: T7-N197 (P-value = 7.6e-08)	BLAST_PRODUM



Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
23	3882482CD1	490	T318 S42 S71 S87 S143 T150 S174 T318 S416 S440 S448 T457 S473 S75 S145 T252 T278 S285 S314 S325 T382 Y377	N271	AAA ATP-binding protease domain: G243-R433 AAA-protein family protease signatures BL00674: W207-P227; W241-A262; S274-R316 G338-K384; G414-R433 KATANIN P60 SUBUNIT PD116869: M1-P135 AAA-PROTEIN FAMILY DM00024 P34808 188-348: D208-L368 AAA motif: V352-R370 ATP/GTP binding site (P-loop): G248-T255 Actin domain: M1-M114 Actins proteins signature BL00406: T5-K39 Actin signature PR00190: E24-V33 PROTEIN STRUCTURAL ACTIN MULTIGENE FAMILY ACETYLATION MUSCLE CYTOSKELETON CYTOPLASMIC ACTINLIKE PD000056: V6-L117 ACTINS AND ACTIN-RELATED PROTEINS DM00167 P20360 3-272: A2-M114 PROTEIN COILED COIL CHAIN MYOSIN REPEAT HEAVY ATPBINDING FILAMENT HEPTAD PD000002: L617-D679 (P-value = 2.5e-05) C2-DOMAIN DM00150 P24506 157-283: R460-S584 (P-value = 7.2e-06)	HMMER-PFAM BLIMPS-BLOCKS  BLAST-PRODOM BLAST-DOMO  MOTIFS MOTIFS HMMER-PFAM BLIMPS-BLOCKS BLIMPS-PRINTS BLAST-PRODOM  BLAST-DOMO  BLAST-DOMO
24	4933451CD1	133	S52 S115			
25	5043904CD1	912	T162 S592 S773 T28 S54 S63 S81 S135 S251 S260 S278 T299 T374 S383 S553 S565 T620 T636 S647 T690 S744 T786 S823 S880 S889 S908 S7 T50 S157 T192 S199 S243 T308 T326 S334 T542 T550 S576 S584 S640 T671 S752 S766 S774 S777 S780 T862 Y681	N483 N742		

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
26	5202390CD1	1076	S602 S903 T10 T45 T176 S204 T206 S393 S422 S454 S469 T475 T492 S494 S509 S518 S575 S592 T630 S632 S657 S732 S748 T759 T776 S822 S828 S880 T954 S982 S22 S57 S140 S184 T185 S203 T238 S253 S273 S316 S406 T410 S505 S548 T606 S626 S647 T835 T842 T856 T885 T898 T950 Y149 Y198 Y733 Y918	N150 N289 N312 N405 N421 N462	Calponin homology (CH) domain: P288-S393 ALPHA-ACTININ ACTIN-BINDING DOMAIN DM00325 P18091 28-252: A290-L385	HMMER-PFAM BLAST-DOMO
27	5526375CD1	542	S223 S313 S36 T80 S162 S280 S113 T158 S179 S218 S231 S272 T355 T366 S540	N188 N292	Kinesin motor domain: R31-N394 Kinesin motor domain BL00411: P25-E39; G100-G121; V157-L175 G216-L240; L264-L305; H314-P344 Kinesin motor domain signature: A247-A297 Kinesin heavy chain signatures PR00380: G100-G121; H225-L242; K263-R281; I315-T336 PROTEIN MOTOR ATP-BINDING COILED COIL MICROTUBULES KINESIN-LIKE KINESIN MITOSIS HEAVY PD000458; R31-L401 KINESIN MOTOR DOMAIN DM00198 P46871 3-343: E23-Q370 ATP/GTP binding site (P-loop): G109-T116 Kinesin motor domain motif: G262-E273 ATP/GTP binding site (P-loop): G38-T45	HMMER-PFAM BLIMPS-BLOCKS PROFILES SCAN BLIMPS-PRINTS BLAST-PRODOM BLAST-DOMO MOTIFS MOTIFS MOTIFS
28	5677408CD1	351	S340 T5 S22 T77 S92 S136 S186 S221 S284 T304 T105 S196 T205 T220 S343 Y336	N20		

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
29	5982278CD1	856	T13 S18 S807 S808 T84 S125 T200 S221 T266 S272 S334 T347 T369 T580 S710 T757 S763 T789 T823 S3 T8 S74 S116 S186 S187 S298 S341 T344 T366 T382 S605 T622 T663 T679 T793 S802	N48 N49 N345 N361 N565 N800	Kinesin motor domain: R31-E466 Kinesin motor domain BL00411: P25-D39; K69-Q85; G103-G124 G130-F140; Y216-L234; G283-I307 L333-L374; M385-P415 Kinesin motor domain signatures: D317-L364 Kinesin heavy chain signatures: PR00380: G103-G124; T292-L309; Q332-E350; V386-V407 PROTEIN MOTOR MICROTUBULES ATPBINDING COILED COIL KINESIN LIKE SIMILAR MITOTIC PROTEIN PD013891: E664-L841 KINESIN MOTOR DOMAIN DM00198   Q02241   4-443: A4-I446 ATP/GTP binding site (P-loop): G112-T119 Kinesin motor domain motif: S331-E342 LIM domain: C986-S1049 Calponin family repeat BL01052: F44-I69 LIM domain: F1008-L1022 LIM domain signature: K964-S1047 Calponin signature PR00889: S25-F39; C55-L72 CALPONIN FAMILY REPEAT DM01491   P51911   6-147: P4-S116 LIM domain motif: C986-L1022	HMMER-PFAM BLIMPS-BLOCKS PROFILESCAN BLIMPS-PRINTS BLAST-PRODROM BLAST-DOMO MOTIFS MOTIFS HMMER-PFAM BLIMPS-BLOCKS BLIMPS-BLOCKS PROFILESCAN BLIMPS-PRINTS BLAST-DOMO MOTIFS
30	6437362CD1	1056	T164 S256 S377 S408 S979 S38 T46 S173 S231 S233 S327 T411 S435 T451 S511 S601 S637 S746 S836 S846 S897 S951 S972 S1027 T35 S109 T113 S120 S169 S185 S251 S373 S403 T529 T537 T553 T630 S647 S702 S710 S753 S889 S974 S981 Y179 Y294	N272 N275 N475 N609	Ank repeat: K347-N379; N77-K109; K110-E142; T144-V176; S177-G209; L212-E244; N246-K278; T279-V311; S314-K346 signal_cleavage: M1-A18 signal_peptide: M1-A18 transmembrane domain: V2-N22, V277-G296, L326-V344 AC133 ANTIGEN PROMININ HOMOLOG: M1-R333	HMMER-PFAM SPSCAN HMMER HMMER BLAST_PRODROM
31	4173970CD1	1569				
32	2772751CD1	680	S113 S212 S340 S343 S357 T104 T159 T276 T328 T342	N210 N233 N251 N56 N66 N89		

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
33	2793768CD1	590	S3 S373 S432 S461 S495 S499 S548 S56 S561 S586 S83 T199 T228 T234 T255 T286 T331 T354 T357 T433 T445 T454 T534 T547 T588	N101 N166 N233	signal_cleavage: M1-S56 Ankyrin repeat: R40-R72, Q73-T102 SIMILAR TO ANKYRIN REPEAT REGION OF FOWLPOX VIRUS BAMHIORF7 PROTEIN: W75-H284, G348-Y469	SPSCAN HMMER_PFAM BLAST_PRODOR
34	3035248CD1	315	S151 S291 T115 T207 T273		signal_cleavage: M1-G26 signal_peptide: M1-S24 transmembrane domain: P4-F20 INTERMEDIATE FILAMENT ASSOCIATED PROTEIN K147-T218 TROPOMYOSIN ALTERNATIVE SPLICING SIGNAL PRECURSOR CHAIN: E39-S213 INTERMEDIATE FILAMENT: K121-T218	SPSCAN HMMER HMMR BLAST_PRODOR BLAST_PRODOR BLAST_DOMO

Table 4

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
35	1889577CB1	2345	902-952	3824958H1 (BRAXNOT01)	1	284
35	1889577CB1	2345		1915360R6 (PROSTUT04)	774	1278
35	1889577CB1	2345		1812980F6 (PROSTUT12)	102	646
35	1889577CB1	2345		3152565H1 (ADREN004)	1799	2066
35	1889577CB1	2345		1369763H1 (BSTMN002)	1608	1840
35	1889577CB1	2345		1784544F6 (BRAINOT10)	1873	2345
35	1889577CB1	2345		674685H1 (CRBLNOT01)	1516	1769
35	1889577CB1	2345		2838122F6 (DRGLNOT01)	1016	1563
35	1889577CB1	2345		1649402F6 (PROSTUT09)	384	896
36	2427982CB1	709		71263527V1	624	709
36	2427982CB1	709		71247061V1	1	683
37	2470833CB1	1569	1-721	1684180F6 (PROSNOT15)	1	497
37	2470833CB1	1569		868966R6 (LUNGAST01)	842	1407
37	2470833CB1	1569		3576193T6 (BRONNOT01)	980	1569
37	2470833CB1	1569		1534629F1 (SPLANNOT04)	677	1197
37	2470833CB1	1569		5296329H1 (COLENOT02)	493	730
37	2470833CB1	1569		1716065F6 (UCMCNOT02)	207	652
38	2080579CB1	1172	1-148, 686-854	868135H1 (BRAITUT03)	368	632
38	2080579CB1	1172		3458305F6 (293TF1T01)	1	433
38	2080579CB1	1172		2080579T6 (UTRSNOT08)	583	1143
38	2080579CB1	1172		2361824H1 (LUNGFET05)	939	1172
38	2080579CB1	1172		5174845H1 (EPIBTXT01)	442	637
39	2156553CB1	2380	1-360, 2126-2380, 1121-1655	5322363H1 (FIBPFEN06)	1	260
39	2156553CB1	2380		2916949T6 (THYMFET03)	1769	2380
39	2156553CB1	2380		866038X304D1 (BRAITUT03)	371	937
39	2156553CB1	2380		2916949F6 (THYMFET03)	1341	1886
39	2156553CB1	2380		2156553F6 (BRAINOT09)	705	1224
39	2156553CB1	2380		1722673F6 (BLADNOT06)	1176	1582
39	2156553CB1	2380		4899945H1 (OVARIT01)	245	531
39	2156553CB1	2380		1758833H1 (PITUNOT03)	236	485
40	2182855CB1	4396	2141-2667, 1- 1505, 1737- 1890, 3803- 4396	2816335F6 (BRSTNOT14)	3897	4396
40	2182855CB1	4396		2967273F6 (SCORNOT04)	580	1116

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
40	2182855CB1	4396		1611084F6 (COLNTUT06)	3554	3992
40	2182855CB1	4396		1785722H1 (BRAINOT10)	2610	2857
40	2182855CB1	4396		2182855F6 (SININOT01)	3389	3914
40	2182855CB1	4396		1484284F6 (CORPNOT02)	1	607
40	2182855CB1	4396		92276318	61	3799
40	2182855CB1	4396		2321435H1 (OVARNOT02)	2609	2794
40	2182855CB1	4396		1578313H1 (DUODNOT01)	1452	1546
40	2182855CB1	4396		1618459F6 (BRAITUT12)	2925	3464
40	2182855CB1	4396		2321435X308F1 (OVARNOT02)	1811	2391
41	2242106CB1	1831	1-509, 626-1018	965728R1 (BRSTNOT05)	1283	1831
41	2242106CB1	1831		1650350F6 (PROSTUT09)	322	1031
41	2242106CB1	1831		1396324T1 (THYRNOT03)	948	1660
41	2242106CB1	1831		6843794H1 (KIDNWTN03)	191	985
41	2242106CB1	1831		956964T1 (KIDNNOT05)	1019	1667
41	2242106CB1	1831		70846228V1	1	216
42	2726877CB1	3249	1979-2045, 2854-2873, 1857-1919, 2543-2608	3728286F6 (SMCCNON03)	1073	1504
42	2726877CB1	3249		3645568F6 (LUNGNOT34)	383	933
42	2726877CB1	3249		4969912H1 (KIDEUNC10)	151	425
42	2726877CB1	3249		2500944T6 (ADRETUT05)	1254	1839
42	2726877CB1	3249		2726877F6 (OVARUT05)	1603	2060
42	2726877CB1	3249		4195125T6 (COLITUT02)	2652	3249
42	2726877CB1	3249		4972430H1 (HELATXT02)	901	1191
42	2726877CB1	3249		4195125F6 (COLITUT02)	2257	2810
42	2726877CB1	3249		5492146H1 (DRGTN04)	2128	2380
42	2726877CB1	3249		3894803H1 (TLYMNOT05)	1	296
42	2726877CB1	3249		3728286T6 (SMCCNON03)	1682	2279
43	2738233CB1	4133	1-194, 4026-4133, 1607-2971, 3066-3186	1649466F6 (PROSTUT09)	2737	3394
43	2738233CB1	4133		2267313R6 (UTRSNOT02)	2211	2730
43	2738233CB1	4133		93882312_CD	126	3516
43	2738233CB1	4133		2945874H2 (BRAITUT23)	2710	3006
43	2738233CB1	4133		2242201F6 (PANCITUT02)	1538	2077

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment (s)	Sequence Fragments	5' Position	3' Position
43	2738233CB1	4133		653470R6 (EOSINOT03)	685	1068
43	2738233CB1	4133		2267313T6 (UTRSNOT02)	3481	4109
43	2738233CB1	4133		2962383H1 (ADRENOT09)	1	277
43	2738233CB1	4133		1467024F6 (PANCUT02)	1948	2420
43	2738233CB1	4133		3555108H1 (SYNONT01)	1394	1704
43	2738233CB1	4133		2965786H1 (SCORNOT04)	217	469
43	2738233CB1	4133		1262951R1 (SYNORAT05)	3959	4133
43	2738233CB1	4133		1260043T1 (MENITUT03)	3370	4017
43	2738233CB1	4133		1486351H1 (CORPNOT02)	1236	1463
43	2738233CB1	4133		3765643F6 (BRSTNOT24)	277	758
44	1833116CB1	1754	1700-1754	2852676F6 (BRSTTUT13)	25	529
44	1833116CB1	1754		3128645H1 (LUNGUTUT12)	539	839
44	1833116CB1	1754		5016828H1 (BRAXNOT03)	632	886
44	1833116CB1	1754		413418R1 (BRSTNOT01)	979	1615
44	1833116CB1	1754		1442616R1 (THYRNOT03)	1247	1754
44	1833116CB1	1754		1785591H1 (BRAINOT10)	1	284
44	1833116CB1	1754		5172858H1 (EPIPTXT01)	331	588
44	1833116CB1	1754		1920612R6 (BRSTTUT01)	866	1402
45	001799CB1	2713	1-27, 1464-2008	5994129H1 (FTUBTUT02)	587	912
45	001799CB1	2713		4245126H1 (BRADIT01)	1982	2239
45	001799CB1	2713		6818763J1 (BRAUNOR01)	1	549
45	001799CB1	2713		5054327H1 (COLATMT01)	1456	1726
45	001799CB1	2713		6739739H1 (BRAFDIT02)	2153	2713
45	001799CB1	2713		71336820V1	276	898
45	001799CB1	2713		3730557H1 (SMCCNON03)	884	1198
45	001799CB1	2713		644891R6 (BRSTTUT02)	1575	1981
45	001799CB1	2713		3515211H1 (LUNGNOT33)	1940	2217
45	001799CB1	2713		2691467T6 (LUNGNOT23)	907	1471
45	001799CB1	2713		3240741H1 (COLAUCT01)	1401	1685
45	001799CB1	2713		4771110H1 (BRATNOT02)	1715	1990
46	119814CB1	1768	1-688	2637776F6 (BONTNOT01)	968	1494
46	119814CB1	1768		119814R1 (MUSCNOT01)	739	1406
46	119814CB1	1768		2638913F6 (BONTNOT01)	1410	1768
46	119814CB1	1768		1993563H1 (CORPNOT02)	656	940
46	119814CB1	1768		g2184959	349	846
46	119814CB1	1768		g1218792	1	579
46	119814CB1	1768		2395927T6 (THP1AZT01)	20	182

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
47	1295420CB1	3287	914-1321, 177-207	148434H1 (FIBRNGT01)	812	1028
47	1295420CB1	3287		2697808H1 (UTRSNOT12)	675	877
47	1295420CB1	3287		5778293H1 (BRAXNOT03)	1185	1447
47	1295420CB1	3287		2149151F6 (BRAINOT09)	1661	2176
47	1295420CB1	3287		2883729F6 (SINJNOT02)	1	515
47	1295420CB1	3287		1888639H1 (BLADTUT07)	868	1147
47	1295420CB1	3287		2154634F6 (BRAINOT09)	2720	3278
47	1295420CB1	3287		3703932T6 (PENCNOT07)	1334	2009
47	1295420CB1	3287		998245R1 (KIDNTUT01)	2118	2637
47	1295420CB1	3287		2223877F6 (SEMNOT01)	323	816
47	1295420CB1	3287		4027885H1 (BRAINOT23)	999	1261
47	1295420CB1	3287		835343R1 (PROSNOT07)	2247	2814
47	1295420CB1	3287		1367731R1 (SCORNON02)	2856	3287
48	1309364CB1	1748	1-49, 1037-1135	4701455H1 (SMCRTXT01)	1162	1416
48	1309364CB1	1748		4904644F6 (TYMNOT08)	799	1399
48	1309364CB1	1748		2914466F6 (THYMFE03)	1	536
48	1309364CB1	1748		5590953H1 (ENDINOT02)	678	933
48	1309364CB1	1748		1309364F6 (COLNFET02)	1227	1748
48	1309364CB1	1748		3727909H1 (SMCCNON03)	513	816
49	1315267CB1	2163	705-799	898915H1 (BRSTTUT03)	1839	2163
49	1315267CB1	2163		465550R6 (LATRNOT01)	673	1257
49	1315267CB1	2163		1575785F6 (LNODNOT03)	1029	1630
49	1315267CB1	2163		5191222F6 (OVARIT06)	1	538
49	1315267CB1	2163		5207783F6 (BRAFNOT02)	426	1041
49	1315267CB1	2163		1315267F6 (BLADTUT02)	1434	2022
50	1403289CB1	1615	1119-1170	2811792T6 (OVARNOT10)	489	1121
50	1403289CB1	1615		059048R6 (MUSCNOT01)	1080	1615
50	1403289CB1	1615		3502723H1 (ADRENOT11)	932	1225
50	1403289CB1	1615		1403289F6 (LATRTUT02)	1	601
51	1607607CB1	1356	1-157, 1263-1356	7262994H1 (UTRETM01)	252	929
51	1607607CB1	1356		3467640F6 (BRAIDIT01)	350	1048
51	1607607CB1	1356		2137437H1 (ENDCNOT01)	1	275
51	1607607CB1	1356		1607607F6 (LUNGNOT15)	872	1356
52	1660025CB1	1268	1-88, 424- 459, 775-836	2580277F6 (KIDNTUT13)	213	932



Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
52	1660025CB1	1268		2172241H1 (ENDCNOT03)	1	240
52	1660025CB1	1268		1988094R6 (LUNGAST01)	750	1268
52	1660025CB1	1268		1756804R6 (PITUNOT03)	248	966
53	1796836CB1	2554	1177-1245, 1-61	2280307T6 (PROSNON01)	687	1302
53	1796836CB1	2554		4971676H1 (HELATXT02)	1	227
53	1796836CB1	2554		2582430T6 (KIDNTUT13)	1198	1545
53	1796836CB1	2554		2497103T6 (ADRETUT05)	1949	2533
53	1796836CB1	2554		2534742H1 (BRAINOT18)	1527	1758
53	1796836CB1	2554		2553754T6 (THYMNOT03)	1817	2528
53	1796836CB1	2554		2726708H1 (OVARFUT05)	1630	1863
53	1796836CB1	2554		276683H1 (TESTNOT03)	1311	1568
53	1796836CB1	2554		2938533H1 (THYMFET02)	1056	1326
53	1796836CB1	2554		6914750J1 (PITUDIR01)	47	684
53	1796836CB1	2554		2300549R6 (BRSTNOT05)	2162	2554
53	1796836CB1	2554		3030841F6 (HEARFET02)	319	877
54	2880670CB1	1216	605-636	2889280T7 (LUNGFET04)	489	1192
54	2880670CB1	1216		1358092F1 (LUNGNOT09)	246	920
54	2880670CB1	1216		816703R1 (OVARFUT01)	657	1216
54	2880670CB1	1216		2529604H1 (GBLANOT02)	1	357
55	2913976CB1	1457	1-446, 1406-1457	3736188F6 (SMCCNOS01)	1173	1428
55	2913976CB1	1457		2913976F6 (KIDNTUT15)	1	520
55	2913976CB1	1457		4645636H1 (PROSTUT20)	1187	1449
55	2913976CB1	1457		4331439H1 (KIDNNOT32)	461	712
55	2913976CB1	1457		4643722H1 (PROSTMT03)	1108	1325
55	2913976CB1	1457		1312116F1 (COLNFET02)	583	1127
56	3092084CB1	1636	857-1636	1709866F6 (PROSNOT16)	1097	1634
56	3092084CB1	1636		2807436F6 (BLADTUT08)	1277	1636
56	3092084CB1	1636		6906626H1 (MUSLTDR02)	1	610
56	3092084CB1	1636		SBMA03169F1	597	1181
56	3092084CB1	1636		3092084F6 (BRSTNOT19)	543	1147
57	3882482CB1	1742	1-82, 923-994	2286328X19F1 (BRAINON01)	742	1251
57	3882482CB1	1742		2804312F6 (PENCNOT01)	299	926
57	3882482CB1	1742		986917T6 (LVENNOT03)	1088	1742
57	3882482CB1	1742		3175528F6 (UTRSTUT04)	1	317
57	3882482CB1	1742		1232503F6 (LUNGFET03)	960	1553

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
57	3882482CB1	1742		2286328X16F1 (BRAINON01)	402	969
58	4933451CB1	602	1-401	4931591H1 (BRSTTUT20)	341	602
58	4933451CB1	602		2502933F6 (CONUTUT01)	1	479
59	5043904CB1	3237	1964-2007, 2911-2969, 993-1057, 1328-1589	2153280F6 (BRAINOT09)	335	888
59	5043904CB1	3237		5043904R6 (PLACFER01)	1	583
59	5043904CB1	3237		075538H1 (THP1PEB01)	1604	1818
59	5043904CB1	3237		3189755X301D1 (THYMNON04)	1131	1674
59	5043904CB1	3237		2460935F6 (THYRNOT08)	614	1177
59	5043904CB1	3237		1795345R6 (PROSTUT05)	2026	2510
59	5043904CB1	3237		1865333F6 (PROSNOT19)	2686	3237
59	5043904CB1	3237		1365052R6 (SCORNON02)	2440	2957
59	5043904CB1	3237		3250182H1 (SEMVNOT03)	1831	2128
59	5043904CB1	3237		4713986H1 (BRAIHCT01)	2266	2513
59	5043904CB1	3237		3804331H1 (BLADTUT03)	1777	2089
60	5202390CB1	3640	1412-1489, 2294-2508, 1-161, 3244-3640	2544502H2 (UTRSNOT11)	2656	2927
60	5202390CB1	3640		2321656R6 (OVARNOT02)	2039	2448
60	5202390CB1	3640		2557486F6 (THYMNOT03)	1	659
60	5202390CB1	3640		1441193F6 (THYRNOT03)	3317	3640
60	5202390CB1	3640		2844888H1 (DRGLNOT01)	864	1009
60	5202390CB1	3640		3705507H1 (PENCNOT07)	3047	3352
60	5202390CB1	3640		2122377F6 (BRSTNOT07)	2844	3297
60	5202390CB1	3640		4313841F6 (BRAFNOT01)	2322	2662
60	5202390CB1	3640		5092148H1 (UTRSTMR01)	685	961
60	5202390CB1	3640		g4240294 CD	525	3562
60	5202390CB1	3640		1944075H1 (PITUNOT01)	1590	1851
60	5202390CB1	3640		3493516H1 (ADRETUT07)	1999	2279
60	5202390CB1	3640		176099H1 (TLYMNOT01)	411	738
60	5202390CB1	3640		2851783H1 (BRSTTUT13)	2600	2796
60	5202390CB1	3640		534176R1 (BRAINOT03)	1041	1567
60	5202390CB1	3640		4107451H1 (BRSTTUT17)	1431	1694
60	5202390CB1	3640		2842892F6 (DRGLNOT01)	1604	2220

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
61	5526375CB1	2111	1-50, 1540-2111	3038423H1 (BRSTNOT16)	1004	1278
61	5526375CB1	2111		2513433F6 (LIVRTUT04)	1600	2111
61	5526375CB1	2111		3534575H1 (KIDNNOT25)	1263	1529
61	5526375CB1	2111		2993194F6 (KIDNFEF02)	729	1224
61	5526375CB1	2111		5109694H1 (PROSTUS19)	1385	1626
61	5526375CB1	2111		2993263H1 (KIDNFEF02)	1242	1479
61	5526375CB1	2111		3534157H1 (KIDNNOT25)	518	788
61	5526375CB1	2111		2580307F6 (KIDNTUT13)	1	519
61	5526375CB1	2111		2070882F6 (ISLTNOT01)	225	715
62	5677408CB1	1389	1-177	535789R1 (ADRENOT03)	1079	1389
62	5677408CB1	1389		881149T6 (THYRNOT02)	793	1378
62	5677408CB1	1389		6023544H1 (TESTNOT11)	736	1017
62	5677408CB1	1389		881149R6 (THYRNOT02)	1	772
63	5982278CB1	3331	809-1149, 1755-1989	5260541H1 (CONDUTUT01)	2203	2470
63	5982278CB1	3331		1390622H1 (EOSINOT01)	2029	2257
63	5982278CB1	3331		4515063H1 (EPIMNOT01)	1740	1993
63	5982278CB1	3331		3584113H1 (293TF4T01)	2462	2796
63	5982278CB1	3331		3405843H1 (ESOGNOT03)	919	1174
63	5982278CB1	3331		5261482H1 (CONDUTUT01)	2611	2862
63	5982278CB1	3331		1637273H1 (UTRSNOT06)	1619	1733
63	5982278CB1	3331		g1521431	2559	3331
63	5982278CB1	3331		3591491H1 (293TF5T01)	1674	1981
63	5982278CB1	3331		4957640H1 (TLYMNOT05)	1464	1726
63	5982278CB1	3331		4666088H1 (MEGBUNT01)	2280	2536
63	5982278CB1	3331		043258H1 (TBLYNOT01)	1370	1562
63	5982278CB1	3331		2907496F6 (THYMNOT05)	24	624
63	5982278CB1	3331		4983673H1 (HELATXT05)	1865	2134
63	5982278CB1	3331		g34671_CD	130	3166
63	5982278CB1	3331		3449505X304D1 (UTRSNON03)	942	1450
63	5982278CB1	3331		2640427T6 (LUNGUT08)	2646	3309
63	5982278CB1	3331		2205131F6 (SPLNFEF02)	1	514
64	6437362CB1	3558	1-428, 3352-3558, 923-1583	720069R6 (SYNOOAT01)	3159	3558
64	6437362CB1	3558		2785980H1 (BRSTNOT13)	2462	2730
64	6437362CB1	3558		1568793H1 (UTRSNOT05)	1264	1467

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
64	6437362CB1	3558		987366H1 (LVENNOT03)	763	1088
64	6437362CB1	3558		g5689540_CD	678	3298
64	6437362CB1	3558		4567664F6 (HELATXT01)	1	500
64	6437362CB1	3558		1988667R6 (LUNGAST01)	2160	2709
64	6437362CB1	3558		4980789H1 (HELATXT04)	981	1248
64	6437362CB1	3558		6437362H1 (LUNGNON07)	432	1052
64	6437362CB1	3558		2903936F6 (DRGCNOT01)	1760	2328
64	6437362CB1	3558		3208061H1 (PENCNOT03)	2706	2948
64	6437362CB1	3558		3288660F6 (BONRFET01)	1433	2050
64	6437362CB1	3558		865171R1 (BRAITUT03)	2929	3483
65	4173970CB1	5373	3418-5373, 1-186, 1641-2444, 857-1058	829704R1 (PROSTUT04)	1356	1950
65	4173970CB1	5373		5604442H1 (MONOTXN03)	2038	2310
65	4173970CB1	5373		1708630F6 (PROSNOT16)	3923	4539
65	4173970CB1	5373		4561514F6 (KERATXT01)	4397	5197
65	4173970CB1	5373		1437088F1 (PANCNOT08)	3691	4247
65	4173970CB1	5373		1433309R1 (BEPINON01)	3666	4214
65	4173970CB1	5373		4167822F6 (PANCNOT21)	1	494
65	4173970CB1	5373		1806736F6 (SINTNOT13)	5043	5373
65	4173970CB1	5373		3508537F6 (CONCNOT01)	2520	3017
65	4173970CB1	5373		4173970F6 (SINTNOT21)	2115	2677
65	4173970CB1	5373		2277402R6 (PROSNON01)	660	1229
65	4173970CB1	5373		1708630T6 (PROSNOT16)	4639	5347
65	4173970CB1	5373		5944958H1 (COLADIT05)	1710	2018
65	4173970CB1	5373		1300156F1 (BRSTNOT07)	1133	1687
65	4173970CB1	5373		g2737563	1885	2171
65	4173970CB1	5373		209752R1 (SPLNNOT02)	3040	3690
65	4173970CB1	5373		287603R1 (EOSIHET02)	210	966
65	4173970CB1	5373		3091106H1 (BRSTNOT19)	4284	4560
65	4173970CB1	5373		516280R6 (MMLR1DT01)	2996	3438
66	2772751CB1	4333	2456-3205, 100-160, 1-23, 1459-1957, 3695-4333	70475866V1	507	1008
66	2772751CB1	4333		70472414V1	2045	2690

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
66	2772751CB1	4333		3402774H1 (ESOGNOT03)	1	256
66	2772751CB1	4333		70475304V1	1481	2093
66	2772751CB1	4333		70475403V1	976	1554
66	2772751CB1	4333		70470913V1	1545	2109
66	2772751CB1	4333		70747026V1	3561	4054
66	2772751CB1	4333		g3923880	319	678
66	2772751CB1	4333		1298255T6 (BRSTNOT07)	3001	3690
66	2772751CB1	4333		70472159V1	1049	1556
66	2772751CB1	4333		70472656V1	2219	2925
66	2772751CB1	4333		g5340324	24	467
66	2772751CB1	4333		6849173H1 (KIDNTMN03)	3801	4333
66	2772751CB1	4333		6221536U1	2737	3452
67	2793768CB1	2213	2186-2213, 1066-1156	70843048V1	1646	2213
67	2793768CB1	2213		2026465R6 (KERANOT02)	709	1268
67	2793768CB1	2213		7712268H1 (TESTTUE02)	1621	2213
67	2793768CB1	2213		g1958420	1	421
67	2793768CB1	2213		6584157H1 (ESOGTMC01)	984	1576
67	2793768CB1	2213		2793768F6 (COLNTUT16)	131	777
67	2793768CB1	2213		71279716V1	1510	2183
68	3035248CB1	1142	1-55, 555-605	71515027V1	585	1140
68	3035248CB1	1142		71486327V1	402	787
68	3035248CB1	1142		71514455V1	1	576

Table 5

Polynucleotide SEQ ID NO:	Incyte Project ID	Representative Library
35	1889577CB1	PROSTUT12
36	2427982CB1	DRGCNOT01
37	2470833CB1	LUNGAST01
38	2080579CB1	UTRSNOT08
39	2156553CB1	THYMFET03
40	2182855CB1	SCORNOT04
41	2242106CB1	COLNPOT01
42	2726877CB1	LUNGNOT34
43	2738233CB1	MENITUT03
44	1833116CB1	THYRNOT03
45	001799CB1	BRSTTUT02
46	119814CB1	MUSCNOT01
47	1295420CB1	BRAITUT12
48	1309364CB1	TYLMONT01
49	1315267CB1	BLADTUT02
50	1403289CB1	LATRTUT02
51	1607607CB1	BRAIDIT01
52	1660025CB1	BRAWNOT01
53	1796836CB1	BRSTNOT05
54	2880670CB1	OVARTUT01
55	2913976CB1	ENDCNOT04
56	3092084CB1	HEAANOT01
57	3882482CB1	SPLNNOT11
58	4933451CB1	BRSTTUT20
59	5043904CB1	PLACFER01
60	5202390CB1	TESTTUT02
61	5526375CB1	KIDNFET02
62	5677408CB1	ADRENOT03
63	5982278CB1	SPLNFET02
64	6437362CB1	BRAINOT23
65	4173970CB1	BRSTNOT07
66	2772751CB1	BRSTNOT07
67	2793768CB1	UTRSNOT12
68	3035248CB1	TYLMNOT05

Table 6

Library	Vector	Library Description
ADRENOT03	PSPORT1	Library was constructed using RNA isolated from the adrenal tissue of a 17-year-old Caucasian male, who died from cerebral anoxia.
BLADTUT02	pINCY	Library was constructed using RNA isolated from bladder tumor tissue removed from an 80-year-old Caucasian female during a radical cystectomy and lymph node excision. Pathology indicated grade 3 invasive transitional cell carcinoma. Family history included acute renal failure, osteoarthritis, and atherosclerosis.
BRAIDIT01	pINCY	Library was constructed using RNA isolated from diseased brain tissue. Patient history included multiple sclerosis, type II lesion.
BRAINOT23	pINCY	Library was constructed using RNA isolated from right temporal lobe tissue removed from a 45-year-old Black male during a brain lobectomy. Pathology for the associated tumor tissue indicated dysembryoplastic neuroepithelial tumor of the right temporal lobe. The right temporal region dura was consistent with calcifying pseudotumor of the neuraxis. The patient presented with convulsive intractable epilepsy, partial epilepsy, and memory disturbance. Patient history included obesity, meningitis, backache, unspecified sleep apnea, acute stressreaction, acquired knee deformity, and chronic sinusitis. Family history included obesity, benign hypertension, cirrhosis of the liver, alcohol abuse, hyperlipidemia, cerebrovascular disease, and type II diabetes.
BRAITUT12	pINCY	Library was constructed using RNA isolated from brain tumor tissue removed from the left frontal lobe of a 40-year-old Caucasian female during excision of a cerebral meningeal lesion. Pathology indicated grade 4 gemistocytic astrocytoma.
BRAWNOT01	pINCY	Library was constructed using RNA isolated from dentate nucleus tissue removed from the brain of a 35-year-old Caucasian male who died from cardiac failure. Pathology indicated moderate leptomeningeal fibrosis and multiple microinfarctions of the cerebral neocortex. Grossly, the brain regions examined and cranial nerves were unremarkable, showing no evidence of atrophy. No atherosclerosis of the major vessels was noted. Microscopically, the cerebral hemisphere revealed moderate fibrosis of the leptomeninges with focal calcifications. There was evidence of shrunken and slightly eosinophilic pyramidal neurons throughout the cerebral hemispheres. There were also multiple small microscopic areas of cavitation with surrounding gliosis scattered throughout the cerebral cortex. Special stains with Bielschowsky silver, Klüver-Barrera, and Congo Red revealed no evidence of neurofibrillary tangles or diffuse anorectic amyloid plaques, demyelination, and cerebral amyloid angiopathy, respectively. Patient history included dilated cardiomyopathy, congestive heart failure, cardiomegaly, and an enlarged spleen and liver. Patient medications included simethicone, Lasix, Digoxin, Colace, Zantac, captopril, and Vasotec.
BRSTNOT05	PSPORT1	Library was constructed using RNA isolated from breast tissue removed from a 58-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology for the associated tumor tissue indicated multicentric invasive grade 4 lobular carcinoma. Patient history included skin cancer, rheumatic heart disease, osteoarthritis, and tuberculosis. Family history included cerebrovascular and cardiovascular disease, breast and prostate cancer, and type I diabetes.

Table 6 (cont.)

Library	Vector	Library Description
BRSTNOT07	pINCY	Library was constructed using RNA isolated from diseased breast tissue removed from a 43-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology indicated mildly proliferative fibrocystic changes with epithelial hyperplasia, papillomatosis, and duct ectasia. Pathology for the associated tumor tissue indicated invasive grade 4, nuclear grade 3 mammary adenocarcinoma with extensive comedo necrosis. Family history included epilepsy, cardiovascular disease, and type II diabetes.
BRSTNOT07	pINCY	Library was constructed using RNA isolated from diseased breast tissue removed from a 43-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology indicated mildly proliferative fibrocystic changes with epithelial hyperplasia, papillomatosis, and duct ectasia. Pathology for the associated tumor tissue indicated invasive grade 4, nuclear grade 3 mammary adenocarcinoma with extensive comedo necrosis. Family history included epilepsy, cardiovascular disease, and type II diabetes.
BRSTTUT02	PSPORT1	Library was constructed using RNA isolated from breast tumor tissue removed from a 54-year-old Caucasian female during a bilateral radical mastectomy with reconstruction. Pathology indicated residual invasive grade 3 mammary ductal adenocarcinoma. The remaining breast parenchyma exhibited proliferative fibrocystic changes without atypia. One of 10 axillary lymph nodes had metastatic tumor as a microscopic intranodal focus. Patient history included kidney infection and condyloma acuminatum. Family history included benign hypertension, hyperlipidemia, and a malignant colon neoplasm.
BRSTTUT20	pINCY	Library was constructed using RNA isolated from left breast tumor tissue removed from a 66-year-old Black female during a unilateral extended simple mastectomy and fine needle breast biopsy. Pathology indicated invasive grade 4, nuclear grade 3 adenocarcinoma ductal type, diffusely replacing the left breast. The skin, nipple and fascia were all involved, including the deep surgical margin. Extensive angiolymphatic invasion was identified, including superficial dermal lymphatics. Metastatic grade 4 adenocarcinoma completely replaced 6 lymph nodes with extranodal extension. Multiple low axillary lymph nodes tissue were positive for metastatic mammary carcinoma. Left chest wall biopsy indicated metastatic grade 4 adenocarcinoma. Prior left breast biopsy indicated metastatic grade 4, nuclear grade 3, metastatic mammary carcinoma. The patient presented with malaise and fatigue. Patient history included secondary malignant neoplasm of the liver, secondary malignant neoplasm of the brain/spine, deficiency anemia, type II diabetes, chronic renal failure, and normal delivery. Patient medications included two cycles of cyclophosphamide/epirubicin and 5-Fluorouracil in November 1995. Family history included benign hypertension, type II diabetes, hyperlipidemia, and depressive disorder in the mother.
COLNPOT01	pINCY	Library was constructed using RNA isolated from colon polyp tissue removed from a 40-year-old Caucasian female during a total colectomy. Pathology indicated an inflammatory pseudopolyp; this tissue was associated with a focally invasive grade 2 adenocarcinoma and multiple tubovillous adenomas. Patient history included a benign neoplasm of the bowel.



Table 6 (cont.)

Library	Vector	Library Description
DRGCNOT01	pINCY	Library was constructed using RNA isolated from dorsal root ganglion tissue removed from the cervical spine of a 32-year-old Caucasian male who died from acute pulmonary edema and bronchopneumonia, bilateral pleural and pericardial effusions, and malignant lymphoma (natural killer cell type). Patient history included probable cytomegalovirus, infection, hepatic congestion and steatosis, splenomegaly, hemorrhagic cystitis, thyroid hemorrhage, and Bell's palsy. Surgeries included colonoscopy, large intestine biopsy, adenotonsillectomy, and nasopharyngeal endoscopy and biopsy; treatment included radiation therapy.
ENDCNOT04	pINCY	Library was constructed using RNA isolated from coronary artery endothelial cell tissue removed from a 3-year-old Caucasian male.
HEANOT01	pINCY	Library was constructed using RNA isolated from right coronary and right circumflex coronary artery tissue removed from the explanted heart of a 46-year-old Caucasian male during a heart transplantation. Patient history included myocardial infarction from total occlusion of the left anterior descending coronary artery, atherosclerotic coronary artery disease, hyperlipidemia, myocardial ischemia, dilated cardiomyopathy, left ventricular dysfunction, and tobacco abuse. Previous surgeries included cardiac catheterization. Family history included atherosclerotic coronary artery disease.
KIDNFET02	pINCY	Library was constructed using RNA isolated from kidney tissue removed from a Caucasian male fetus, who was stillborn with a hypoplastic left heart and died at 23 weeks' gestation.
LATRTUT02	pINCY	Library was constructed using RNA isolated from a myxoma removed from the left atrium of a 43-year-old Caucasian male during annuloplasty. Pathology indicated atrial myxoma. Patient history included pulmonary insufficiency, acute myocardial infarction, atherosclerotic coronary artery disease, hyperlipidemia, and tobacco use. Family history included benign hypertension, acute myocardial infarction, atherosclerotic coronary artery disease, and type II diabetes.
LUNGAST01	PSPORT1	Library was constructed using RNA isolated from the lung tissue of a 17-year-old Caucasian male, who died from head trauma. Patient history included asthma.
LUNGNOT34	pINCY	Library was constructed using RNA isolated from lung tissue removed from a 12-year-old Caucasian male.
MENITUT03	pINCY	Library was constructed using RNA isolated from brain meningioma tissue removed from a 35-year-old Caucasian female during excision of a cerebral meningeal lesion. Pathology indicated a benign neoplasm in the right cerebellopontine angle of the brain. Patient history included hypothyroidism. Family history included myocardial infarction and breast cancer.
MUSCNOT01	PBLUESCRIPT	Library was constructed at Stratagene (STR937209), using RNA isolated from the skeletal muscle tissue of a patient with malignant hyperthermia.

Table 6 (cont.)

Library	Vector	Library Description
OVARTUT01	PSPORT1	Library was constructed using RNA isolated from ovarian tumor tissue removed from a 43-year-old Caucasian female during removal of the fallopian tubes and ovaries. Pathology indicated grade 2 mucinous cystadenocarcinoma involving the entire left ovary. Patient history included mitral valve disorder, pneumonia, and viral hepatitis. Family history included atherosclerotic coronary artery disease, pancreatic cancer, stress reaction, cerebrovascular disease, breast cancer, and uterine cancer.
PLACFER01	pINCY	The library was constructed using RNA isolated from placental tissue removed from a Caucasian fetus, who died after 16 weeks' gestation from fetal demise and hydrocephalus. Patient history included umbilical cord wrapped around the head (3 times) and the shoulders (1 time). Serology was positive for anti-CMV. Family history included multiple pregnancies and live births, and an abortion.
PROSTUT12	pINCY	Library was constructed using RNA isolated from prostate tumor tissue removed from a 65-year-old Caucasian male during a radical prostatectomy. Pathology indicated an adenocarcinoma (Gleason grade 2+2). Adenofibromatous hyperplasia was also present. The patient presented with elevated prostate specific antigen (PSA).
SCORNOT04	pINCY	Library was constructed using RNA isolated from cervical spinal cord tissue removed from a 32-year-old Caucasian male who died from acute pulmonary edema and bronchopneumonia, bilateral pleural and pericardial effusions, and malignant lymphoma (natural killer cell type). Patient history included probable cytomegalovirus, infection, hepatic congestion and steatosis, splenomegaly, hemorrhagic cystitis, thyroid hemorrhage, and Bell's palsy. Surgeries included colonoscopy, large intestine biopsy, adenotonsillectomy, and nasopharyngeal endoscopy and biopsy; treatment included radiation therapy.
SPLNFET02	pINCY	Library was constructed using RNA isolated from spleen tissue removed from a Caucasian male fetus, who died at 23 weeks' gestation.
SPLNNOT11	pINCY	Library was constructed using RNA isolated from diseased spleen tissue removed from a 14-year-old Asian male during a total splenectomy. Pathology indicated changes consistent with idiopathic thrombocytopenic purpura. The patient presented with bruising. Patient medications included Vincristine.
TESTTUT02	pINCY	Library was constructed using RNA isolated from testicular tumor removed from a 31-year-old Caucasian male during unilateral orchiectomy. Pathology indicated embryonal carcinoma.
THYMFET03	pINCY	Library was constructed using RNA isolated from thymus tissue removed from a Caucasian male fetus.
THYRNOT03	pINCY	Library was constructed using RNA isolated from thyroid tissue removed from the left thyroid of a 28-year-old Caucasian female during a complete thyroidectomy. Pathology indicated a small nodule of adenomatous hyperplasia present in the left thyroid. Pathology for the associated tumor tissue indicated dominant follicular adenoma, forming a well-encapsulated mass in the left thyroid.
TYLMNOT05	pINCY	Library was constructed RNA isolated from nonactivated Th2 cells. These cells were differentiated from umbilical cord CD4 T cells with IL-4 in the presence of anti-IL-12 antibodies and B7-transfected COS cells.

Table 6 (cont.)

Library	Vector	Library Description
TLYMUNT01	pINCY	Library was constructed using RNA isolated from restingallogenic T-lymphocyte tissue removed from an adult (40-50-year-old) Caucasian male.
UTRSNOT08	pINCY	Library was constructed using RNA isolated from uterine tissue removed from a 35-year-old Caucasian female during a vaginal hysterectomy with dilation and curettage. Pathology indicated that the endometrium was secretory phase with a benign endometrial polyp 1 cm in diameter. The cervix showed mild chronic cervicitis. Family history included atherosclerotic coronary artery disease and type II diabetes.
UTRSNOT12	pINCY	Library was constructed using RNA isolated from uterine myometrial tissue removed from a 41-year-old Caucasian female during a vaginal hysterectomy with dilation and curettage. The endometrium was secretory and contained fragments of endometrial polyps. Benign endo- and ectocervical mucosa were identified in the endocervix. Pathology for the associated tumor tissue indicated uterine leiomyoma. Patient history included ventral hernia and a benign ovarian neoplasm.

Table 7

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25:3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. USA 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183:63-98; and Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value=1.06E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value=1.0E-8 or less Full Length sequences: fastx score=100 or greater
BLIMPS	A BLocks IMProved Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff (1991) Nucleic Acids Res. 19:6565-6572; Henikoff, J.G. and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37:417-424.	Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) J. Mol. Biol. 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322; Durbin, R. et al. (1998) Our World View, in a Nutshell, Cambridge Univ. Press, pp. 1-350.	PFAM hits: Probability value= 1.0E-3 or less Signal peptide hits: Score= 0 or greater

Table 7 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, M. et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221.	Normalized quality score>GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M.S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies.	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12:431-439.	Score=3.5 or greater
TMAP	A program that uses weight matrices to delineate transmembrane segments on protein sequences and determine orientation.	Persson, B. and P. Argos (1994) J. Mol. Biol. 237:182-192; Persson, B. and P. Argos (1996) Protein Sci. 5:363-371.	
TMHMMER	A program that uses a hidden Markov model (HMM) to delineate transmembrane segments on protein sequences and determine orientation.	Sonnhammer, E.L. et al. (1998) Proc. Sixth Intl. Conf. on Intelligent Systems for Mol. Biol., Glasgow et al., eds., The Am. Assoc. for Artificial Intelligence Press, Menlo Park, CA, pp. 175-182.	
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. An isolated polypeptide selected from the group consisting of:

a) a polypeptide comprising an amino acid sequence selected from the group consisting of

5 SEQ ID NO:1-34,

b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-34,

c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, and

10 d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34.

2. An isolated polypeptide of claim 1 selected from the group consisting of SEQ ID NO:1-34.

15

3. An isolated polynucleotide encoding a polypeptide of claim 1.

4. An isolated polynucleotide encoding a polypeptide of claim 2.

20 5. An isolated polynucleotide of claim 4 selected from the group consisting of SEQ ID NO:35-68.

6. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.

25

7. A cell transformed with a recombinant polynucleotide of claim 6.

8. A transgenic organism comprising a recombinant polynucleotide of claim 6.

30 9. A method for producing a polypeptide of claim 1, the method comprising:

a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and

35 b) recovering the polypeptide so expressed.

10. An isolated antibody which specifically binds to a polypeptide of claim 1.

11. An isolated polynucleotide selected from the group consisting of:

- a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting  
5 of SEQ ID NO:35-68,  
b) a naturally occurring polynucleotide comprising a polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:35-68,  
c) a polynucleotide complementary to a polynucleotide of a),  
d) a polynucleotide complementary to a polynucleotide of b), and  
10 e) an RNA equivalent of a)-d).

12. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 11.

13. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

- a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization  
20 complex is formed between said probe and said target polynucleotide or fragments thereof, and  
b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

14. A method of claim 13, wherein the probe comprises at least 60 contiguous nucleotides.

15. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

- a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and  
30 b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

16. A composition comprising a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

17. A composition of claim 16, wherein the polypeptide has an amino acid sequence selected from the group consisting of SEQ ID NO:1-34.

18. A method for treating a disease or condition associated with decreased expression of functional CYSKP, comprising administering to a patient in need of such treatment the composition of claim 16.

19. A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

- 10       a) exposing a sample comprising a polypeptide of claim 1 to a compound, and  
         b) detecting agonist activity in the sample.

20. A composition comprising an agonist compound identified by a method of claim 19 and a pharmaceutically acceptable excipient.

15

21. A method for treating a disease or condition associated with decreased expression of functional CYSKP, comprising administering to a patient in need of such treatment a composition of claim 20.

22. A method for screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

- 20       a) exposing a sample comprising a polypeptide of claim 1 to a compound, and  
         b) detecting antagonist activity in the sample.

23. A composition comprising an antagonist compound identified by a method of claim 22 and a pharmaceutically acceptable excipient.

25

24. A method for treating a disease or condition associated with overexpression of functional CYSKP, comprising administering to a patient in need of such treatment a composition of claim 23.

30

25. A method of screening for a compound that specifically binds to the polypeptide of claim 1, said method comprising the steps of:

         a) combining the polypeptide of claim 1 with at least one test compound under suitable conditions, and

35       b) detecting binding of the polypeptide of claim 1 to the test compound, thereby identifying a



compound that specifically binds to the polypeptide of claim 1.

26. A method of screening for a compound that modulates the activity of the polypeptide of claim 1, said method comprising:

- 5           a) combining the polypeptide of claim 1 with at least one test compound under conditions permissive for the activity of the polypeptide of claim 1,
- b) assessing the activity of the polypeptide of claim 1 in the presence of the test compound, and
- c) comparing the activity of the polypeptide of claim 1 in the presence of the test compound with the activity of the polypeptide of claim 1 in the absence of the test compound, wherein a change in
- 10          the activity of the polypeptide of claim 1 in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide of claim 1.

27. A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 5, the method

- 15          comprising:
- a) exposing a sample comprising the target polynucleotide to a compound, under conditions suitable for the expression of the target polynucleotide,
- b) detecting altered expression of the target polynucleotide, and
- c) comparing the expression of the target polynucleotide in the presence of varying amounts of
- 20          the compound and in the absence of the compound.

28. A method for assessing toxicity of a test compound, said method comprising:

- a) treating a biological sample containing nucleic acids with the test compound;
- b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at
- 25          least 20 contiguous nucleotides of a polynucleotide of claim 11 under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a polynucleotide of claim 11 or fragment thereof;
- c) quantifying the amount of hybridization complex; and
- 30          d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

29. A diagnostic test for a condition or disease associated with the expression of CYSKP in a biological sample comprising the steps of:

a) combining the biological sample with an antibody of claim 10, under conditions suitable for the antibody to bind the polypeptide and form an antibody:polypeptide complex; and

5 b) detecting the complex, wherein the presence of the complex correlates with the presence of the polypeptide in the biological sample.

30. The antibody of claim 10, wherein the antibody is:

- 10 a) a chimeric antibody,  
b) a single chain antibody,  
c) a Fab fragment,  
d) a F(ab')<sub>2</sub> fragment, or  
e) a humanized antibody.

15 31. A composition comprising an antibody of claim 10 and an acceptable excipient.

32. A method of diagnosing a condition or disease associated with the expression of CYSKP in a subject, comprising administering to said subject an effective amount of the composition of claim 31.

20

33. A composition of claim 31, wherein the antibody is labeled.

34. A method of diagnosing a condition or disease associated with the expression of CYSKP in a subject, comprising administering to said subject an effective amount of the composition of claim 33.

25

35. A method of preparing a polyclonal antibody with the specificity of the antibody of claim 10 comprising:

- 30 a) immunizing an animal with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34, or an immunogenic fragment thereof, under conditions to elicit an antibody response;  
b) isolating antibodies from said animal; and  
c) screening the isolated antibodies with the polypeptide, thereby identifying a polyclonal antibody which binds specifically to a polypeptide having an amino acid sequence selected from the

group consisting of SEQ ID NO:1-34.

36. An antibody produced by a method of claim 35.

5 37. A composition comprising the antibody of claim 36 and a suitable carrier.

38. A method of making a monoclonal antibody with the specificity of the antibody of claim  
10 comprising:

- 10 a) immunizing an animal with a polypeptide having an amino acid sequence selected from  
the group consisting of SEQ ID NO:1-34, or an immunogenic fragment thereof, under conditions to  
elicit an antibody response;
- b) isolating antibody producing cells from the animal;
- c) fusing the antibody producing cells with immortalized cells to form monoclonal antibody-  
producing hybridoma cells;
- 15 d) culturing the hybridoma cells; and
- e) isolating from the culture monoclonal antibody which binds specifically to a polypeptide  
having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34.

20 39. A monoclonal antibody produced by a method of claim 38.

40. A composition comprising the antibody of claim 39 and a suitable carrier.

41. The antibody of claim 10, wherein the antibody is produced by screening a Fab  
expression library.

25 42. The antibody of claim 10, wherein the antibody is produced by screening a recombinant  
immunoglobulin library.

30 43. A method for detecting a polypeptide having an amino acid sequence selected from the  
group consisting of SEQ ID NO:1-34 in a sample, comprising the steps of:

- a) incubating the antibody of claim 10 with a sample under conditions to allow specific  
binding of the antibody and the polypeptide; and
- b) detecting specific binding, wherein specific binding indicates the presence of a  
polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34 in

the sample.

44. A method of purifying a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34 from a sample, the method comprising:

- 5       a) incubating the antibody of claim 10 with a sample under conditions to allow specific binding of the antibody and the polypeptide; and
- b) separating the antibody from the sample and obtaining the purified polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-34.

10       45. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:1.

46. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:2.

47. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:3.

15

48. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:4.

49. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:5.

20

50. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:6.

51. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:7.

52. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:8.

25

53. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:9.

54. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:10.

30

55. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:11.

56. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:12.

57. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:13.

58. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:14.
59. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:15.
60. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:16.
61. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:17.
62. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:18.
63. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:19.
64. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:20.
65. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:21.
66. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:22.
67. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:23.
68. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:24.
69. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:25.
70. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:26.
71. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:27.
72. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:28.
73. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:29.
74. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:30.

75. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:31.

76. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:32.

5

77. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:33.

78. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:34.

10

79. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:35.

80. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:36.

15

81. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:37.

82. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:38.

20

83. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:39.

84. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:40.

25

85. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:41.

30

86. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:42.

87. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID

NO:43.

88. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
NO:44.

5

89. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
NO:45.

90. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
10 NO:46.

91. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
NO:47.

92. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
15 NO:48.

93. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
NO:49.

20

94. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
NO:50.

95. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
25 NO:51.

96. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
NO:52.

97. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
30 NO:53.

98. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
NO:54.

99. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:55.

5        100. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:56.

101. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:57.

10        102. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:58.

15        103. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:59.

104. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:60.

20        105. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:61.

106. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:62.

25        107. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:63.

30        108. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:64.

109. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:65.



110. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
NO:66.

111. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
5 NO:67.

112. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
NO:68.

<110> INCYTE GENOMICS, INC.  
 YUE, Henry  
 TANG, Y. Tom  
 AU-YOUNG, Janice  
 LU, Dyung Aina M.  
 BAUGHN, Mariah R.  
 HILLMAN, Jennifer L.  
 AZIMZAI, Yalda  
 LAL, Preeti  
 YAO, Monique G.  
 BANDMAN, Olga  
 BURFORD, Neil  
 BATRA, Sajeew  
 KEARNEY, Liam  
 POLICKY, Jennifer L.

<120> CYTOSKELETON-ASSOCIATED PROTEINS

<130> PF-0772 PCT

<140> To Be Assigned

<141> Herewith

<150> 60/201,960; 60/202,729; 60/209,705; 60/210,149; 60/213,215

<151> 2000-05-05; 2000-05-08; 2000-06-05; 2000-06-07; 2000-06-21

<160> 68

<170> PERL Program

<210> 1

<211> 622

<212> PRT

<213> Homo sapiens

<220>

<221> misc\_feature

<223> Incyte ID No: 1889577CD1

<400> 1

Met	Ala	Met	Met	Val	Phe	Pro	Arg	Glu	Glu	Lys	Leu	Ser	Gln	Asp
1				5					10					15
Glu	Ile	Val	Leu	Gly	Thr	Lys	Ala	Val	Ile	Gln	Gly	Leu	Glu	Thr
				20					25					30
Leu	Arg	Gly	Glu	His	Arg	Ala	Leu	Leu	Ala	Pro	Leu	Val	Ala	Pro
				35					40					45
Glu	Ala	Gly	Glu	Pro	Glu	Pro	Gly	Ser	Gln	Glu	Arg	Cys	Ile	Leu
				50					55					60
Leu	Arg	Arg	Ser	Leu	Glu	Ala	Ile	Glu	Leu	Gly	Leu	Gly	Glu	Ala
				65					70					75
Gln	Val	Ile	Leu	Ala	Leu	Ser	Ser	His	Leu	Gly	Ala	Val	Glu	Ser
				80					85					90
Glu	Lys	Gln	Lys	Leu	Arg	Ala	Gln	Val	Arg	Arg	Leu	Val	Gln	Glu
				95					100					105
Asn	Gln	Trp	Leu	Arg	Glu	Glu	Leu	Ala	Gly	Thr	Gln	Gln	Lys	Leu
				110					115					120
Gln	Arg	Ser	Glu	Gln	Ala	Val	Ala	Gln	Leu	Glu	Glu	Glu	Lys	Gln
				125					130					135
His	Leu	Leu	Phe	Met	Ser	Gln	Ile	Arg	Lys	Leu	Asp	Glu	Asp	Ala
				140					145					150
Ser	Pro	Asn	Glu	Glu	Lys	Gly	Asp	Val	Pro	Lys	Asp	Thr	Leu	Asp
				155					160					165
Asp	Leu	Phe	Pro	Asn	Glu	Asp	Glu	Gln	Ser	Pro	Ala	Pro	Ser	Pro
				170					175					180
Gly	Gly	Gly	Asp	Val	Ser	Gly	Gln	His	Gly	Gly	Tyr	Glu	Ile	Pro
				185					190					195
Ala	Arg	Leu	Arg	Thr	Leu	His	Asn	Leu	Val	Ile	Gln	Tyr	Ala	Ser

Gln Gly Arg Tyr	200	Val Ala Val Pro	205	Cys Lys Gln Ala	210
Glu Asp Leu Glu	215	Lys Thr Ser Gly His	220	Asp His Pro Asp Val	225
Thr Met Leu Asn	230	Ile Leu Ala Leu Val	235	Tyr Arg Asp Gln Asn	240
Tyr Lys Glu Ala	245	Ala His Leu Leu Asn	250	Asp Ala Leu Ala Ile	255
Glu Lys Thr Leu	260	Gly Lys Asp His Pro	265	Ala Val Ala Ala Thr	270
Asn Asn Leu Ala	275	Val Leu Tyr Gly Lys	280	Arg Gly Lys Tyr Lys	285
Ala Glu Pro Leu	290	Cys Lys Arg Ala Leu	295	Glu Ile Arg Glu Lys	300
Leu Gly Lys Phe	305	His Pro Asp Val Ala	310	Lys Gln Leu Ser Asn	315
Ala Leu Leu Cys	320	Gln Asn Gln Gly Lys	325	Ala Glu Glu Val Glu	330
Tyr Tyr Arg Arg	335	Ala Leu Glu Ile Tyr	340	Ala Thr Arg Leu Gly	345
Asp Asp Pro Asn	350	Val Ala Lys Thr Lys	355	Asn Asn Leu Ala Ser	360
Tyr Leu Lys Gln	365	Gly Lys Tyr Gln Asp	370	Ala Glu Thr Leu Tyr	375
Glu Ile Leu Thr	380	Arg Ala His Glu Lys	385	Glu Phe Gly Ser Val	390
Gly Asp Asn Lys	395	Pro Ile Trp Met His	400	Ala Glu Glu Arg Glu	405
Ser Lys Asp Lys	410	Arg Arg Asp Ser Ala	415	Pro Tyr Gly Glu Tyr	420
Ser Trp Tyr Lys	425	Ala Cys Lys Val Asp	430	Ser Pro Thr Val Asn	435
Thr Leu Arg Thr	440	Leu Gly Ala Leu Tyr	445	Arg Arg Gln Gly Lys	450
Glu Ala Ala His	455	Thr Leu Glu Asp Cys	460	Ala Ser Arg Asn Arg	465
Gln Gly Leu Asp	470	Pro Ala Ser Gln Thr	475	Lys Val Val Glu Leu	480
Lys Asp Gly Ser	485	Gly Arg Arg Gly Asp	490	Pro Arg Ser Ser Arg	495
Met Ala Gly Gly	500	Ala Gly Pro Arg Ser	505	Glu Ser Asp Leu Glu	510
Val Gly Pro Thr	515	Ala Glu Trp Asn Gly	520	Asp Gly Ser Gly Ser	525
Arg Arg Ser Gly	530	Ser Phe Gly Lys Leu	535	Arg Asp Ala Leu Arg	540
Ser Ser Glu Met	545	Leu Val Lys Lys Leu	550	Gln Gly Gly Thr Pro	555
Glu Pro Pro Asn	560	Pro Arg Met Lys Arg	565	Ala Ser Ser Leu Asn	570
Leu Asn Lys Ser	575	Val Glu Glu Pro Thr	580	Gln Pro Gly Gly Thr	585
Leu Ser Asp Ser	590	Arg Thr Leu Ser Ser	595	Ser Ser Met Asp Leu	600
Arg Arg Ser Ser	605	Val Gly	610		615
	620				

&lt;210&gt; 2

&lt;211&gt; 190

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2427982CD1

&lt;400&gt; 2

Met	Ala	Lys	Ala	Thr	Thr	Ile	Lys	Glu	Ala	Leu	Ala	Arg	Trp	Glu	
1				5					10					15	
Glu	Lys	Thr	Gly	Gln	Arg	Pro	Ser	Glu	Ala	Lys	Glu	Ile	Lys	Leu	
				20					25					30	
Tyr	Ala	Gln	Ile	Pro	Pro	Ile	Glu	Lys	Met	Asp	Ala	Ser	Leu	Ser	
				35					40					45	
Met	Leu	Ala	Asn	Cys	Glu	Lys	Leu	Ser	Leu	Ser	Thr	Asn	Cys	Ile	
				50					55					60	
Glu	Lys	Ile	Ala	Asn	Leu	Asn	Gly	Leu	Lys	Asn	Leu	Arg	Ile	Leu	
				65					70					75	
Ser	Leu	Gly	Arg	Asn	Asn	Ile	Lys	Asn	Leu	Asn	Gly	Leu	Glu	Ala	
				80					85					90	
Val	Gly	Asp	Thr	Leu	Glu	Glu	Leu	Trp	Ile	Ser	Tyr	Asn	Phe	Ile	
				95					100					105	
Glu	Lys	Leu	Lys	Gly	Ile	His	Ile	Met	Lys	Lys	Leu	Lys	Ile	Leu	
				110					115					120	
Tyr	Met	Ser	Asn	Asn	Leu	Val	Lys	Asp	Trp	Ala	Glu	Phe	Val	Lys	
				125					130					135	
Leu	Ala	Glu	Leu	Pro	Cys	Leu	Glu	Asp	Leu	Val	Phe	Val	Gly	Asn	
				140					145					150	
Pro	Leu	Glu	Glu	Lys	His	Ser	Ala	Glu	Asn	Asn	Trp	Ile	Glu	Glu	
				155					160					165	
Ala	Thr	Lys	Arg	Val	Pro	Lys	Leu	Lys	Lys	Leu	Asp	Gly	Thr	Pro	
				170					175					180	
Val	Ile	Lys	Gly	Asp	Glu	Glu	Glu	Asp	Asn						
				185					190						

&lt;210&gt; 3

&lt;211&gt; 331

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2470833CD1

&lt;400&gt; 3

Met	Glu	Pro	Glu	Phe	Leu	Tyr	Asp	Leu	Leu	Gln	Leu	Pro	Lys	Gly	
1				5					10					15	
Val	Glu	Pro	Pro	Ala	Glu	Glu	Glu	Leu	Ser	Lys	Gly	Gly	Lys	Lys	
				20					25					30	
Lys	Tyr	Leu	Pro	Pro	Thr	Ser	Arg	Lys	Asp	Pro	Lys	Phe	Glu	Glu	
				35					40					45	
Leu	Gln	Lys	Val	Leu	Met	Glu	Trp	Ile	Asn	Ala	Thr	Leu	Leu	Pro	
				50					55					60	
Glu	His	Ile	Val	Val	Arg	Ser	Leu	Glu	Glu	Asp	Met	Phe	Asp	Gly	
				65					70					75	
Leu	Ile	Leu	His	His	Leu	Phe	Gln	Arg	Leu	Ala	Ala	Leu	Lys	Leu	
				80					85					90	
Glu	Ala	Glu	Asp	Ile	Ala	Leu	Thr	Ala	Thr	Ser	Gln	Lys	His	Lys	
				95					100					105	
Leu	Thr	Val	Val	Leu	Glu	Ala	Val	Asn	Arg	Ser	Leu	Gln	Leu	Glu	
				110					115					120	
Glu	Trp	Gln	Ala	Lys	Trp	Ser	Val	Glu	Ser	Ile	Phe	Asn	Lys	Asp	
				125					130					135	
Leu	Leu	Ser	Thr	Leu	His	Leu	Leu	Val	Ala	Leu	Ala	Lys	Arg	Phe	
				140					145					150	
Gln	Pro	Asp	Leu	Ser	Leu	Pro	Thr	Asn	Val	Gln	Val	Glu	Val	Ile	
				155					160					165	
Thr	Ile	Glu	Ser	Thr	Lys	Ser	Gly	Leu	Lys	Ser	Glu	Lys	Leu	Val	
				170					175					180	
Glu	Gln	Leu	Thr	Glu	Tyr	Ser	Thr	Asp	Lys	Asp	Glu	Pro	Pro	Lys	
				185					190					195	
Asp	Val	Phe	Asp	Glu	Leu	Phe	Lys	Leu	Ala	Pro	Glu	Lys	Val	Asn	
				200					205					210	
Ala	Val	Lys	Glu	Ala	Ile	Val	Asn	Phe	Val	Asn	Gln	Lys	Leu	Asp	

Arg	Leu	Gly	Leu	215	Ser	Val	Gln	Asn	Leu	220	Asp	Thr	Gln	Phe	Ala	225
				230												240
Gly	Val	Ile	Leu	245	Leu	Leu	Ile	Gly	250	Gln	Leu	Glu	Gly	Phe		255
Leu	His	Leu	Lys	260	Glu	Phe	Tyr	Leu	265	Pro	Asn	Ser	Pro	Ala	Glu	270
Met	Leu	His	Asn	275	Val	Thr	Leu	Ala	280	Glu	Leu	Leu	Lys	Asp	Glu	285
Gly	Leu	Leu	Ser	290	Cys	Pro	Val	Ser	295	Glu	Asp	Ile	Val	Asn	Lys	300
Asp	Ala	Lys	Ser	305	Thr	Leu	Arg	Val	310	Tyr	Gly	Leu	Phe	Cys	Lys	315
His	Thr	Gln	Lys	320	Ala	His	Arg	Asp	325	Thr	Pro	His	Gly	Ala	Pro	330
Asn																

<210> 4  
 <211> 239  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2080579CD1

Met	Met	Glu	Ile	Gln	Met	Asp	Glu	Gly	Gly	Gly	Val	Val	Val	Tyr	
1				5					10					15	
Gln	Asp	Asp	Tyr	Cys	Ser	Gly	Ser	Val	Met	Ser	Glu	Arg	Val	Ser	
				20					25					30	
Gly	Leu	Ala	Gly	Ser	Ile	Tyr	Arg	Glu	Phe	Glu	Arg	Leu	Ile	His	
				35					40					45	
Cys	Tyr	Asp	Glu	Glu	Val	Val	Lys	Glu	Leu	Met	Pro	Leu	Val	Val	
				50					55					60	
Asn	Val	Leu	Glu	Asn	Leu	Asp	Ser	Val	Leu	Ser	Glu	Asn	Gln	Glu	
				65					70					75	
His	Glu	Val	Glu	Leu	Glu	Leu	Leu	Arg	Glu	Asp	Asn	Glu	Gln	Leu	
				80					85					90	
Leu	Thr	Gln	Tyr	Glu	Arg	Glu	Lys	Ala	Leu	Arg	Arg	Gln	Ala	Glu	
				95					100					105	
Glu	Lys	Phe	Ile	Glu	Phe	Glu	Asp	Ala	Leu	Glu	Gln	Glu	Lys	Lys	
				110					115					120	
Glu	Leu	Gln	Ile	Gln	Val	Glu	His	Tyr	Glu	Phe	Gln	Thr	Arg	Gln	
				125					130					135	
Leu	Glu	Leu	Lys	Ala	Lys	Asn	Tyr	Ala	Asp	Gln	Ile	Ser	Arg	Leu	
				140					145					150	
Glu	Glu	Arg	Glu	Ser	Glu	Met	Lys	Lys	Glu	Tyr	Asn	Ala	Leu	His	
				155					160					165	
Gln	Arg	His	Thr	Glu	Met	Ile	Gln	Thr	Tyr	Val	Glu	His	Ile	Glu	
				170					175					180	
Arg	Ser	Lys	Met	Gln	Gln	Val	Gly	Gly	Asn	Ser	Gln	Thr	Glu	Ser	
				185					190					195	
Ser	Leu	Pro	Gly	Arg	Arg	Tyr	Ala	Gly	Arg	Gly	Gly	Val	Glu	Val	
				200					205					210	
Arg	Gly	Ala	Arg	Arg	Gly	Gly	Gly	Thr	Gln	Asp	Ala	Ala	His	Ala	
				215					220					225	
Arg	Val	Val	Val	Leu	Val	Met	Ala	Arg	Ala	Leu	Gly	Ser	Gly		
				230					235						

<210> 5  
 <211> 488  
 <212> PRT  
 <213> Homo sapiens

<220>

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2156553CD1

&lt;400&gt; 5

Met	Asp	Ile	Asp	Lys	Asp	Leu	Glu	Ala	Pro	Leu	Tyr	Leu	Thr	Pro
1				5					10					15
Glu	Gly	Trp	Ser	Leu	Phe	Leu	Gln	Arg	Tyr	Tyr	Gln	Val	Val	His
				20					25					30
Glu	Gly	Ala	Glu	Leu	Arg	His	Leu	Asp	Thr	Gln	Val	Gln	Arg	Cys
				35					40					45
Glu	Asp	Ile	Leu	Gln	Gln	Leu	Gln	Ala	Val	Val	Pro	Gln	Ile	Asp
				50					55					60
Met	Glu	Gly	Asp	Arg	Asn	Ile	Trp	Ile	Val	Lys	Pro	Gly	Ala	Lys
				65					70					75
Ser	Arg	Gly	Arg	Gly	Ile	Met	Cys	Met	Asp	His	Leu	Glu	Glu	Met
				80					85					90
Leu	Lys	Leu	Val	Asn	Gly	Asn	Pro	Val	Val	Met	Lys	Asp	Gly	Lys
				95					100					105
Trp	Val	Val	Gln	Lys	Tyr	Ile	Glu	Arg	Pro	Leu	Leu	Ile	Phe	Gly
				110					115					120
Thr	Lys	Phe	Asp	Leu	Arg	Gln	Trp	Phe	Leu	Val	Thr	Asp	Trp	Asn
				125					130					135
Pro	Leu	Thr	Val	Trp	Phe	Tyr	Arg	Asp	Ser	Tyr	Ile	Arg	Phe	Ser
				140					145					150
Thr	Gln	Pro	Phe	Ser	Leu	Lys	Asn	Leu	Asp	Asn	Ser	Val	His	Leu
				155					160					165
Cys	Asn	Asn	Ser	Ile	Gln	Lys	His	Leu	Glu	Asn	Ser	Cys	His	Arg
				170					175					180
His	Pro	Leu	Leu	Pro	Pro	Asp	Asn	Met	Trp	Ser	Ser	Gln	Arg	Phe
				185					190					195
Gln	Ala	His	Leu	Gln	Glu	Met	Gly	Ala	Pro	Asn	Ala	Trp	Ser	Thr
				200					205					210
Ile	Ile	Val	Pro	Gly	Met	Lys	Asp	Ala	Val	Ile	His	Ala	Leu	Gln
				215					220					225
Thr	Ser	Gln	Asp	Thr	Val	Gln	Cys	Arg	Lys	Ala	Ser	Phe	Glu	Leu
				230					235					240
Tyr	Gly	Ala	Asp	Phe	Val	Phe	Gly	Glu	Asp	Phe	Gln	Pro	Trp	Leu
				245					250					255
Ile	Glu	Ile	Asn	Ala	Ser	Pro	Thr	Met	Ala	Pro	Ser	Thr	Ala	Val
				260					265					270
Thr	Ala	Arg	Leu	Cys	Ala	Gly	Val	Gln	Ala	Asp	Thr	Leu	Arg	Val
				275					280					285
Val	Ile	Asp	Arg	Met	Leu	Asp	Arg	Asn	Cys	Asp	Thr	Gly	Ala	Phe
				290					295					300
Glu	Leu	Ile	Tyr	Lys	Gln	Pro	Ala	Val	Glu	Val	Pro	Gln	Tyr	Val
				305					310					315
Gly	Ile	Arg	Leu	Gly	Val	Glu	Gly	Phe	Thr	Ile	Lys	Lys	Pro	Met
				320					325					330
Ala	Met	Cys	His	Arg	Arg	Met	Gly	Val	Arg	Pro	Ala	Val	Pro	Leu
				335					340					345
Leu	Thr	Gln	Arg	Gly	Ser	Gly	Glu	Gly	Lys	Asp	Ser	Gly	Ile	Pro
				350					355					360
Thr	His	Arg	Ser	Ala	Ser	Arg	Lys	Gly	Thr	Gly	Ala	Arg	Ser	Leu
				365					370					375
Gly	His	Ser	Glu	Lys	Pro	Val	Ser	Thr	Ala	Thr	Thr	Ser	Ala	Pro
				380					385					390
Gly	Lys	Gly	Lys	Lys	Gly	Lys	Ala	Lys	Arg	Ala	Thr	Ala	Leu	Val
				395					400					405
Cys	Pro	Asn	Leu	Trp	Glu	Trp	Asp	Ala	Pro	Ser	Thr	Arg	Met	Gly
				410					415					420
Cys	Ile	Phe	Thr	Met	Thr	Phe	Ser	Ser	Gly	Asp	Arg	Gln	Pro	His
				425					430					435
His	Leu	Asn	Arg	Leu	Pro	Leu	Ser	Pro	Lys	Asn	Pro	Gln	Ala	Leu
				440					445					450
Gly	Lys	Thr	Ile	Pro	Pro	Lys	His	Pro	Ser	Val	Pro	Arg	Arg	Phe
				455					460					465
Ile	Pro	Ala	Leu	Gln	Ala	Pro	Pro	Asn	His	Leu	Asp	Gln	Pro	Pro

470  
 His Gln Arg Ala Thr Ser Ser Lys  
 485

475  
 480

<210> 6  
 <211> 1190  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2182855CD1

<400> 6  
 Met Pro Arg Gly Val Phe Gln Gln Leu Ser Asn Leu Val Leu Gln  
 1 5 10 15  
 Glu Leu Asn Ala Asn Leu Ser Asn Leu Thr Ser Ala Phe Glu Lys  
 20 25 30  
 Ala Thr Ala Glu Lys Ile Lys Cys Gln Glu Ala Asp Ala Thr  
 35 40 45  
 Asn Arg Val Ile Leu Leu Ala Asn Arg Leu Val Gly Gly Leu Ala  
 50 55 60  
 Ser Glu Asn Ile Arg Trp Ala Glu Ser Val Glu Asn Phe Arg Ser  
 65 70 75  
 Gln Gly Val Thr Leu Cys Gly Asp Val Leu Leu Ile Ser Ala Phe  
 80 85 90  
 Val Ser Tyr Val Gly Tyr Phe Thr Lys Lys Tyr Arg Asn Glu Leu  
 95 100 105  
 Met Glu Lys Phe Trp Ile Pro Tyr Ile His Asn Leu Lys Val Pro  
 110 115 120  
 Ile Pro Ile Thr Asn Gly Leu Asp Pro Leu Ser Leu Leu Thr Asp  
 125 130 135  
 Asp Ala Asp Val Ala Thr Trp Asn Asn Gln Gly Leu Pro Ser Asp  
 140 145 150  
 Arg Met Ser Thr Glu Asn Ala Thr Ile Leu Gly Asn Thr Glu Arg  
 155 160 165  
 Trp Pro Leu Ile Val Asp Ala Gln Leu Gln Gly Ile Lys Trp Ile  
 170 175 180  
 Lys Asn Lys Tyr Arg Ser Glu Leu Lys Ala Ile Arg Leu Gly Gln  
 185 190 195  
 Lys Ser Tyr Leu Asp Val Ile Glu Gln Ala Ile Ser Glu Gly Asp  
 200 205 210  
 Thr Leu Leu Ile Glu Asn Ile Gly Glu Thr Val Asp Pro Val Leu  
 215 220 225  
 Asp Pro Leu Leu Gly Arg Asn Thr Ile Lys Lys Gly Lys Tyr Ile  
 230 235 240  
 Lys Ile Gly Asp Lys Glu Val Glu Tyr His Pro Lys Phe Arg Leu  
 245 250 255  
 Ile Leu His Thr Lys Tyr Phe Asn Pro His Tyr Lys Pro Glu Met  
 260 265 270  
 Gln Ala Gln Cys Thr Leu Ile Asn Phe Leu Val Thr Arg Asp Gly  
 275 280 285  
 Leu Glu Asp Gln Leu Leu Ala Ala Val Val Ala Lys Glu Arg Pro  
 290 295 300  
 Asp Leu Glu Gln Leu Lys Ala Asn Leu Thr Lys Ser Gln Asn Glu  
 305 310 315  
 Phe Lys Ile Val Leu Lys Glu Leu Glu Asp Ser Leu Leu Ala Arg  
 320 325 330  
 Leu Ser Ala Ala Ser Gly Asn Phe Leu Gly Asp Thr Ala Leu Val  
 335 340 345  
 Glu Asn Leu Glu Thr Thr Lys His Thr Ala Ser Glu Ile Glu Glu  
 350 355 360  
 Lys Val Val Glu Ala Lys Ile Thr Glu Val Lys Ile Asn Glu Ala  
 365 370 375  
 Arg Glu Asn Tyr Arg Pro Ala Ala Glu Arg Ala Ser Leu Leu Tyr  
 380 385 390  
 Phe Ile Leu Asn Asp Leu Asn Lys Ile Asn Pro Val Tyr Gln Phe

	395		400		405
Ser Leu Lys Ala	Phe Asn Val Val Phe	Glu Lys Ala Ile Gln	Arg		
	410		415		420
Thr Thr Pro Ala	Asn Glu Val Lys Gln	Arg Val Ile Asn Leu	Thr		
	425		430		435
Asp Glu Ile Thr	Tyr Ser Val Tyr Met	Tyr Thr Ala Arg Gly	Leu		
	440		445		450
Phe Glu Arg Asp	Lys Leu Ile Phe Leu	Ala Gln Val Thr Phe	Gln		
	455		460		465
Val Leu Ser Met	Lys Lys Glu Leu Asn	Pro Val Glu Leu Asp	Phe		
	470		475		480
Leu Leu Arg Phe	Pro Phe Lys Ala Gly	Val Val Ser Pro Val	Asp		
	485		490		495
Phe Leu Gln His	Gln Gly Trp Gly Gly	Ile Lys Ala Leu Ser	Glu		
	500		505		510
Met Asp Glu Phe	Lys Asn Leu Asp Ser	Asp Ile Glu Gly Ser	Ala		
	515		520		525
Lys Arg Trp Lys	Lys Leu Val Glu Ser	Glu Ala Pro Glu Lys	Glu		
	530		535		540
Ile Phe Pro Lys	Glu Trp Lys Asn Lys	Thr Ala Leu Gln Lys	Leu		
	545		550		555
Cys Met Val Arg	Cys Leu Arg Pro Asp	Arg Met Thr Tyr Ala	Ile		
	560		565		570
Lys Asn Phe Val	Glu Glu Lys Met Gly	Ser Lys Phe Val Glu	Gly		
	575		580		585
Arg Ser Val Glu	Phe Ser Lys Ser Tyr	Glu Glu Ser Ser Pro	Ser		
	590		595		600
Thr Ser Ile Phe	Phe Ile Leu Ser Pro	Gly Val Asp Pro Leu	Lys		
	605		610		615
Asp Val Glu Ala	Leu Gly Lys Lys Leu	Gly Phe Thr Ile Asp	Asn		
	620		625		630
Gly Lys Leu His	Asn Val Ser Leu Gly	Gln Gly Gln Glu Val	Val		
	635		640		645
Ala Glu Asn Ala	Leu Asp Val Ala Ala	Glu Lys Gly His Trp	Val		
	650		655		660
Ile Leu Gln Asn	Ile His Leu Val Ala	Arg Trp Leu Gly Thr	Leu		
	665		670		675
Asp Lys Lys Leu	Glu Arg Tyr Ser Thr	Gly Ser His Glu Asp	Tyr		
	680		685		690
Arg Val Phe Ile	Ser Ala Glu Pro Ala	Pro Ser Pro Glu Thr	His		
	695		700		705
Ile Ile Pro Gln	Gly Ile Leu Glu Asn	Ala Ile Lys Ile Thr	Asn		
	710		715		720
Glu Pro Pro Thr	Gly Met Tyr Ala Asn	Leu His Lys Ala Leu	Asp		
	725		730		735
Leu Phe Thr Gln	Asp Thr Leu Glu Met	Cys Thr Lys Glu Met	Glu		
	740		745		750
Phe Lys Cys Met	Leu Phe Ala Leu Cys	Tyr Phe His Ala Val	Val		
	755		760		765
Ala Glu Arg Arg	Lys Phe Gly Ala Gln	Gly Trp Asn Arg Ser	Tyr		
	770		775		780
Pro Phe Asn Asn	Gly Asp Leu Thr Ile	Ser Ile Asn Val Leu	Tyr		
	785		790		795
Asn Tyr Leu Glu	Ala Asn Pro Lys Val	Pro Trp Asp Asp Leu	Arg		
	800		805		810
Tyr Leu Phe Gly	Glu Ile Met Tyr Gly	Gly His Ile Thr Asp	Asp		
	815		820		825
Trp Asp Arg Arg	Leu Cys Arg Thr Tyr	Leu Ala Glu Tyr Ile	Arg		
	830		835		840
Thr Glu Met Leu	Glu Gly Asp Val Leu	Leu Ala Pro Gly Phe	Gln		
	845		850		855
Ile Pro Pro Asn	Leu Asp Tyr Lys Gly	Tyr His Glu Tyr Ile	Asp		
	860		865		870
Glu Asn Leu Pro	Pro Glu Ser Pro Tyr	Leu Tyr Gly Leu His	Pro		
	875		880		885
Asn Ala Glu Ile	Gly Phe Leu Thr Val	Thr Ser Glu Lys Leu	Phe		
	890		895		900



```

Arg Thr Val Leu Glu Met Gln Pro Lys Glu Thr Asp Ser Gly Ala
      905      910
Gly Thr Gly Val Ser Arg Glu Glu Lys Val Lys Ala Val Leu Asp
      920      925
Asp Ile Leu Glu Lys Ile Pro Glu Thr Phe Asn Met Ala Glu Ile
      935      940
Met Ala Lys Ala Ala Glu Lys Thr Pro Tyr Val Val Val Ala Phe
      950      955
Gln Glu Cys Glu Arg Met Asn Ile Leu Thr Asn Glu Met Arg Arg
      965      970
Ser Leu Lys Glu Leu Asn Leu Gly Leu Lys Gly Glu Leu Thr Ile
      980      985
Thr Thr Asp Val Glu Asp Leu Ser Thr Ala Leu Phe Tyr Asp Thr
      995      1000
Val Pro Asp Thr Trp Val Ala Arg Ala Tyr Pro Ser Met Met Gly
      1010      1015
Leu Ala Ala Trp Tyr Ala Asp Leu Leu Leu Arg Ile Arg Glu Leu
      1025      1030
Glu Ala Trp Thr Thr Asp Phe Ala Leu Pro Thr Thr Val Trp Leu
      1040      1045
Ala Gly Phe Phe Asn Pro Gln Ser Phe Leu Thr Ala Ile Met Gln
      1055      1060
Ser Met Ala Arg Lys Asn Glu Trp Pro Leu Asp Lys Met Cys Leu
      1070      1075
Ser Val Glu Val Thr Lys Lys Asn Arg Glu Asp Met Thr Ala Pro
      1085      1090
Pro Arg Glu Gly Ser Tyr Val Tyr Gly Leu Phe Met Glu Gly Ala
      1100      1105
Arg Trp Asp Thr Gln Thr Gly Val Ile Ala Glu Ala Arg Leu Lys
      1115      1120
Glu Leu Thr Pro Ala Met Pro Val Ile Phe Ile Lys Ala Ile Pro
      1130      1135
Val Asp Arg Met Glu Thr Lys Asn Ile Tyr Glu Cys Pro Val Tyr
      1145      1150
Lys Thr Arg Ile Arg Gly Pro Thr Tyr Val Trp Thr Phe Asn Leu
      1160      1165
Lys Thr Lys Glu Lys Ala Ala Lys Trp Ile Leu Ala Ala Val Ala
      1175      1180
Leu Leu Leu Gln Val
      1190

```

```

<210> 7
<211> 270
<212> PRT
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 2242106CD1

```

```

<400> 7
Met Leu Leu Thr Gln Val Val Trp Leu Pro Glu Pro Gly His Ser
  1      5      10
His Arg Phe Gln Val Leu Ser Val Ala Thr Asp Gly Lys Val Leu
      20      25
Leu Trp Gln Gly Ile Gly Val Gly Gln Leu Gln Leu Thr Glu Gly
      35      40
Phe Ala Leu Val Met Gln Gln Leu Pro Arg Ser Thr Lys Leu Lys
      50      55
Lys His Pro Arg Gly Glu Thr Glu Val Gly Ala Thr Ala Val Ala
      65      70
Phe Ser Ser Phe Asp Pro Arg Leu Phe Ile Leu Gly Thr Glu Gly
      80      85
Gly Phe Pro Leu Lys Cys Ser Leu Ala Ala Gly Glu Ala Ala Leu
      95      100
Thr Arg Met Pro Ser Ser Val Pro Leu Arg Ala Pro Ala Gln Phe
      110      115

```

Thr	Phe	Ser	Pro	His	Gly	Gly	Pro	Ile	Tyr	Ser	Val	Ser	Cys	Ser
				125					130					135
Pro	Phe	His	Arg	Asn	Leu	Phe	Leu	Ser	Ala	Gly	Thr	Asp	Gly	His
				140					145					150
Val	His	Leu	Tyr	Ser	Met	Leu	Gln	Ala	Pro	Pro	Leu	Thr	Ser	Leu
				155					160					165
Gln	Leu	Ser	Leu	Lys	Tyr	Leu	Phe	Ala	Val	Arg	Trp	Ser	Pro	Val
				170					175					180
Arg	Pro	Leu	Val	Phe	Ala	Ala	Ala	Ser	Gly	Lys	Gly	Asp	Val	Gln
				185					190					195
Leu	Phe	Asp	Leu	Gln	Lys	Ser	Ser	Gln	Lys	Pro	Thr	Val	Leu	Ile
				200					205					210
Lys	Gln	Thr	Gln	Asp	Glu	Ser	Pro	Val	Tyr	Cys	Leu	Glu	Phe	Asn
				215					220					225
Ser	Gln	Gln	Thr	Gln	Leu	Leu	Ala	Ala	Gly	Asp	Ala	Gln	Gly	Thr
				230					235					240
Val	Lys	Val	Trp	Gln	Leu	Ser	Thr	Glu	Phe	Thr	Glu	Gln	Gly	Pro
				245					250					255
Arg	Glu	Ala	Glu	Asp	Leu	Asp	Cys	Leu	Ala	Ala	Glu	Val	Ala	Ala
				260					265					270

&lt;210&gt; 8

&lt;211&gt; 647

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2726877CD1

&lt;400&gt; 8

Met	Asp	Leu	Leu	Lys	Asn	Pro	Lys	Ile	Ala	Asp	Tyr	Leu	Thr	Arg
1				5					10					15
Tyr	Glu	His	Phe	Ser	Ser	Cys	Leu	His	Gln	Val	Leu	Gly	Leu	Leu
				20					25					30
Asn	Gly	Lys	Asp	Pro	Asp	Ser	Ser	Ser	Lys	Val	Leu	Glu	Leu	Leu
				35					40					45
Leu	Ala	Phe	Cys	Ser	Val	Thr	Gln	Leu	Arg	His	Met	Leu	Thr	Gln
				50					55					60
Met	Met	Phe	Glu	Gln	Ser	Pro	Pro	Gly	Ser	Ala	Thr	Leu	Gly	Ser
				65					70					75
His	Thr	Lys	Cys	Leu	Glu	Pro	Thr	Val	Ala	Leu	Leu	Arg	Trp	Leu
				80					85					90
Ser	Gln	Pro	Leu	Asp	Gly	Ser	Glu	Asn	Cys	Ser	Val	Leu	Ala	Leu
				95					100					105
Glu	Leu	Phe	Lys	Glu	Ile	Phe	Glu	Asp	Val	Ile	Asp	Ala	Ala	Asn
				110					115					120
Cys	Ser	Ser	Ala	Asp	Arg	Phe	Val	Thr	Leu	Leu	Leu	Pro	Thr	Ile
				125					130					135
Leu	Asp	Gln	Leu	Gln	Phe	Thr	Glu	Gln	Asn	Leu	Asp	Glu	Ala	Leu
				140					145					150
Thr	Arg	Gln	Lys	Cys	Glu	Arg	Ile	Ala	Lys	Ala	Phe	Glu	Val	Leu
				155					160					165
Leu	Thr	Leu	Cys	Gly	Asp	Asp	Thr	Leu	Lys	Met	His	Ile	Ala	Lys
				170					175					180
Ile	Leu	Thr	Thr	Val	Lys	Cys	Thr	Thr	Leu	Ile	Glu	Gln	Gln	Phe
				185					190					195
Thr	Tyr	Gly	Lys	Ile	Asp	Leu	Gly	Phe	Gly	Thr	Lys	Val	Ala	Asp
				200					205					210
Ser	Glu	Leu	Cys	Lys	Leu	Ala	Ala	Asp	Val	Ile	Leu	Lys	Thr	Leu
				215					220					225
Asp	Leu	Ile	Asn	Lys	Leu	Lys	Pro	Leu	Val	Pro	Gly	Met	Glu	Val
				230					235					240
Ser	Phe	Tyr	Lys	Ile	Leu	Gln	Asp	Pro	Arg	Leu	Ile	Thr	Pro	Leu
				245					250					255
Ala	Phe	Ala	Leu	Thr	Ser	Asp	Asn	Arg	Glu	Gln	Val	Gln	Ser	Gly

Leu Arg Ile Leu	260	Leu Glu Ala Ala Pro	265	Leu Pro Asp Phe Pro	270
	275		280		285
Leu Val Leu Gly	290	Glu Ser Ile Ala Ala	295	Asn Asn Ala Tyr Arg	300
Gln Glu Thr Glu	305	His Ile Pro Arg Lys	310	Met Pro Trp Gln Ser	315
Asn His Ser Phe	320	Pro Thr Ser Ile Lys	325	Cys Leu Thr Pro His	330
Lys Asp Gly Val	335	Pro Gly Leu Asn Ile	340	Glu Glu Leu Ile Glu	345
Leu Gln Ser Gly	350	Met Val Val Lys Asp	355	Gln Ile Cys Asp Val	360
Ile Ser Asp Ile	365	Met Asp Val Tyr Glu	370	Met Lys Leu Ser Thr	375
Ala Ser Lys Glu	380	Ser Arg Leu Gln Asp	385	Leu Leu Glu Thr Lys	390
Leu Ala Leu Ala	395	Gln Ala Asp Arg Leu	400	Ile Ala Gln His Arg	405
Gln Arg Thr Gln	410	Ala Glu Thr Glu Ala	415	Arg Thr Leu Ala Ser	420
Leu Arg Glu Val	425	Glu Arg Lys Asn Glu	430	Glu Leu Ser Val Leu	435
Lys Ala Gln Gln	440	Val Glu Ser Glu Arg	445	Ala Gln Ser Asp Ile	450
His Leu Phe Gln	455	His Asn Arg Lys Leu	460	Glu Ser Val Ala Glu	465
His Glu Ile Leu	470	Thr Lys Ser Tyr Met	475	Glu Leu Leu Gln Arg	480
Glu Ser Thr Glu	485	Lys Lys Asn Lys Asp	490	Leu Gln Ile Thr Cys	495
Ser Leu Asn Lys	500	Gln Ile Glu Thr Val	505	Lys Lys Leu Asn Glu	510
Leu Lys Glu Gln	515	Asn Glu Lys Ser Ile	520	Ala Gln Leu Ile Glu	525
Glu Glu Gln Arg	530	Lys Glu Val Gln Asn	535	Gln Leu Val Asp Arg	540
His Lys Leu Ala	545	Asn Leu His Gln Lys	550	Thr Lys Val Gln Glu	555
Lys Ile Lys Thr	560	Leu Gln Lys Glu Arg	565	Glu Asp Lys Glu Glu	570
Ile Asp Ile Leu	575	Arg Lys Glu Leu Ser	580	Arg Thr Glu Gln Ile	585
Lys Glu Leu Ser	590	Ile Lys Ala Ser Ser	595	Leu Glu Val Gln Lys	600
Gln Leu Glu Gly	605	Arg Leu Glu Glu Lys	610	Glu Ser Leu Val Lys	615
Gln Gln Glu Glu	620	Leu Asn Lys His Ser	625	His Met Ile Ala Met	630
His Ser Leu Ser	635	Gly Gly Lys Ile Asn	640	Pro Glu Thr Val Asn	645
Ser Ile					

<210> 9  
 <211> 1086  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2738233CD1

<400> 9  
 Met Arg Cys Lys Glu Leu Glu Asn Ala Val Gly Ser Trp Thr Asp  
 1 5 10 15  
 Asp Leu Thr Gln Leu Ser Leu Leu Lys Asp Thr Leu Ser Ala Tyr

Ile Ser Ala Asp	20	Ile Ser Ile Leu	25	Glu Arg Val Glu	30
Asp	35	Asn	40	Leu	45
Leu Gln Arg Gln Trp	50	Glu Glu Leu Cys	55	Gln Leu Ser Leu Arg	60
Arg Gln Gln Ile Gly	65	Glu Arg Leu Asn	70	Trp Ala Val Phe Ser	75
Glu Lys Asn Lys Glu	80	Leu Cys Glu Trp	85	Thr Gln Met Glu Ser	90
Lys Val Ser Gln Asn	95	Gly Asp Ile Leu	100	Ile Glu Glu Met Ile Glu	105
Lys Leu Lys Lys Asp	110	Tyr Gln Glu Glu	115	Ile Ala Ile Ala Gln Glu	120
Asn Lys Ile Gln Leu	125	Gln Gln Met Gly	130	Arg Leu Ala Lys Ala	135
Ser His Glu Ser Lys	140	Ala Ser Glu Ile	145	Glu Tyr Lys Leu Gly Lys	150
Val Asn Asp Arg Trp	155	Gln His Leu Leu	160	Leu Ile Ala Ala Arg	165
Val Lys Lys Leu Lys	170	Glu Thr Leu Val	175	Ala Val Gln Gln Leu Asp	180
Lys Asn Met Ser Ser	185	Leu Arg Thr Trp	190	Leu Ala His Ile Glu Ser	195
Glu Leu Ala Lys Pro	200	Ile Val Tyr Gly	205	Ser Cys Asn Ser Glu Glu	210
Ile Gln Arg Lys Leu	215	Asn Glu Gln Gln	220	Glu Leu Gln Arg Asp Ile	225
Glu Lys His Ser Thr	230	Gly Val Ala Ser	235	Val Leu Asn Leu Cys Glu	240
Val Leu Leu His Asp	245	Cys Asp Ala Cys	250	Ala Thr Asp Ala Glu Cys	255
Asp Ser Ile Gln Gln	260	Ala Thr Arg Asn	265	Leu Asp Arg Arg Trp Arg	270
Asn Ile Cys Ala Met	275	Ser Met Glu Arg	280	Arg Leu Lys Ile Glu Glu	285
Thr Trp Arg Leu Trp	290	Gln Lys Phe Leu	295	Asp Asp Tyr Ser Arg Phe	300
Glu Asp Trp Leu Lys	305	Ser Ser Glu Arg	310	Thr Ala Ala Phe Pro Ser	315
Ser Ser Gly Val Ile	320	Tyr Thr Val Ala	325	Lys Glu Glu Leu Lys Lys	330
Phe Glu Ala Phe Gln	335	Arg Gln Val His	340	Glu Cys Leu Thr Gln Leu	345
Glu Leu Ile Asn Lys	350	Gln Tyr Arg Arg	355	Leu Ala Arg Glu Asn Arg	360
Thr Asp Ser Ala Cys	365	Ser Leu Lys Gln	370	Met Val His Glu Gly Asn	375
Gln Arg Trp Asp Asn	380	Leu Gln Lys Arg	385	Val Thr Ser Ile Leu Arg	390
Arg Leu Lys His Phe	395	Ile Gly Gln Arg	400	Glu Glu Phe Glu Thr Ala	405
Arg Asp Ser Ile Leu	410	Val Trp Leu Thr	415	Glu Met Asp Leu Gln Leu	420
Thr Asn Ile Glu His	425	Phe Ser Glu Cys	430	Asp Val Gln Ala Lys Ile	435
Lys Gln Leu Lys Ala	440	Phe Gln Gln Glu	445	Ile Ser Leu Asn His Asn	450
Lys Ile Glu Gln Ile	455	Ile Ala Gln Gly	460	Glu Gln Leu Ile Glu Lys	465
Ser Glu Pro Leu Asp	470	Ala Ala Ile Ile	475	Glu Glu Glu Leu Asp Glu	480
Leu Arg Arg Tyr Cys	485	Gln Glu Val Phe	490	Gly Arg Val Glu Arg Tyr	495
His Lys Lys Leu Ile	500	Arg Leu Pro Leu	505	Pro Asp Asp Glu His Asp	510
Leu Ser Asp Arg Glu	515	Leu Glu Leu Glu	520	Asp Ser Ala Ala Leu Ser	525

Asp	Leu	His	Trp	His	Asp	Arg	Ser	Ala	Asp	Ser	Leu	Leu	Ser	Pro
				530					535					540
Gln	Pro	Ser	Ser	Asn	Leu	Ser	Leu	Ser	Leu	Ala	Gln	Pro	Leu	Arg
				545					550					555
Ser	Glu	Arg	Ser	Gly	Arg	Asp	Thr	Pro	Ala	Ser	Val	Asp	Ser	Ile
				560					565					570
Pro	Leu	Glu	Trp	Asp	His	Asp	Tyr	Asp	Leu	Ser	Arg	Asp	Leu	Glu
				575					580					585
Ser	Ala	Met	Ser	Arg	Ala	Leu	Pro	Ser	Glu	Asp	Glu	Glu	Gly	Gln
				590					595					600
Asp	Asp	Lys	Asp	Phe	Tyr	Leu	Arg	Gly	Ala	Val	Gly	Leu	Ser	Gly
				605					610					615
Asp	His	Ser	Ala	Leu	Glu	Ser	Gln	Ile	Arg	Gln	Leu	Gly	Lys	Ala
				620					625					630
Leu	Asp	Asp	Ser	Arg	Phe	Gln	Ile	Gln	Gln	Thr	Glu	Asn	Ile	Ile
				635					640					645
Arg	Ser	Lys	Thr	Pro	Thr	Gly	Pro	Glu	Leu	Asp	Thr	Ser	Tyr	Lys
				650					655					660
Gly	Tyr	Met	Lys	Leu	Leu	Gly	Glu	Cys	Ser	Ser	Ser	Ile	Asp	Ser
				665					670					675
Val	Lys	Arg	Leu	Glu	His	Lys	Leu	Lys	Glu	Glu	Glu	Glu	Ser	Leu
				680					685					690
Pro	Gly	Phe	Val	Asn	Leu	His	Ser	Thr	Glu	Thr	Gln	Thr	Ala	Gly
				695					700					705
Val	Ile	Asp	Arg	Trp	Glu	Leu	Leu	Gln	Ala	Gln	Ala	Leu	Ser	Lys
				710					715					720
Glu	Leu	Arg	Met	Lys	Gln	Asn	Leu	Gln	Lys	Trp	Gln	Gln	Phe	Asn
				725					730					735
Ser	Asp	Leu	Asn	Ser	Ile	Trp	Ala	Trp	Leu	Gly	Asp	Thr	Glu	Glu
				740					745					750
Glu	Leu	Glu	Gln	Leu	Gln	Arg	Leu	Glu	Leu	Ser	Thr	Asp	Ile	Gln
				755					760					765
Thr	Ile	Glu	Leu	Gln	Ile	Lys	Lys	Leu	Lys	Glu	Leu	Gln	Lys	Ala
				770					775					780
Val	Asp	His	Arg	Lys	Ala	Ile	Ile	Leu	Ser	Ile	Asn	Leu	Cys	Ser
				785					790					795
Pro	Glu	Phe	Thr	Gln	Ala	Asp	Ser	Lys	Glu	Ser	Arg	Asp	Leu	Gln
				800					805					810
Asp	Arg	Leu	Ser	Gln	Met	Asn	Gly	Arg	Trp	Asp	Arg	Val	Cys	Ser
				815					820					825
Leu	Leu	Glu	Glu	Trp	Arg	Gly	Leu	Leu	Gln	Asp	Ala	Leu	Met	Gln
				830					835					840
Cys	Gln	Gly	Phe	His	Glu	Met	Ser	His	Gly	Leu	Leu	Leu	Met	Leu
				845					850					855
Glu	Asn	Ile	Asp	Arg	Arg	Lys	Asn	Glu	Ile	Val	Pro	Ile	Asp	Ser
				860					865					870
Asn	Leu	Asp	Ala	Glu	Ile	Leu	Gln	Asp	His	His	Lys	Gln	Leu	Met
				875					880					885
Gln	Ile	Lys	His	Glu	Leu	Leu	Glu	Ser	Gln	Leu	Arg	Val	Ala	Ser
				890					895					900
Leu	Gln	Asp	Met	Ser	Cys	Gln	Leu	Leu	Val	Asn	Ala	Glu	Gly	Thr
				905					910					915
Asp	Cys	Leu	Glu	Ala	Lys	Glu	Lys	Val	His	Val	Ile	Gly	Asn	Arg
				920					925					930
Leu	Lys	Leu	Leu	Leu	Lys	Glu	Val	Ser	Arg	His	Ile	Lys	Glu	Leu
				935					940					945
Glu	Lys	Leu	Leu	Asp	Val	Ser	Ser	Ser	Gln	Gln	Asp	Leu	Ser	Ser
				950					955					960
Trp	Ser	Ser	Ala	Asp	Glu	Leu	Asp	Thr	Ser	Gly	Ser	Val	Ser	Pro
				965					970					975
Thr	Ser	Gly	Arg	Ser	Thr	Pro	Asn	Arg	Gln	Lys	Thr	Pro	Arg	Gly
				980					985					990
Lys	Cys	Ser	Leu	Ser	Gln	Pro	Gly	Pro	Ser	Val	Ser	Ser	Pro	His
				995					1000					1005
Ser	Arg	Ser	Thr	Lys	Gly	Gly	Ser	Asp	Ser	Ser	Leu	Ser	Glu	Pro
				1010					1015					1020
Gly	Pro	Gly	Arg	Ser	Gly	Arg	Gly	Phe	Leu	Phe	Arg	Val	Leu	Arg

1025	1030	1035
Ala Ala Leu Pro Leu Gln Leu Leu Leu Leu Leu Leu Ile Gly Leu		
1040	1045	1050
Ala Cys Leu Val Pro Met Ser Glu Glu Asp Tyr Ser Cys Ala Leu		
1055	1060	1065
Ser Asn Asn Phe Ala Arg Ser Phe His Pro Met Leu Arg Tyr Thr		
1070	1075	1080
Asn Gly Pro Pro Pro Leu		
1085		

&lt;210&gt; 10

&lt;211&gt; 396

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1833116CD1

&lt;400&gt; 10

Met Thr Thr Leu Val Leu Asp Asn Gly Ala Tyr Asn Ala Lys Ile		
1	5	10
Gly Tyr Ser His Glu Asn Val Ser Val Ile Pro Asn Cys Gln Phe		15
	20	25
Arg Ser Lys Thr Ala Arg Leu Lys Thr Phe Thr Ala Asn Gln Ile		30
	35	40
Asp Glu Ile Lys Asp Pro Ser Gly Leu Phe Tyr Ile Leu Pro Phe		45
	50	55
Gln Lys Gly Tyr Leu Val Asn Trp Asp Val Gln Arg Gln Val Trp		60
	65	70
Asp Tyr Leu Phe Gly Lys Glu Met Tyr Gln Val Asp Phe Leu Asp		75
	80	85
Thr Asn Ile Ile Ile Thr Glu Pro Tyr Phe Asn Phe Thr Ser Ile		90
	95	100
Gln Glu Ser Met Asn Glu Ile Leu Phe Glu Glu Tyr Gln Phe Gln		105
	110	115
Ala Val Leu Arg Val Asn Ala Gly Ala Leu Ser Ala His Arg Tyr		120
	125	130
Phe Arg Asp Asn Pro Ser Glu Leu Cys Cys Ile Ile Val Asp Ser		135
	140	145
Gly Tyr Ser Phe Thr His Ile Val Pro Tyr Cys Arg Ser Lys Lys		150
	155	160
Lys Lys Glu Ala Ile Ile Arg Ile Asn Val Gly Gly Lys Leu Leu		165
	170	175
Thr Asn His Leu Lys Glu Ile Ile Ser Tyr Arg Gln Leu His Val		180
	185	190
Met Asp Glu Thr His Val Ile Asn Gln Val Lys Glu Asp Val Cys		195
	200	205
Tyr Val Ser Gln Asp Phe Tyr Arg Asp Met Asp Ile Ala Lys Leu		210
	215	220
Lys Gly Glu Glu Asn Thr Val Met Ile Asp Tyr Val Leu Pro Asp		225
	230	235
Phe Ser Thr Ile Lys Lys Gly Phe Cys Lys Pro Arg Glu Glu Met		240
	245	250
Val Leu Ser Gly Lys Tyr Lys Ser Gly Glu Gln Ile Leu Arg Leu		255
	260	265
Ala Asn Glu Arg Phe Ala Val Pro Glu Ile Leu Phe Asn Pro Ser		270
	275	280
Asp Ile Gly Ile Gln Glu Met Gly Ile Pro Glu Ala Ile Val Tyr		285
	290	295
Ser Ile Gln Asn Leu Pro Glu Glu Met Gln Pro His Phe Phe Lys		300
	305	310
Asn Ile Val Leu Thr Gly Gly Asn Ser Leu Phe Pro Gly Phe Arg		315
	320	325
Asp Arg Val Tyr Ser Glu Val Arg Cys Leu Thr Pro Thr Asp Tyr		330
	335	340
Asp Val Ser Val Val Leu Pro Glu Asn Pro Ile Thr Tyr Ala Trp		345

Glu Gly Gly Lys	350	Ile Ser Glu Asn	355	Asp Asp Phe Glu Asp	360
Val Val Thr Arg	365	Glu Asp Tyr Glu Glu	370	Asn Gly His Ser Val	375
Glu Glu Lys Phe	380	Asp Ile	385		390
	395				

<210> 11  
 <211> 304  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 001799CD1

<400> 11

Met Ala Ser Glu Thr	His Asn Val Lys Lys	Arg Asn Phe Cys Asn
1 5	10	15
Lys Ile Glu Asp His	Phe Ile Asp Leu Pro	Arg Lys Lys Ile Ser
20	25	30
Asn Phe Thr Asn Lys	Asn Met Lys Glu Val	Lys Lys Ser Pro Lys
35	40	45
Gln Leu Ala Ala Tyr	Ile Asn Arg Thr Val	Gly Gln Thr Val Lys
50	55	60
Ser Pro Asp Lys Leu	Arg Lys Val Ile Tyr	Arg Arg Lys Lys Val
65	70	75
His His Pro Phe Pro	Asn Pro Cys Tyr Arg	Lys Lys Gln Ser Pro
80	85	90
Gly Ser Gly Gly Cys	Asp Met Ala Asn Lys	Glu Asn Glu Leu Ala
95	100	105
Cys Ala Gly His Leu	Pro Glu Lys Leu His	His Asp Ser Arg Thr
110	115	120
Tyr Leu Val Asn Ser	Ser Asp Ser Gly Ser	Ser Gln Thr Glu Ser
125	130	135
Pro Ser Ser Lys Tyr	Ser Gly Phe Phe Ser	Glu Val Ser Gln Asp
140	145	150
His Glu Thr Met Ala	Gln Val Leu Phe Ser	Arg Asn Met Arg Leu
155	160	165
Asn Val Ala Leu Thr	Phe Trp Arg Lys Arg	Ser Ile Ser Glu Leu
170	175	180
Val Ala Tyr Leu Leu	Arg Ile Glu Asp Leu	Gly Val Val Val Asp
185	190	195
Cys Leu Pro Val Leu	Thr Asn Cys Leu Gln	Glu Glu Lys Gln Tyr
200	205	210
Ile Ser Leu Gly Cys	Cys Val Asp Leu Leu	Pro Leu Val Lys Ser
215	220	225
Leu Leu Lys Ser Lys	Phe Glu Glu Tyr Val	Ile Val Gly Leu Asn
230	235	240
Trp Leu Gln Ala Val	Ile Lys Arg Trp Trp	Ser Glu Leu Ser Ser
245	250	255
Lys Thr Glu Ile Ile	Asn Asp Gly Asn Ile	Gln Ile Leu Lys Gln
260	265	270
Gln Leu Ser Gly Leu	Trp Glu Gln Glu Asn	His Leu Thr Leu Val
275	280	285
Pro Gly Tyr Thr Gly	Asn Ile Ala Lys Asp	Val Asp Ala Tyr Leu
290	295	300

Leu Gln Leu His

<210> 12  
 <211> 201  
 <212> PRT  
 <213> Homo sapiens

<220>

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 119814CD1

&lt;400&gt; 12

```

Met Ile Val Ser Glu Lys Gly Leu His Ser Leu Ile Phe Glu Val
 1          5          10          15
Val Arg Ala Ser Asp Ala Gly Ala Tyr Ala Cys Val Ala Lys Asn
          20          25          30
Arg Ala Gly Glu Ala Thr Phe Thr Val Gln Leu Asp Val Leu Ala
          35          40          45
Lys Glu His Lys Arg Ala Pro Met Phe Ile Tyr Lys Pro Gln Ser
          50          55          60
Lys Lys Val Leu Glu Gly Asp Ser Val Lys Leu Glu Cys Gln Ile
          65          70          75
Ser Ala Ile Pro Pro Pro Lys Leu Phe Trp Lys Arg Asn Asn Glu
          80          85          90
Met Val Gln Phe Asn Thr Asp Arg Ile Ser Leu Tyr Gln Asp Asn
          95          100          105
Thr Gly Arg Val Thr Leu Leu Ile Lys Asp Val Asn Lys Lys Asp
          110          115          120
Ala Gly Trp Tyr Thr Val Ser Ala Val Asn Glu Ala Gly Val Thr
          125          130          135
Thr Cys Asn Thr Arg Leu Asp Val Thr Ala Arg Pro Asn Gln Thr
          140          145          150
Leu Pro Ala Pro Lys Gln Leu Arg Val Arg Pro Thr Phe Ser Lys
          155          160          165
Tyr Leu Ala Leu Asn Gly Lys Gly Leu Asn Val Lys Gln Ala Phe
          170          175          180
Asn Pro Glu Gly Glu Phe Gln Arg Leu Ala Ala Gln Ser Gly Leu
          185          190          195
Tyr Glu Ser Glu Glu Leu
          200

```

&lt;210&gt; 13

&lt;211&gt; 547

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1295420CD1

&lt;400&gt; 13

```

Met Thr Lys Thr Asp Pro Ala Pro Met Ala Pro Pro Pro Arg Gly
 1          5          10          15
Glu Glu Glu Glu Glu Glu Glu Glu Asp Glu Pro Val Pro Glu Ala
          20          25          30
Pro Ser Pro Thr Gln Glu Arg Arg Gln Lys Pro Val Val His Pro
          35          40          45
Ser Ala Pro Ala Pro Leu Pro Lys Asp Tyr Ala Phe Thr Phe Phe
          50          55          60
Asp Pro Asn Asp Pro Ala Cys Gln Glu Ile Leu Phe Asp Pro Gln
          65          70          75
Thr Thr Ile Pro Glu Leu Phe Ala Ile Val Arg Gln Trp Val Pro
          80          85          90
Gln Val Gln His Lys Ile Asp Val Ile Gly Asn Glu Ile Leu Arg
          95          100          105
Arg Gly Cys His Val Asn Asp Arg Asp Gly Leu Thr Asp Met Thr
          110          115          120
Leu Leu His Tyr Ala Cys Lys Ala Gly Ala His Gly Val Gly Asp
          125          130          135
Pro Ala Ala Ala Val Arg Leu Ser Gln Gln Leu Leu Ala Leu Gly
          140          145          150
Ala Asp Val Thr Leu Arg Ser Arg Trp Thr Asn Met Asn Ala Leu
          155          160          165
His Tyr Ala Ala Tyr Phe Asp Val Pro Asp Leu Val Arg Val Leu
          170          175          180

```



Leu	Lys	Gly	Ala	Arg	Pro	Arg	Val	Val	Asn	Ser	Thr	Cys	Ser	Asp
				185					190					195
Phe	Asn	His	Gly	Ser	Ala	Leu	His	Ile	Ala	Ala	Ser	Ser	Leu	Cys
				200					205					210
Leu	Gly	Ala	Ala	Lys	Cys	Leu	Leu	Glu	His	Gly	Ala	Asn	Pro	Ala
				215					220					225
Leu	Arg	Asn	Arg	Lys	Gly	Gln	Val	Pro	Ala	Glu	Val	Val	Pro	Asp
				230					235					240
Pro	Met	Asp	Met	Ser	Leu	Asp	Lys	Ala	Glu	Ala	Ala	Leu	Val	Ala
				245					250					255
Lys	Glu	Leu	Arg	Thr	Leu	Leu	Glu	Glu	Ala	Val	Pro	Leu	Ser	Cys
				260					265					270
Ala	Leu	Pro	Lys	Val	Thr	Leu	Pro	Asn	Tyr	Asp	Asn	Val	Pro	Gly
				275					280					285
Asn	Leu	Met	Leu	Ser	Ala	Leu	Gly	Leu	Arg	Leu	Gly	Asp	Arg	Val
				290					295					300
Leu	Leu	Asp	Gly	Gln	Lys	Thr	Gly	Thr	Leu	Arg	Phe	Cys	Gly	Thr
				305					310					315
Thr	Glu	Phe	Ala	Ser	Gly	Gln	Trp	Val	Gly	Val	Glu	Leu	Asp	Glu
				320					325					330
Pro	Glu	Gly	Lys	Asn	Asp	Gly	Ser	Val	Gly	Gly	Val	Arg	Tyr	Phe
				335					340					345
Ile	Cys	Pro	Pro	Lys	Gln	Gly	Leu	Phe	Ala	Ser	Val	Ser	Lys	Ile
				350					355					360
Ser	Lys	Ala	Val	Asp	Ala	Pro	Pro	Ser	Ser	Val	Thr	Ser	Thr	Pro
				365					370					375
Arg	Thr	Pro	Arg	Met	Asp	Phe	Ser	Arg	Val	Thr	Gly	Lys	Gly	Arg
				380					385					390
Arg	Glu	His	Lys	Gly	Lys	Lys	Lys	Thr	Pro	Ser	Ser	Pro	Ser	Leu
				395					400					405
Gly	Ser	Leu	Gln	Gln	Arg	Asp	Gly	Ala	Lys	Ala	Glu	Val	Gly	Asp
				410					415					420
Gln	Val	Leu	Val	Ala	Gly	Gln	Lys	Gln	Gly	Ile	Val	Arg	Phe	Tyr
				425					430					435
Gly	Lys	Thr	Asp	Phe	Ala	Pro	Gly	Tyr	Trp	Tyr	Gly	Ile	Glu	Leu
				440					445					450
Asp	Gln	Pro	Thr	Gly	Lys	His	Asp	Gly	Ser	Val	Phe	Gly	Val	Arg
				455					460					465
Tyr	Phe	Thr	Cys	Pro	Pro	Arg	His	Gly	Val	Phe	Ala	Pro	Ala	Ser
				470					475					480
Arg	Ile	Gln	Arg	Ile	Gly	Gly	Ser	Thr	Asp	Ser	Pro	Gly	Asp	Ser
				485					490					495
Val	Gly	Ala	Lys	Lys	Val	His	Gln	Val	Thr	Met	Thr	Gln	Pro	Lys
				500					505					510
Arg	Thr	Phe	Thr	Thr	Val	Arg	Thr	Pro	Lys	Asp	Ile	Ala	Ser	Glu
				515					520					525
Asn	Ser	Ile	Ser	Arg	Leu	Leu	Phe	Cys	Cys	Trp	Phe	Pro	Trp	Met
				530					535					540
Leu	Arg	Ala	Glu	Met	Gln	Ser								
				545										

&lt;210&gt; 14

&lt;211&gt; 464

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1309364CD1

&lt;400&gt; 14

Met	Glu	Thr	Leu	Ser	Phe	Pro	Arg	Tyr	Asn	Val	Ala	Glu	Ile	Val
1				5					10					15
Ile	His	Ile	Arg	Asn	Lys	Ile	Leu	Thr	Gly	Ala	Asp	Gly	Lys	Asn
				20					25					30
Leu	Thr	Lys	Asn	Asp	Leu	Tyr	Pro	Asn	Pro	Lys	Pro	Glu	Val	Leu
				35					40					45

His Met Ile Tyr	Met Arg Ala Leu Gln	Ile Val Tyr Gly	Ile Arg
	50	55	60
Leu Glu His Phe	Tyr Met Met Pro Val	Asn Ser Glu Val	Met Tyr
	65	70	75
Pro His Leu Met	Glu Gly Phe Leu Pro	Phe Ser Asn Leu	Val Thr
	80	85	90
His Leu Asp Ser	Phe Leu Pro Ile Cys	Arg Val Asn Asp	Phe Glu
	95	100	105
Thr Ala Asp Ile	Leu Cys Pro Lys Ala	Lys Arg Thr Ser	Arg Phe
	110	115	120
Leu Ser Gly Ile	Ile Asn Phe Ile His	Phe Arg Glu Ala	Cys Arg
	125	130	135
Glu Thr Tyr Met	Glu Phe Leu Trp Gln	Tyr Lys Ser Ser	Ala Asp
	140	145	150
Lys Met Gln Gln	Leu Asn Ala Ala His	Gln Glu Ala Leu	Met Lys
	155	160	165
Leu Glu Arg Leu	Asp Ser Val Pro Val	Glu Glu Gln Glu	Glu Phe
	170	175	180
Lys Gln Leu Ser	Asp Gly Ile Gln Glu	Leu Gln Gln Ser	Leu Asn
	185	190	195
Gln Asp Phe His	Gln Lys Thr Ile Val	Leu Gln Glu Gly	Asn Pro
	200	205	210
Gln Lys Lys Ser	Asn Ile Ser Glu Lys	Thr Lys Arg Leu	Asn Glu
	215	220	225
Leu Lys Leu Leu	Val Val Ser Leu Lys	Glu Ile Gln Glu	Ser Leu
	230	235	240
Lys Thr Lys Ile	Val Asp Ser Pro Glu	Lys Leu Lys Asn	Tyr Lys
	245	250	255
Glu Lys Met Lys	Asp Thr Val Gln Lys	Leu Lys Asn Ala	Arg Gln
	260	265	270
Glu Val Val Glu	Lys Tyr Glu Ile Tyr	Gly Asp Ser Val	Asp Cys
	275	280	285
Leu Pro Ser Cys	Gln Leu Glu Val Gln	Leu Tyr Gln Lys	Lys Ile
	290	295	300
Gln Asp Leu Ser	Asp Asn Arg Glu Lys	Leu Ala Ser Ile	Leu Lys
	305	310	315
Glu Ser Leu Asn	Leu Glu Asp Gln Ile	Glu Ser Asp Glu	Ser Glu
	320	325	330
Leu Lys Lys Leu	Lys Thr Glu Glu Asn	Ser Phe Lys Arg	Leu Met
	335	340	345
Ile Val Lys Lys	Glu Lys Leu Ala Thr	Ala Gln Phe Lys	Ile Asn
	350	355	360
Lys Lys His Glu	Asp Val Lys Gln Tyr	Lys Arg Thr Val	Ile Glu
	365	370	375
Asp Cys Asn Lys	Val Gln Glu Lys Arg	Gly Ala Val Tyr	Glu Arg
	380	385	390
Val Thr Thr Ile	Asn Gln Glu Ile Gln	Lys Ile Lys Leu	Gly Ile
	395	400	405
Gln Gln Leu Lys	Asp Ala Ala Glu Arg	Glu Lys Leu Lys	Ser Gln
	410	415	420
Glu Ile Phe Leu	Asn Leu Lys Thr Ala	Leu Glu Lys Tyr	His Asp
	425	430	435
Gly Ile Glu Lys	Ala Ala Glu Asp Ser	Tyr Ala Lys Ile	Asp Glu
	440	445	450
Lys Thr Ala Glu	Leu Lys Arg Lys Met	Phe Lys Met Ser	Thr
	455	460	

&lt;210&gt; 15

&lt;211&gt; 569

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1315267CD1

&lt;400&gt; 15

Met	Glu	Ser	Ile	Lys	His	Lys	Val	Ser	Glu	Pro	Ser	Arg	Ser	Ser
1				5					10					15
Ser	Leu	Ser	Leu	Ser	Lys	Met	Asp	Phe	Asp	Asp	Glu	Arg	Thr	Trp
				20					25					30
Thr	Asp	Leu	Glu	Glu	Asn	Leu	Cys	Asn	His	Asp	Val	Val	Leu	Gly
				35					40					45
Asn	Glu	Ser	Thr	Tyr	Gly	Thr	Pro	Gln	Thr	Cys	Tyr	Pro	Asn	Asn
				50					55					60
Glu	Ile	Gly	Ile	Leu	Asp	Lys	Thr	Ile	Lys	Arg	Lys	Ile	Ala	Pro
				65					70					75
Val	Lys	Arg	Gly	Glu	Asp	Leu	Ser	Lys	Ser	Arg	Arg	Ser	Arg	Ser
				80					85					90
Pro	Pro	Thr	Ser	Glu	Leu	Met	Met	Lys	Phe	Phe	Pro	Ser	Leu	Lys
				95					100					105
Pro	Lys	Pro	Lys	Ser	Asp	Ser	His	Leu	Gly	Asn	Glu	Leu	Lys	Leu
				110					115					120
Asn	Ile	Ser	Gln	Asp	Gln	Pro	Pro	Gly	Asp	Asn	Ala	Arg	Ser	Gln
				125					130					135
Val	Leu	Arg	Glu	Lys	Ile	Ile	Glu	Leu	Glu	Thr	Glu	Ile	Glu	Lys
				140					145					150
Phe	Lys	Ala	Glu	Asn	Ala	Ser	Leu	Ala	Lys	Leu	Arg	Ile	Glu	Arg
				155					160					165
Glu	Ser	Ala	Leu	Glu	Lys	Leu	Arg	Lys	Glu	Ile	Ala	Asp	Phe	Glu
				170					175					180
Gln	Gln	Lys	Ala	Lys	Glu	Leu	Ala	Arg	Ile	Glu	Glu	Phe	Lys	Lys
				185					190					195
Glu	Glu	Met	Arg	Lys	Leu	Gln	Lys	Glu	Arg	Lys	Val	Phe	Glu	Lys
				200					205					210
Tyr	Thr	Thr	Ala	Ala	Arg	Thr	Phe	Pro	Asp	Lys	Lys	Glu	Arg	Glu
				215					220					225
Glu	Ile	Gln	Thr	Leu	Lys	Gln	Gln	Ile	Ala	Asp	Leu	Arg	Glu	Asp
				230					235					240
Leu	Lys	Arg	Lys	Glu	Thr	Lys	Trp	Ser	Ser	Thr	His	Ser	Arg	Leu
				245					250					255
Arg	Ser	Gln	Ile	Gln	Met	Leu	Val	Arg	Glu	Asn	Thr	Asp	Leu	Arg
				260					265					270
Glu	Glu	Ile	Lys	Val	Met	Glu	Arg	Phe	Arg	Leu	Asp	Ala	Trp	Lys
				275					280					285
Arg	Ala	Glu	Ala	Ile	Glu	Ser	Ser	Leu	Glu	Val	Glu	Lys	Lys	Asp
				290					295					300
Lys	Leu	Ala	Asn	Thr	Ser	Val	Arg	Phe	Gln	Asn	Ser	Gln	Ile	Ser
				305					310					315
Ser	Gly	Thr	Gln	Val	Glu	Lys	Tyr	Lys	Lys	Asn	Tyr	Leu	Pro	Met
				320					325					330
Gln	Gly	Asn	Pro	Pro	Arg	Arg	Ser	Lys	Ser	Ala	Pro	Pro	Arg	Asp
				335					340					345
Leu	Gly	Asn	Leu	Asp	Lys	Gly	Gln	Ala	Ala	Ser	Pro	Arg	Glu	Pro
				350					355					360
Leu	Glu	Pro	Leu	Asn	Phe	Pro	Asp	Pro	Glu	Tyr	Lys	Glu	Glu	Glu
				365					370					375
Glu	Asp	Gln	Asp	Ile	Gln	Gly	Glu	Ile	Ser	His	Pro	Asp	Gly	Lys
				380					385					390
Val	Glu	Lys	Val	Tyr	Lys	Asn	Gly	Cys	Arg	Val	Ile	Leu	Phe	Pro
				395					400					405
Asn	Gly	Thr	Arg	Lys	Glu	Val	Ser	Ala	Asp	Gly	Lys	Thr	Ile	Thr
				410					415					420
Val	Thr	Phe	Phe	Asn	Gly	Asp	Val	Lys	Gln	Val	Met	Pro	Asp	Gln
				425					430					435
Arg	Val	Ile	Tyr	Tyr	Tyr	Ala	Ala	Ala	Gln	Thr	Thr	His	Thr	Thr
				440					445					450
Tyr	Pro	Glu	Gly	Leu	Glu	Val	Leu	His	Phe	Ser	Ser	Gly	Gln	Ile
				455					460					465
Glu	Lys	His	Tyr	Pro	Asp	Gly	Arg	Lys	Glu	Ile	Thr	Phe	Pro	Asp
				470					475					480
Gln	Thr	Val	Lys	Asn	Leu	Phe	Pro	Asp	Gly	Gln	Glu	Glu	Ser	Ile
				485					490					495
Phe	Pro	Asp	Gly	Thr	Ile	Val	Arg	Val	Gln	Arg	Asp	Gly	Asn	Lys

Leu	Ile	Glu	Phe	500	Asn	Asn	Gly	Gln	Arg	505	Glu	Leu	His	Thr	Ala	Gln	510
				515						520							525
Phe	Lys	Arg	Arg	530	Glu	Tyr	Pro	Asp	Gly	535	Thr	Val	Lys	Thr	Val	Tyr	540
Ala	Asn	Gly	His	545	Gln	Glu	Thr	Lys	Tyr	550	Arg	Ser	Gly	Arg	Ile	Arg	555
Val	Lys	Asp	Lys	560	Glu	Gly	Asn	Val	Leu	565	Met	Asp	Thr	Glu	Leu		

&lt;210&gt; 16

&lt;211&gt; 436

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1403289CD1

&lt;400&gt; 16

Met	Leu	Leu	Ser	Pro	Lys	Phe	Ser	Leu	Ser	Thr	Ile	His	Ile	Arg			
1				5					10					15			
Leu	Thr	Ala	Lys	Gly	Leu	Leu	Arg	Asn	Leu	Arg	Leu	Pro	Ser	Gly			
				20					25					30			
Phe	Arg	Arg	Ser	Thr	Val	Val	Phe	His	Thr	Val	Glu	Lys	Ser	Arg			
				35					40					45			
Gln	Lys	Asn	Pro	Arg	Ser	Leu	Cys	Ile	Gln	Pro	Gln	Thr	Ala	Pro			
				50					55					60			
Asp	Ala	Leu	Pro	Pro	Glu	Lys	Thr	Leu	Glu	Leu	Thr	Gln	Tyr	Lys			
				65					70					75			
Thr	Lys	Cys	Glu	Asn	Gln	Ser	Gly	Phe	Ile	Leu	Gln	Leu	Lys	Gln			
				80					85					90			
Leu	Leu	Ala	Cys	Gly	Asn	Thr	Lys	Phe	Glu	Ala	Leu	Thr	Val	Val			
				95					100					105			
Ile	Gln	His	Leu	Leu	Ser	Glu	Arg	Glu	Glu	Ala	Leu	Lys	Gln	His			
				110					115					120			
Lys	Thr	Leu	Ser	Gln	Glu	Leu	Val	Asn	Leu	Arg	Gly	Glu	Leu	Val			
				125					130					135			
Thr	Ala	Ser	Thr	Thr	Cys	Glu	Lys	Leu	Glu	Lys	Ala	Arg	Asn	Glu			
				140					145					150			
Leu	Gln	Thr	Val	Tyr	Glu	Ala	Phe	Val	Gln	Gln	His	Gln	Ala	Glu			
				155					160					165			
Lys	Thr	Glu	Arg	Glu	Asn	Arg	Leu	Lys	Glu	Phe	Tyr	Thr	Arg	Glu			
				170					175					180			
Tyr	Glu	Lys	Leu	Arg	Asp	Thr	Tyr	Ile	Glu	Glu	Ala	Glu	Lys	Tyr			
				185					190					195			
Lys	Met	Gln	Leu	Gln	Glu	Gln	Phe	Asp	Asn	Leu	Asn	Ala	Ala	His			
				200					205					210			
Glu	Thr	Ser	Lys	Leu	Glu	Ile	Glu	Ala	Ser	His	Ser	Glu	Lys	Leu			
				215					220					225			
Glu	Leu	Leu	Lys	Lys	Ala	Tyr	Glu	Ala	Ser	Leu	Ser	Glu	Ile	Lys			
				230					235					240			
Lys	Gly	His	Glu	Ile	Glu	Lys	Lys	Ser	Leu	Glu	Asp	Leu	Leu	Ser			
				245					250					255			
Glu	Lys	Gln	Glu	Ser	Leu	Glu	Lys	Gln	Ile	Asn	Asp	Leu	Lys	Ser			
				260					265					270			
Glu	Asn	Asp	Ala	Leu	Asn	Glu	Lys	Leu	Lys	Ser	Glu	Glu	Gln	Lys			
				275					280					285			
Arg	Arg	Ala	Arg	Glu	Lys	Ala	Asn	Leu	Lys	Asn	Pro	Gln	Ile	Met			
				290					295					300			
Tyr	Leu	Glu	Gln	Glu	Leu	Glu	Ser	Leu	Lys	Ala	Val	Leu	Glu	Ile			
				305					310					315			
Lys	Asn	Glu	Lys	Leu	His	Gln	Gln	Asp	Ile	Lys	Leu	Met	Lys	Met			
				320					325					330			
Glu	Lys	Leu	Val	Asp	Asn	Asn	Thr	Ala	Leu	Val	Asp	Lys	Leu	Lys			
				335					340					345			
Arg	Phe	Gln	Gln	Glu	Asn	Glu	Glu	Leu	Lys	Ala	Arg	Met	Asp	Lys			

His Met Ala Ile	350	Ser Arg Gln Leu Ser	355	Thr Glu Gln Ala Val	360
	365		370		375
Gln Glu Ser Leu	380	Glu Lys Glu Ser Lys	385	Val Asn Lys Arg Leu	390
Met Glu Asn Glu	395	Glu Leu Leu Trp Lys	400	Leu His Asn Gly Asp	405
Cys Ser Pro Lys	410	Arg Ser Pro Thr Ser	415	Ser Ala Ile Pro Leu	420
Ser Pro Arg Asn	425	Ser Gly Ser Leu His	430	Ser Pro Ser Ile Ser	435
Arg					

<210> 17  
 <211> 363  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1607607CD1

<400> 17

Met Ala Ser Ala	Glu Leu Gln Gly Lys Tyr	Gln Lys Leu Ala Gln
1	5	10
Glu Tyr Ser Lys	Leu Arg Ala Gln Asn Gln	Val Leu Lys Lys Gly
	20	25
Val Val Asp Glu	Gln Ala Asn Ser Ala Ala	Leu Lys Glu Gln Leu
	35	40
Lys Met Lys Asp	Gln Ser Leu Arg Lys Leu	Gln Gln Glu Met Asp
	50	55
Ser Leu Thr Phe	Arg Asn Leu Gln Leu Ala	Lys Arg Val Glu Leu
	65	70
Leu Gln Asp Glu	Leu Ala Leu Ser Glu Pro	Arg Gly Lys Lys Asn
	80	85
Lys Lys Ser Gly	Glu Ser Ser Ser Gln Leu	Ser Gln Glu Gln Lys
	95	100
Ser Val Phe Asp	Glu Asp Leu Gln Lys Lys	Ile Glu Glu Asn Glu
	110	115
Arg Leu His Ile	Gln Phe Phe Glu Ala Asp	Glu Gln His Lys His
	125	130
Val Glu Ala Glu	Leu Arg Ser Arg Leu Ala	Thr Leu Glu Thr Glu
	140	145
Ala Ala Gln His	Gln Ala Val Val Asp Gly	Leu Thr Arg Lys Tyr
	155	160
Met Glu Thr Ile	Glu Lys Leu Gln Asn Asp	Lys Ala Lys Leu Glu
	170	175
Val Lys Ser Gln	Thr Leu Glu Lys Glu Ala	Lys Glu Cys Arg Leu
	185	190
Arg Thr Glu Glu	Cys Gln Leu Gln Leu Lys	Thr Leu His Glu Asp
	200	205
Leu Ser Gly Arg	Leu Glu Glu Ser Leu Ser	Ile Ile Asn Glu Lys
	215	220
Val Pro Phe Asn	Asp Thr Lys Tyr Ser Gln	Tyr Asn Ala Leu Asn
	230	235
Val Pro Leu His	Asn Arg Arg His Gln Leu	Lys Met Arg Asp Ile
	245	250
Ala Gly Gln Ala	Leu Ala Phe Val Gln Asp	Leu Val Thr Ala Leu
	260	265
Leu Asn Phe His	Thr Tyr Thr Glu Gln Arg	Ile Gln Ile Phe Pro
	275	280
Val Asp Ser Ala	Ile Asp Thr Ile Ser Pro	Leu Asn Gln Lys Phe
	290	295
Ser Gln Tyr Leu	His Glu Asn Ala Ser Tyr	Val Arg Pro Leu Glu
	305	310
Glu Gly Met Leu	His Leu Phe Glu Ser Ile	Thr Glu Asp Thr Val

Thr Val Leu Glu	320	Thr Val Lys Leu	325	Thr Phe Ser Glu	330
	335		340		345
Leu Thr Ser Tyr	Ile Cys Phe Leu Arg	Lys Ile Leu Pro Tyr	Gln		
	350		355		360
Leu Lys Arg					

<210> 18  
 <211> 247  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1660025CD1

<400> 18

Met Gly Lys Arg Asp	Asn Arg Val Ala Tyr	Met Asn Pro Ile Ala
1	5	10
Met Ala Arg Ser Arg	Gly Pro Ile Gln Ser	Ser Gly Pro Thr Ile
	20	25
Gln Asp Tyr Leu Asn	Arg Pro Arg Pro Thr	Trp Glu Glu Val Lys
	35	40
Glu Gln Leu Glu Lys	Lys Lys Lys Gly Ser	Lys Ala Leu Ala Glu
	50	55
Phe Glu Glu Lys Met	Asn Glu Asn Trp Lys	Lys Glu Leu Glu Lys
	65	70
His Arg Glu Lys Leu	Leu Ser Gly Ser Glu	Ser Ser Ser Lys Lys
	80	85
Arg Gln Arg Lys Lys	Lys Glu Lys Lys Lys	Ser Gly Arg Tyr Ser
	95	100
Ser Ser Ser Ser Ser	Ser Ser Asp Ser Ser	Ser Ser Ser Ser Asp
	110	115
Ser Glu Asp Glu Asp	Lys Lys Gln Gly Lys	Arg Arg Lys Lys Lys
	125	130
Lys Asn Arg Ser His	Lys Ser Ser Glu Ser	Ser Met Ser Glu Thr
	140	145
Glu Ser Asp Ser Lys	Asp Ser Leu Lys Lys	Lys Lys Lys Ser Lys
	155	160
Asp Gly Thr Glu Lys	Glu Lys Asp Ile Lys	Gly Leu Ser Lys Lys
	170	175
Arg Lys Met Tyr Ser	Glu Asp Lys Pro Leu	Ser Ser Glu Ser Leu
	185	190
Ser Glu Ser Glu Tyr	Ile Glu Glu Val Arg	Ala Lys Lys Lys Lys
	200	205
Ser Ser Glu Glu Arg	Glu Lys Ala Thr Glu	Lys Thr Lys Lys Lys
	215	220
Lys Lys His Lys Lys	His Ser Lys Lys Lys	Lys Lys Lys Ala Ala
	230	235
Ser Ser Ser Pro Asp	Ser Pro	
	245	

<210> 19  
 <211> 441  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1796836CD1

<400> 19

Met Asp Asp Asp Asp	Phe Gly Gly Phe Glu	Ala Ala Glu Thr Phe
1	5	10
Asp Gly Gly Ser Gly	Glu Thr Gln Thr Thr	Ser Pro Ala Ile Pro
	20	25

Trp	Ala	Ala	Phe	Pro	Ala	Val	Ser	Gly	Val	His	Leu	Ser	Pro	Ser
				35					40					45
Ser	Pro	Glu	Ile	Val	Leu	Asp	Arg	Asp	His	Ser	Ser	Ser	Ile	Gly
				50					55					60
Cys	Leu	Ser	Ser	Asp	Ala	Ile	Ile	Ser	Ser	Pro	Glu	Asn	Thr	His
				65					70					75
Ala	Ala	Asn	Ser	Ile	Val	Ser	Gln	Thr	Ile	Pro	Lys	Ala	Gln	Ile
				80					85					90
Gln	Gln	Ser	Thr	His	Thr	His	Leu	Asp	Ile	Ser	Leu	Phe	Pro	Leu
				95					100					105
Gly	Leu	Thr	Asp	Glu	Lys	Ser	Asn	Gly	Thr	Ile	Ala	Leu	Val	Asp
				110					115					120
Asp	Ser	Glu	Asp	Pro	Gly	Ala	Asn	Val	Ser	Asn	Ile	Gln	Leu	Gln
				125					130					135
Gln	Lys	Ile	Ser	Ser	Leu	Glu	Ile	Lys	Leu	Lys	Val	Ser	Glu	Glu
				140					145					150
Glu	Lys	Gln	Arg	Ile	Lys	Gln	Asp	Val	Glu	Ser	Leu	Met	Glu	Lys
				155					160					165
His	Asn	Val	Leu	Glu	Lys	Gly	Phe	Leu	Lys	Glu	Lys	Glu	Gln	Glu
				170					175					180
Ala	Ile	Ser	Phe	Gln	Asp	Arg	Tyr	Lys	Glu	Leu	Gln	Glu	Lys	His
				185					190					195
Lys	Gln	Glu	Leu	Glu	Asp	Met	Arg	Lys	Ala	Gly	His	Glu	Ala	Leu
				200					205					210
Ser	Ile	Ile	Val	Asp	Glu	Tyr	Lys	Ala	Leu	Leu	Gln	Ser	Ser	Val
				215					220					225
Lys	Gln	Gln	Val	Glu	Ala	Ile	Glu	Lys	Gln	Tyr	Ile	Ser	Ala	Ile
				230					235					240
Glu	Lys	Gln	Ala	His	Lys	Cys	Glu	Glu	Leu	Leu	Asn	Ala	Gln	His
				245					250					255
Gln	Arg	Leu	Leu	Glu	Met	Leu	Asp	Thr	Glu	Lys	Glu	Leu	Leu	Lys
				260					265					270
Glu	Lys	Ile	Lys	Glu	Ala	Leu	Ile	Gln	Gln	Ser	Gln	Glu	Gln	Lys
				275					280					285
Glu	Ile	Leu	Glu	Lys	Cys	Leu	Glu	Glu	Glu	Arg	Gln	Arg	Asn	Lys
				290					295					300
Glu	Ala	Leu	Val	Ser	Ala	Ala	Lys	Leu	Glu	Lys	Glu	Ala	Met	Lys
				305					310					315
Asp	Ala	Val	Leu	Lys	Val	Val	Glu	Glu	Glu	Arg	Lys	Asn	Leu	Glu
				320					325					330
Lys	Ala	His	Ala	Glu	Glu	Arg	Glu	Leu	Trp	Lys	Thr	Glu	His	Ala
				335					340					345
Lys	Asp	Gln	Glu	Lys	Val	Ser	Gln	Glu	Ile	Gln	Lys	Ala	Ile	Gln
				350					355					360
Glu	Gln	Arg	Lys	Ile	Ser	Gln	Glu	Thr	Val	Lys	Ala	Ala	Ile	Ile
				365					370					375
Glu	Glu	Gln	Lys	Arg	Ser	Glu	Lys	Ala	Val	Glu	Glu	Ala	Val	Lys
				380					385					390
Arg	Thr	Arg	Asp	Glu	Leu	Ile	Glu	Tyr	Ile	Lys	Glu	Gln	Lys	Arg
				395					400					405
Leu	Asp	Gln	Val	Ile	Arg	Gln	Arg	Ser	Leu	Ser	Ser	Leu	Glu	Leu
				410					415					420
Phe	Leu	Ser	Cys	Ala	Gln	Lys	Gln	Leu	Ser	Ala	Leu	Ile	Ala	Thr
				425					430					435
Glu	Pro	Val	Asp	Ile	Glu									
				440										

&lt;210&gt; 20

&lt;211&gt; 183

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2880670CD1

&lt;400&gt; 20

Met	Ala	Ala	Gln	Arg	Gly	Met	Pro	Ser	Ser	Ala	Val	Arg	Val	Leu
1				5					10					15
Glu	Glu	Ala	Leu	Gly	Met	Gly	Leu	Thr	Ala	Ala	Gly	Asp	Ala	Arg
				20					25					30
Asp	Thr	Ala	Asp	Ala	Val	Ala	Ala	Glu	Gly	Ala	Tyr	Tyr	Leu	Glu
				35					40					45
Gln	Val	Thr	Ile	Thr	Glu	Ala	Ser	Glu	Asp	Asp	Tyr	Glu	Tyr	Glu
				50					55					60
Glu	Ile	Pro	Asp	Asp	Asn	Phe	Ser	Ile	Pro	Glu	Gly	Glu	Glu	Asp
				65					70					75
Leu	Ala	Lys	Ala	Ile	Gln	Met	Ala	Gln	Glu	Gln	Ala	Thr	Asp	Thr
				80					85					90
Glu	Ile	Leu	Glu	Arg	Lys	Thr	Val	Leu	Pro	Ser	Lys	His	Ala	Val
				95					100					105
Pro	Glu	Val	Ile	Glu	Asp	Phe	Leu	Cys	Asn	Phe	Leu	Ile	Lys	Met
				110					115					120
Gly	Met	Thr	Arg	Thr	Leu	Asp	Cys	Phe	Gln	Ser	Glu	Trp	Tyr	Glu
				125					130					135
Leu	Ile	Gln	Lys	Gly	Val	Thr	Glu	Leu	Arg	Thr	Val	Gly	Asn	Val
				140					145					150
Pro	Asp	Val	Tyr	Thr	Gln	Ile	Met	Leu	Leu	Glu	Asn	Glu	Asn	Lys
				155					160					165
Asn	Leu	Lys	Lys	Asp	Leu	Lys	His	Tyr	Lys	Gln	Ala	Ala	Glu	Tyr
				170					175					180
Val	Ile	Phe												

<210> 21  
 <211> 212  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2913976CD1

Met	Glu	Glu	Glu	Leu	Pro	Pro	Pro	Pro	Ala	Glu	Pro	Val	Glu	Lys
1				5					10					15
Gly	Ala	Ser	Thr	Asp	Ile	Cys	Ala	Phe	Cys	His	Lys	Thr	Val	Phe
				20					25					30
Pro	Arg	Glu	Leu	Ala	Val	Glu	Ala	Met	Lys	Arg	Gln	Tyr	His	Ala
				35					40					45
Gln	Cys	Phe	Thr	Cys	Arg	Thr	Cys	Arg	Arg	Gln	Leu	Ala	Gly	Gln
				50					55					60
Ser	Phe	Tyr	Gln	Lys	Asp	Gly	Arg	Pro	Leu	Cys	Glu	Pro	Cys	Tyr
				65					70					75
Gln	Asp	Thr	Leu	Glu	Arg	Cys	Gly	Lys	Cys	Gly	Glu	Val	Val	Arg
				80					85					90
Asp	His	Ile	Ile	Arg	Ala	Leu	Gly	Gln	Ala	Phe	His	Pro	Ser	Cys
				95					100					105
Phe	Thr	Cys	Val	Thr	Cys	Ala	Arg	Cys	Ile	Gly	Asp	Glu	Ser	Phe
				110					115					120
Ala	Leu	Gly	Ser	Gln	Asn	Glu	Val	Tyr	Cys	Leu	Asp	Asp	Phe	Tyr
				125					130					135
Arg	Lys	Phe	Ala	Pro	Val	Cys	Ser	Ile	Cys	Glu	Asn	Pro	Ile	Ile
				140					145					150
Pro	Arg	Asp	Gly	Lys	Asp	Ala	Phe	Lys	Ile	Glu	Cys	Met	Gly	Arg
				155					160					165
Asn	Phe	His	Glu	Asn	Cys	Tyr	Arg	Cys	Glu	Asp	Cys	Arg	Ile	Leu
				170					175					180
Leu	Ser	Val	Glu	Pro	Thr	Asp	Gln	Gly	Cys	Tyr	Pro	Leu	Asn	Asn
				185					190					195
His	Leu	Phe	Cys	Lys	Pro	Cys	His	Val	Lys	Arg	Ser	Ala	Ala	Gly
				200					205					210
Cys	Cys													



<210> 22  
 <211> 227  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3092084CD1

<400> 22  
 Met Gly Gly Thr Thr Ser Thr Arg Arg Val Thr Phe Glu Ala Asp  
 1 5 10 15  
 Glu Asn Glu Asn Ile Thr Val Val Lys Gly Ile Arg Leu Ser Glu  
 20 25 30  
 Asn Val Ile Asp Arg Met Lys Glu Ser Ser Pro Ser Gly Ser Lys  
 35 40 45  
 Ser Gln Arg Tyr Ser Gly Ala Tyr Gly Ala Ser Val Ser Asp Glu  
 50 55 60  
 Glu Leu Lys Arg Arg Val Ala Glu Glu Leu Ala Leu Glu Gln Ala  
 65 70 75  
 Lys Lys Glu Ser Glu Asp Gln Lys Arg Leu Lys Gln Ala Lys Glu  
 80 85 90  
 Leu Asp Arg Glu Arg Ala Ala Ala Asn Glu Gln Leu Thr Arg Ala  
 95 100 105  
 Ile Leu Arg Glu Arg Ile Cys Ser Glu Glu Glu Arg Ala Lys Ala  
 110 115 120  
 Lys His Leu Ala Arg Gln Leu Glu Glu Lys Asp Arg Val Leu Lys  
 125 130 135  
 Lys Gln Asp Ala Phe Tyr Lys Glu Gln Leu Ala Arg Leu Glu Glu  
 140 145 150  
 Arg Ser Ser Glu Phe Tyr Arg Val Thr Thr Glu Gln Tyr Gln Lys  
 155 160 165  
 Ala Ala Glu Glu Val Glu Ala Lys Phe Lys Arg Tyr Glu Ser His  
 170 175 180  
 Pro Val Cys Ala Asp Leu Gln Ala Lys Ile Leu Gln Cys Tyr Arg  
 185 190 195  
 Glu Asn Thr His Gln Thr Leu Lys Cys Ser Ala Leu Ala Thr Gln  
 200 205 210  
 Tyr Met His Cys Val Asn His Ala Lys Gln Ser Met Leu Glu Lys  
 215 220 225  
 Gly Gly

<210> 23  
 <211> 490  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3882482CD1

<400> 23  
 Met Asn Leu Ala Glu Ile Cys Asp Asn Ala Lys Lys Gly Arg Glu  
 1 5 10 15  
 Tyr Ala Leu Leu Gly Asn Tyr Asp Ser Ser Met Val Tyr Tyr Gln  
 20 25 30  
 Gly Val Met Gln Gln Ile Gln Arg His Cys Gln Ser Val Arg Asp  
 35 40 45  
 Pro Ala Ile Lys Gly Lys Trp Gln Gln Val Arg Gln Glu Leu Leu  
 50 55 60  
 Glu Glu Tyr Glu Gln Val Lys Ser Ile Val Ser Thr Leu Glu Ser  
 65 70 75  
 Phe Lys Ile Asp Lys Pro Pro Asp Phe Pro Val Ser Cys Gln Asp  
 80 85 90  
 Glu Pro Phe Arg Asp Pro Ala Val Trp Pro Pro Val Pro Ala  
 95 100 105

Glu	His	Arg	Ala	Pro	Pro	Gln	Ile	Arg	Arg	Pro	Asn	Arg	Glu	Val
				110					115					120
Arg	Pro	Leu	Arg	Lys	Glu	Met	Ala	Gly	Val	Gly	Ala	Arg	Gly	Pro
				125					130					135
Val	Gly	Arg	Ala	His	Pro	Ile	Ser	Lys	Ser	Glu	Lys	Pro	Ser	Thr
				140					145					150
Ser	Arg	Asp	Lys	Asp	Tyr	Arg	Ala	Arg	Gly	Arg	Asp	Asp	Lys	Gly
				155					160					165
Arg	Lys	Asn	Met	Gln	Asp	Gly	Ala	Ser	Asn	Gly	Glu	Met	Pro	Lys
				170					175					180
Phe	Asp	Gly	Ala	Gly	Tyr	Asp	Lys	Asp	Leu	Val	Glu	Ala	Leu	Glu
				185					190					195
Arg	Asp	Ile	Val	Ser	Arg	Asn	Pro	Ser	Ile	His	Trp	Asp	Asp	Ile
				200					205					210
Ala	Asp	Leu	Glu	Glu	Ala	Lys	Lys	Leu	Leu	Arg	Glu	Ala	Val	Val
				215					220					225
Leu	Pro	Met	Trp	Met	Pro	Asp	Phe	Phe	Lys	Gly	Ile	Arg	Arg	Pro
				230					235					240
Trp	Lys	Gly	Val	Leu	Met	Val	Gly	Pro	Pro	Gly	Thr	Gly	Lys	Thr
				245					250					255
Met	Leu	Ala	Lys	Ala	Val	Ala	Thr	Glu	Cys	Gly	Thr	Thr	Phe	Phe
				260					265					270
Asn	Val	Ser	Ser	Ser	Thr	Leu	Thr	Ser	Lys	Tyr	Arg	Gly	Glu	Ser
				275					280					285
Glu	Lys	Leu	Val	Arg	Leu	Leu	Phe	Glu	Met	Ala	Arg	Phe	Tyr	Ala
				290					295					300
Pro	Thr	Thr	Ile	Phe	Ile	Asp	Glu	Ile	Asp	Ser	Ile	Cys	Ser	Arg
				305					310					315
Arg	Gly	Thr	Ser	Asp	Glu	His	Glu	Ala	Ser	Arg	Arg	Val	Lys	Ser
				320					325					330
Glu	Leu	Leu	Ile	Gln	Met	Asp	Gly	Val	Gly	Gly	Ala	Leu	Glu	Asn
				335					340					345
Asp	Asp	Pro	Ser	Lys	Met	Val	Met	Val	Leu	Ala	Ala	Thr	Asn	Phe
				350					355					360
Pro	Trp	Asp	Ile	Asp	Glu	Ala	Leu	Arg	Arg	Arg	Leu	Glu	Lys	Arg
				365					370					375
Ile	Tyr	Ile	Pro	Leu	Pro	Thr	Ala	Lys	Gly	Arg	Ala	Glu	Leu	Leu
				380					385					390
Lys	Ile	Asn	Leu	Arg	Glu	Val	Glu	Leu	Asp	Pro	Asp	Ile	Gln	Leu
				395					400					405
Glu	Asp	Ile	Ala	Glu	Lys	Ile	Glu	Gly	Tyr	Ser	Gly	Ala	Asp	Ile
				410					415					420
Thr	Asn	Val	Cys	Arg	Asp	Ala	Ser	Leu	Met	Ala	Met	Arg	Arg	Arg
				425					430					435
Ile	Asn	Gly	Leu	Ser	Pro	Glu	Glu	Ile	Arg	Ala	Leu	Ser	Lys	Glu
				440					445					450
Glu	Leu	Gln	Met	Pro	Val	Thr	Lys	Gly	Asp	Phe	Glu	Leu	Ala	Leu
				455					460					465
Lys	Lys	Ile	Ala	Lys	Ser	Val	Ser	Ala	Ala	Asp	Leu	Glu	Lys	Tyr
				470					475					480
Glu	Lys	Trp	Met	Val	Glu	Phe	Gly	Ser	Ala					
				485					490					

&lt;210&gt; 24

&lt;211&gt; 133

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 4933451CD1

&lt;400&gt; 24

Met	Ala	Ala	Arg	Thr	Val	Ile	Ile	Asp	His	Gly	Ser	Gly	Phe	Leu
1				5					10					15
Lys	Ala	Gly	Thr	Ala	Gly	Trp	Asn	Glu	Pro	Gln	Met	Val	Phe	Pro
				20					25					30

Asn	Ile	Val	Asn	Tyr	Leu	Pro	Cys	Lys	Glu	Asn	Pro	Gly	Pro	Ser
				35					40					45
Tyr	Ala	Arg	Lys	Arg	Val	Ser	Leu	Gly	Ile	Asp	Ile	Cys	His	Pro
				50					55					60
Asp	Thr	Phe	Ser	Tyr	Pro	Ile	Glu	Arg	Gly	Arg	Ile	Leu	Asn	Trp
				65					70					75
Glu	Gly	Val	Gln	Tyr	Leu	Trp	Ser	Phe	Val	Leu	Glu	Asn	His	Arg
				80					85					90
Arg	Glu	Gln	Glu	Val	Pro	Pro	Val	Ile	Ile	Thr	Glu	Thr	Pro	Leu
				95					100					105
Arg	Glu	Pro	Ala	Asp	Arg	Lys	Lys	Met	Ser	Ser	Leu	Glu	Thr	Leu
				110					115					120
Gln	Gly	Thr	Val	Phe	Pro	Gly	Trp	Pro	Ile	Ile	Gly	Val		
				125					130					

&lt;210&gt; 25

&lt;211&gt; 912

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5043904CD1

&lt;400&gt; 25

Met	Ser	Asp	Ala	Gln	Gly	Ser	Tyr	Lys	Leu	Asp	Glu	Ala	Gln	Ala
1				5					10					15
Val	Leu	Arg	Glu	Thr	Lys	Ala	Ile	Lys	Lys	Ala	Ile	Thr	Cys	Gly
				20					25					30
Glu	Lys	Glu	Lys	Gln	Asp	Leu	Ile	Lys	Ser	Leu	Ala	Met	Leu	Lys
				35					40					45
Asp	Gly	Phe	Cys	Thr	Asp	Arg	Gly	Ser	His	Ser	Asp	Leu	Trp	Ser
				50					55					60
Ser	Ser	Ser	Ser	Leu	Glu	Ser	Ser	Ser	Phe	Pro	Leu	Pro	Lys	Gln
				65					70					75
Tyr	Leu	Asp	Val	Ser	Ser	Gln	Thr	Asp	Ile	Ser	Gly	Ser	Phe	Gly
				80					85					90
Ile	Asn	Ser	Asn	Asn	Gln	Leu	Ala	Glu	Lys	Val	Arg	Leu	Arg	Leu
				95					100					105
Arg	Tyr	Glu	Glu	Ala	Lys	Arg	Arg	Ile	Ala	Asn	Leu	Lys	Ile	Gln
				110					115					120
Leu	Ala	Lys	Leu	Asp	Ser	Glu	Ala	Trp	Pro	Gly	Val	Leu	Asp	Ser
				125					130					135
Glu	Arg	Asp	Arg	Leu	Ile	Leu	Ile	Asn	Glu	Lys	Glu	Glu	Leu	Leu
				140					145					150
Lys	Glu	Met	Arg	Phe	Ile	Ser	Pro	Arg	Lys	Trp	Thr	Gln	Gly	Glu
				155					160					165
Val	Glu	Gln	Leu	Glu	Met	Ala	Arg	Lys	Arg	Leu	Glu	Lys	Asp	Leu
				170					175					180
Gln	Ala	Ala	Arg	Asp	Thr	Gln	Ser	Lys	Ala	Leu	Thr	Glu	Arg	Leu
				185					190					195
Lys	Leu	Asn	Ser	Lys	Arg	Asn	Gln	Leu	Val	Arg	Glu	Leu	Glu	Glu
				200					205					210
Ala	Thr	Arg	Gln	Val	Ala	Thr	Leu	His	Ser	Gln	Leu	Lys	Ser	Leu
				215					220					225
Ser	Ser	Ser	Met	Gln	Ser	Leu	Ser	Ser	Gly	Ser	Ser	Pro	Gly	Ser
				230					235					240
Leu	Thr	Ser	Ser	Arg	Gly	Ser	Leu	Val	Ala	Ser	Ser	Leu	Asp	Ser
				245					250					255
Ser	Thr	Ser	Ala	Ser	Phe	Thr	Asp	Leu	Tyr	Tyr	Asp	Pro	Phe	Glu
				260					265					270
Gln	Leu	Asp	Ser	Glu	Leu	Gln	Ser	Lys	Val	Glu	Phe	Leu	Leu	Leu
				275					280					285
Glu	Gly	Ala	Thr	Gly	Phe	Arg	Pro	Ser	Gly	Cys	Ile	Thr	Thr	Ile
				290					295					300
His	Glu	Asp	Glu	Val	Ala	Lys	Thr	Gln	Lys	Ala	Glu	Gly	Gly	Gly
				305					310					315

Arg	Leu	Gln	Ala	Leu	Arg	Ser	Leu	Ser	Gly	Thr	Pro	Lys	Ser	Met
				320					325					330
Thr	Ser	Leu	Ser	Pro	Arg	Ser	Ser	Leu	Ser	Ser	Pro	Ser	Pro	Pro
				335					340					345
Cys	Ser	Pro	Leu	Met	Ala	Asp	Pro	Leu	Leu	Ala	Gly	Asp	Ala	Phe
				350					355					360
Leu	Asn	Ser	Leu	Glu	Phe	Glu	Asp	Pro	Glu	Leu	Ser	Ala	Thr	Leu
				365					370					375
Cys	Glu	Leu	Ser	Leu	Gly	Asn	Ser	Ala	Gln	Glu	Arg	Tyr	Arg	Leu
				380					385					390
Glu	Glu	Pro	Gly	Thr	Glu	Gly	Lys	Gln	Leu	Gly	Gln	Ala	Val	Asn
				395					400					405
Thr	Ala	Gln	Gly	Cys	Gly	Leu	Lys	Val	Ala	Cys	Val	Ser	Ala	Ala
				410					415					420
Val	Ser	Asp	Glu	Ser	Val	Ala	Gly	Asp	Ser	Gly	Val	Tyr	Glu	Ala
				425					430					435
Ser	Val	Gln	Arg	Leu	Gly	Ala	Ser	Glu	Ala	Ala	Ala	Phe	Asp	Ser
				440					445					450
Asp	Glu	Ser	Glu	Ala	Val	Gly	Ala	Thr	Arg	Ile	Gln	Ile	Ala	Leu
				455					460					465
Lys	Tyr	Asp	Glu	Lys	Asn	Lys	Gln	Phe	Ala	Ile	Leu	Ile	Ile	Gln
				470					475					480
Leu	Ser	Asn	Leu	Ser	Ala	Leu	Leu	Gln	Gln	Gln	Asp	Gln	Lys	Val
				485					490					495
Asn	Ile	Arg	Val	Ala	Val	Leu	Pro	Cys	Ser	Glu	Ser	Thr	Thr	Cys
				500					505					510
Leu	Phe	Arg	Thr	Arg	Pro	Leu	Asp	Ala	Ser	Asp	Thr	Leu	Val	Phe
				515					520					525
Asn	Glu	Val	Phe	Trp	Val	Ser	Met	Ser	Tyr	Pro	Ala	Leu	His	Gln
				530					535					540
Lys	Thr	Leu	Arg	Val	Asp	Val	Cys	Thr	Thr	Asp	Arg	Ser	His	Leu
				545					550					555
Glu	Glu	Cys	Leu	Gly	Gly	Ala	Gln	Ile	Ser	Leu	Ala	Glu	Val	Cys
				560					565					570
Arg	Ser	Gly	Glu	Arg	Ser	Thr	Arg	Trp	Tyr	Asn	Leu	Leu	Ser	Tyr
				575					580					585
Lys	Tyr	Leu	Lys	Lys	Gln	Ser	Arg	Glu	Leu	Lys	Pro	Val	Gly	Val
				590					595					600
Met	Ala	Pro	Ala	Ser	Gly	Pro	Ala	Ser	Thr	Asp	Ala	Val	Ser	Ala
				605					610					615
Leu	Leu	Glu	Gln	Thr	Ala	Val	Glu	Leu	Glu	Lys	Arg	Gln	Glu	Gly
				620					625					630
Arg	Ser	Ser	Thr	Gln	Thr	Leu	Glu	Asp	Ser	Trp	Arg	Tyr	Glu	Glu
				635					640					645
Thr	Ser	Glu	Asn	Glu	Ala	Val	Ala	Glu	Glu	Glu	Glu	Glu	Glu	Val
				650					655					660
Glu	Glu	Glu	Glu	Gly	Glu	Glu	Asp	Val	Phe	Thr	Glu	Lys	Ala	Ser
				665					670					675
Pro	Asp	Met	Asp	Gly	Tyr	Pro	Ala	Leu	Lys	Val	Asp	Lys	Glu	Thr
				680					685					690
Asn	Thr	Glu	Thr	Pro	Ala	Pro	Ser	Pro	Thr	Val	Val	Arg	Pro	Lys
				695					700					705
Asp	Arg	Arg	Val	Gly	Thr	Pro	Ser	Gln	Gly	Pro	Phe	Leu	Arg	Gly
				710					715					720
Ser	Thr	Ile	Ile	Arg	Ser	Lys	Thr	Phe	Ser	Pro	Gly	Pro	Gln	Ser
				725					730					735
Gln	Tyr	Val	Cys	Arg	Leu	Asn	Arg	Ser	Asp	Ser	Asp	Ser	Ser	Thr
				740					745					750
Leu	Ser	Lys	Lys	Pro	Pro	Phe	Val	Arg	Asn	Ser	Leu	Glu	Arg	Arg
				755					760					765
Ser	Val	Arg	Met	Lys	Arg	Pro	Ser	Ser	Val	Lys	Ser	Leu	Arg	Ser
				770					775					780
Glu	Arg	Leu	Ile	Arg	Thr	Ser	Leu	Asp	Leu	Glu	Leu	Asp	Leu	Gln
				785					790					795
Ala	Thr	Arg	Thr	Trp	His	Ser	Gln	Leu	Thr	Gln	Glu	Ile	Ser	Val
				800					805					810
Leu	Lys	Glu	Leu	Lys	Glu	Gln	Leu	Glu	Gln	Ala	Lys	Ser	His	Gly

				815					820					825
Glu	Lys	Glu	Leu	Pro	Gln	Trp	Leu	Arg	Glu	Asp	Glu	Arg	Phe	Arg
				830					835					840
Leu	Leu	Leu	Arg	Met	Leu	Glu	Lys	Arg	Gln	Met	Asp	Arg	Ala	Glu
				845					850					855
His	Lys	Gly	Glu	Leu	Gln	Thr	Asp	Lys	Met	Met	Arg	Ala	Ala	Ala
				860					865					870
Lys	Asp	Val	His	Arg	Leu	Arg	Gly	Gln	Ser	Cys	Lys	Glu	Pro	Pro
				875					880					885
Glu	Val	Gln	Ser	Phe	Arg	Glu	Lys	Met	Ala	Phe	Phe	Thr	Arg	Pro
				890					895					900
Arg	Met	Asn	Ile	Pro	Ala	Leu	Ser	Ala	Asp	Asp	Val			
				905					910					

&lt;210&gt; 26

&lt;211&gt; 1076

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5202390CD1

&lt;400&gt; 26

Met	Lys	Gln	Tyr	Ala	Ser	Pro	Met	Pro	Thr	Gln	Thr	Asp	Val	Lys
1				5					10					15
Leu	Lys	Phe	Lys	Pro	Leu	Ser	Lys	Lys	Val	Val	Ser	Ala	Ala	Leu
				20					25					30
Gln	Phe	Ser	Leu	Ser	Cys	Ile	Phe	Leu	Arg	Glu	Gly	Lys	Ala	Thr
				35					40					45
Asp	Glu	Asp	Met	Gln	Ser	Leu	Ala	Ser	Leu	Met	Ser	Met	Lys	Gln
				50					55					60
Ala	Asp	Ile	Gly	Asn	Leu	Asp	Asp	Phe	Glu	Glu	Asp	Asn	Glu	Asp
				65					70					75
Asp	Asp	Glu	Asn	Arg	Val	Asn	Gln	Glu	Glu	Lys	Ala	Ala	Lys	Ile
				80					85					90
Thr	Glu	Leu	Ile	Asn	Lys	Leu	Asn	Phe	Leu	Asp	Glu	Ala	Glu	Lys
				95					100					105
Asp	Leu	Ala	Thr	Val	Asn	Ser	Asn	Pro	Phe	Asp	Asp	Pro	Asp	Ala
				110					115					120
Ala	Glu	Leu	Asn	Pro	Phe	Gly	Asp	Pro	Asp	Ser	Glu	Glu	Pro	Ile
				125					130					135
Thr	Glu	Thr	Ala	Ser	Pro	Arg	Lys	Thr	Glu	Asp	Ser	Phe	Tyr	Asn
				140					145					150
Asn	Ser	Tyr	Asn	Pro	Phe	Lys	Glu	Val	Gln	Thr	Pro	Gln	Tyr	Leu
				155					160					165
Asn	Pro	Phe	Asp	Glu	Pro	Glu	Ala	Phe	Val	Thr	Ile	Lys	Asp	Ser
				170					175					180
Pro	Pro	Gln	Ser	Thr	Lys	Arg	Lys	Asn	Ile	Arg	Pro	Val	Asp	Met
				185					190					195
Ser	Lys	Tyr	Leu	Tyr	Ala	Asp	Ser	Ser	Lys	Thr	Glu	Glu	Glu	Glu
				200					205					210
Leu	Asp	Glu	Ser	Asn	Pro	Phe	Tyr	Glu	Pro	Lys	Ser	Thr	Pro	Pro
				215					220					225
Pro	Asn	Asn	Leu	Val	Asn	Pro	Val	Gln	Glu	Leu	Glu	Thr	Glu	Arg
				230					235					240
Arg	Val	Lys	Arg	Lys	Ala	Pro	Ala	Pro	Pro	Val	Leu	Ser	Pro	Lys
				245					250					255
Thr	Gly	Val	Leu	Asn	Glu	Asn	Thr	Val	Ser	Ala	Gly	Lys	Asp	Leu
				260					265					270
Ser	Thr	Ser	Pro	Lys	Pro	Ser	Pro	Ile	Pro	Ser	Pro	Val	Leu	Gly
				275					280					285
Arg	Lys	Pro	Asn	Ala	Ser	Gln	Ser	Leu	Leu	Val	Trp	Cys	Lys	Glu
				290					295					300
Val	Thr	Lys	Asn	Tyr	Arg	Gly	Val	Lys	Ile	Thr	Asn	Phe	Thr	Thr
				305					310					315
Ser	Trp	Arg	Asn	Gly	Leu	Ser	Phe	Cys	Ala	Ile	Leu	His	His	Phe

	320		325		330
Arg Pro Asp Leu	Ile Asp Tyr Lys Ser	Leu Asn Pro Gln Asp	Ile		
	335		340		345
Lys Glu Asn Asn	Lys Lys Ala Tyr Asp	Gly Phe Ala Ser Ile	Gly		
	350		355		360
Ile Ser Arg Leu	Leu Glu Pro Ser Asp	Met Val Leu Leu Ala	Ile		
	365		370		375
Pro Asp Lys Leu	Thr Val Met Thr Tyr	Leu Tyr Gln Ile Arg	Ala		
	380		385		390
His Phe Ser Gly	Gln Glu Leu Asn Val	Val Gln Ile Glu Glu	Asn		
	395		400		405
Ser Ser Lys Ser	Thr Tyr Lys Val Gly	Asn Tyr Glu Thr Asp	Thr		
	410		415		420
Asn Ser Ser Val	Asp Gln Glu Lys Phe	Tyr Ala Glu Leu Ser	Asp		
	425		430		435
Leu Lys Arg Glu	Pro Glu Leu Gln Gln	Pro Ile Ser Gly Ala	Val		
	440		445		450
Asp Phe Leu Ser	Gln Asp Asp Ser Val	Phe Val Asn Asp Ser	Gly		
	455		460		465
Val Gly Glu Ser	Glu Ser Glu His Gln	Thr Pro Asp Asp His	Leu		
	470		475		480
Ser Pro Ser Thr	Ala Ser Pro Tyr Cys	Arg Arg Thr Lys Ser	Asp		
	485		490		495
Thr Glu Pro Gln	Lys Ser Gln Gln Ser	Ser Gly Arg Thr Ser	Gly		
	500		505		510
Ser Asp Asp Pro	Gly Ile Cys Ser Asn	Thr Asp Ser Thr Gln	Ala		
	515		520		525
Gln Val Leu Leu	Gly Lys Lys Arg Leu	Leu Lys Ala Glu Thr	Leu		
	530		535		540
Glu Leu Ser Asp	Leu Tyr Val Ser Asp	Lys Lys Lys Asp Met	Ser		
	545		550		555
Pro Pro Phe Ile	Cys Glu Glu Thr Asp	Glu Gln Lys Leu Gln	Thr		
	560		565		570
Leu Asp Ile Gly	Ser Asn Leu Glu Lys	Glu Lys Leu Glu Asn	Ser		
	575		580		585
Arg Ser Leu Glu	Cys Arg Ser Asp Pro	Glu Ser Pro Ile Lys	Lys		
	590		595		600
Thr Ser Leu Ser	Pro Thr Ser Lys Leu	Gly Tyr Ser Tyr Ser	Arg		
	605		610		615
Asp Leu Asp Leu	Ala Lys Lys Lys His	Ala Ser Leu Arg Gln	Thr		
	620		625		630
Glu Ser Asp Pro	Asp Ala Asp Arg Thr	Thr Leu Asn His Ala	Asp		
	635		640		645
His Ser Ser Lys	Ile Val Gln His Arg	Leu Leu Ser Arg Gln	Glu		
	650		655		660
Glu Leu Lys Glu	Arg Ala Arg Val Leu	Leu Glu Gln Ala Arg	Arg		
	665		670		675
Asp Ala Ala Leu	Lys Ala Gly Asn Lys	His Asn Thr Asn Thr	Ala		
	680		685		690
Thr Pro Phe Cys	Asn Arg Gln Leu Ser	Asp Gln Gln Asp Glu	Glu		
	695		700		705
Arg Arg Arg Gln	Leu Arg Glu Arg Ala	Arg Gln Leu Ile Ala	Glu		
	710		715		720
Ala Arg Ser Gly	Val Lys Met Ser Glu	Leu Pro Ser Tyr Gly	Glu		
	725		730		735
Met Ala Ala Glu	Lys Leu Lys Glu Arg	Ser Lys Ala Ser Gly	Asp		
	740		745		750
Glu Asn Asp Asn	Ile Glu Ile Asp Thr	Asn Glu Glu Ile Pro	Glu		
	755		760		765
Gly Phe Val Val	Gly Gly Gly Asp Glu	Leu Thr Asn Leu Glu	Asn		
	770		775		780
Asp Leu Asp Thr	Pro Glu Gln Asn Ser	Lys Leu Val Asp Leu	Lys		
	785		790		795
Leu Lys Lys Leu	Leu Glu Val Gln Pro	Gln Val Ala Asn Ser	Pro		
	800		805		810
Ser Ser Ala Ala	Gln Lys Ala Val Thr	Glu Ser Ser Glu Gln	Asp		
	815		820		825

Met	Lys	Ser	Gly	Thr	Glu	Asp	Leu	Arg	Thr	Glu	Arg	Leu	Gln	Lys
				830						835				840
Thr	Thr	Glu	Arg	Phe	Arg	Asn	Pro	Val	Val	Phe	Ser	Lys	Asp	Ser
				845						850				855
Thr	Val	Arg	Lys	Thr	Gln	Leu	Gln	Ser	Phe	Ser	Gln	Tyr	Ile	Glu
				860						865				870
Asn	Arg	Pro	Glu	Met	Lys	Arg	Gln	Arg	Ser	Ile	Gln	Glu	Asp	Thr
				875						880				885
Lys	Lys	Gly	Asn	Glu	Glu	Lys	Ala	Ala	Ile	Thr	Glu	Thr	Gln	Arg
				890						895				900
Lys	Pro	Ser	Glu	Asp	Glu	Val	Leu	Asn	Lys	Gly	Phe	Lys	Asp	Thr
				905						910				915
Ser	Gln	Tyr	Val	Val	Gly	Glu	Leu	Ala	Ala	Leu	Glu	Asn	Glu	Gln
				920						925				930
Lys	Gln	Ile	Asp	Thr	Arg	Ala	Ala	Leu	Val	Glu	Lys	Arg	Leu	Arg
				935						940				945
Tyr	Leu	Met	Asp	Thr	Gly	Arg	Asn	Thr	Glu	Glu	Glu	Glu	Ala	Met
				950						955				960
Met	Gln	Glu	Trp	Phe	Met	Leu	Val	Asn	Lys	Lys	Asn	Ala	Leu	Ile
				965						970				975
Arg	Arg	Met	Asn	Gln	Leu	Ser	Leu	Leu	Glu	Lys	Glu	His	Asp	Leu
				980						985				990
Glu	Arg	Arg	Tyr	Glu	Leu	Leu	Asn	Arg	Glu	Leu	Arg	Ala	Met	Leu
				995					1000					1005
Ala	Ile	Glu	Asp	Trp	Gln	Lys	Thr	Glu	Ala	Gln	Lys	Arg	Arg	Glu
				1010					1015					1020
Gln	Leu	Leu	Leu	Asp	Glu	Leu	Val	Ala	Leu	Val	Asn	Lys	Arg	Asp
				1025					1030					1035
Ala	Leu	Val	Arg	Asp	Leu	Asp	Ala	Gln	Glu	Lys	Gln	Ala	Glu	Glu
				1040					1045					1050
Glu	Asp	Glu	His	Leu	Glu	Arg	Thr	Leu	Glu	Gln	Asn	Lys	Gly	Lys
				1055					1060					1065
Met	Ala	Lys	Lys	Glu	Glu	Lys	Cys	Val	Leu	Gln				
				1070					1075					

&lt;210&gt; 27

&lt;211&gt; 542

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5526375CD1

&lt;400&gt; 27

Met	Glu	Glu	Arg	Gly	Ser	Pro	Asp	Gly	Asp	Leu	Ala	Arg	Ser	Leu
1				5					10					15
Glu	Gln	Gly	Pro	Glu	Gly	Pro	Glu	Thr	Pro	Ile	Gln	Val	Val	Leu
				20					25					30
Arg	Val	Arg	Pro	Met	Ser	Ala	Ala	Glu	Leu	Arg	Arg	Gly	Gln	Gln
				35					40					45
Ser	Val	Leu	His	Cys	Ser	Gly	Thr	Arg	Thr	Leu	Gln	Val	Ser	Pro
				50					55					60
Pro	Gly	Gly	Gly	Pro	Glu	Val	Ala	Phe	Arg	Phe	Gly	Ala	Val	Leu
				65					70					75
Asp	Ala	Ala	Arg	Thr	Gln	Glu	Asp	Val	Phe	Arg	Ala	Cys	Gly	Val
				80					85					90
Arg	Arg	Leu	Gly	Glu	Leu	Ala	Leu	Arg	Gly	Phe	Ser	Cys	Thr	Val
				95					100					105
Phe	Thr	Phe	Gly	Gln	Thr	Gly	Ser	Gly	Lys	Thr	Tyr	Thr	Leu	Thr
				110					115					120
Gly	Pro	Pro	Pro	Gln	Gly	Glu	Gly	Val	Pro	Val	Pro	Pro	Ser	Leu
				125					130					135
Ala	Gly	Ile	Met	Gln	Arg	Thr	Phe	Ala	Trp	Leu	Leu	Asp	Arg	Val
				140					145					150
Gln	His	Leu	Gly	Ala	Pro	Val	Thr	Leu	Arg	Ala	Ser	Tyr	Leu	Glu
				155					160					165

Ile	Tyr	Asn	Gly	Gln	Val	Arg	Asp	Leu	Leu	Ser	Leu	Gly	Ser	Pro
				170					175					180
Arg	Pro	Leu	Pro	Val	Arg	Trp	Asn	Lys	Thr	Arg	Gly	Phe	Tyr	Val
				185					190					195
Glu	Gln	Leu	Arg	Val	Val	Glu	Phe	Gly	Ser	Leu	Glu	Ala	Leu	Met
				200					205					210
Glu	Leu	Leu	Gln	Thr	Gly	Leu	Ser	Arg	Arg	Arg	Asn	Ser	Ala	His
				215					220					225
Thr	Leu	Asn	Gln	Ala	Ser	Ser	Arg	Ser	His	Ala	Leu	Leu	Thr	Leu
				230					235					240
Tyr	Ile	Ser	Arg	Gln	Thr	Ala	Gln	Gln	Met	Pro	Ser	Val	Asp	Pro
				245					250					255
Gly	Glu	Pro	Pro	Val	Gly	Gly	Lys	Leu	Cys	Phe	Val	Asp	Leu	Ala
				260					265					270
Gly	Ser	Glu	Lys	Val	Ala	Ala	Thr	Gly	Ser	Arg	Gly	Glu	Leu	Met
				275					280					285
Leu	Glu	Ala	Asn	Ser	Ile	Asn	Arg	Ser	Leu	Leu	Ala	Leu	Gly	His
				290					295					300
Cys	Ile	Ser	Leu	Leu	Leu	Asp	Pro	Gln	Arg	Lys	Gln	Ser	His	Ile
				305					310					315
Pro	Phe	Arg	Asp	Ser	Lys	Leu	Thr	Lys	Leu	Leu	Ala	Asp	Ser	Leu
				320					325					330
Gly	Gly	Arg	Gly	Val	Thr	Leu	Met	Val	Ala	Cys	Val	Ser	Pro	Ser
				335					340					345
Ala	Gln	Cys	Leu	Pro	Glu	Thr	Leu	Ser	Thr	Leu	Arg	Tyr	Ala	Ser
				350					355					360
Arg	Ala	Gln	Arg	Val	Thr	Thr	Arg	Pro	Gln	Ala	Pro	Lys	Ser	Pro
				365					370					375
Val	Ala	Lys	Gln	Pro	Gln	Arg	Leu	Glu	Thr	Glu	Met	Leu	Gln	Leu
				380					385					390
Gln	Glu	Glu	Asn	Arg	Arg	Leu	Gln	Phe	Gln	Leu	Asp	Gln	Met	Asp
				395					400					405
Cys	Lys	Ala	Ser	Gly	Leu	Ser	Gly	Ala	Arg	Val	Ala	Trp	Ala	Gln
				410					415					420
Arg	Asn	Leu	Tyr	Gly	Met	Leu	Gln	Glu	Phe	Met	Leu	Glu	Asn	Glu
				425					430					435
Arg	Leu	Arg	Lys	Glu	Lys	Ser	Gln	Leu	Gln	Asn	Ser	Arg	Glu	Leu
				440					445					450
Ala	Gln	Asn	Glu	Gln	Arg	Ile	Leu	Ala	Gln	Gln	Val	His	Ala	Leu
				455					460					465
Glu	Arg	Arg	Leu	Leu	Ser	Ala	Cys	Tyr	His	His	Gln	Gln	Gly	Pro
				470					475					480
Gly	Leu	Thr	Pro	Pro	Cys	Pro	Cys	Leu	Met	Ala	Pro	Ala	Pro	Pro
				485					490					495
Cys	His	Ala	Leu	Pro	Pro	Leu	Tyr	Ser	Cys	Pro	Cys	Cys	His	Ile
				500					505					510
Cys	Pro	Leu	Cys	Arg	Val	Pro	Leu	Ala	His	Trp	Gly	Cys	Leu	Pro
				515					520					525
Gly	Glu	His	His	Leu	Pro	Gln	Pro	Leu	Phe	Trp	Ala	Leu	Arg	Ser
				530					535					540
Gln	Lys													

&lt;210&gt; 28

&lt;211&gt; 351

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5677408CD1

&lt;400&gt; 28

Met	Pro	Ser	Glu	Thr	Leu	Trp	Glu	Ile	Ala	Lys	Ala	Glu	Val	Glu
1				5					10					15
Lys	Arg	Gly	Ile	Asn	Gly	Ser	Glu	Gly	Asp	Gly	Ala	Glu	Ile	Ala
				20					25					30



Glu	Lys	Phe	Val	Phe	Phe	Ile	Gly	Ser	Lys	Asn	Gly	Gly	Lys	Thr
				35					40					45
Thr	Ile	Ile	Leu	Arg	Cys	Leu	Asp	Arg	Asp	Glu	Pro	Pro	Lys	Pro
				50					55					60
Thr	Leu	Ala	Leu	Glu	Tyr	Thr	Tyr	Gly	Arg	Arg	Ala	Lys	Gly	His
				65					70					75
Asn	Thr	Pro	Lys	Asp	Ile	Ala	His	Phe	Trp	Glu	Leu	Gly	Gly	Gly
				80					85					90
Thr	Ser	Leu	Leu	Asp	Leu	Ile	Ser	Ile	Pro	Ile	Thr	Gly	Asp	Thr
				95					100					105
Leu	Arg	Thr	Phe	Ser	Leu	Val	Leu	Val	Leu	Asp	Leu	Ser	Lys	Pro
				110					115					120
Asn	Asp	Leu	Trp	Pro	Thr	Met	Glu	Asn	Leu	Leu	Gln	Ala	Thr	Lys
				125					130					135
Ser	His	Val	Asp	Lys	Val	Ile	Met	Lys	Leu	Gly	Lys	Thr	Asn	Ala
				140					145					150
Lys	Ala	Val	Ser	Glu	Met	Arg	Gln	Lys	Ile	Trp	Asn	Asn	Met	Pro
				155					160					165
Lys	Asp	His	Pro	Asp	His	Glu	Leu	Ile	Asp	Pro	Phe	Pro	Val	Pro
				170					175					180
Leu	Val	Ile	Ile	Gly	Ser	Lys	Tyr	Asp	Val	Phe	Gln	Asp	Phe	Glu
				185					190					195
Ser	Glu	Lys	Arg	Lys	Val	Ile	Cys	Lys	Thr	Leu	Arg	Phe	Val	Ala
				200					205					210
His	Tyr	Tyr	Gly	Ala	Ser	Leu	Met	Phe	Thr	Ser	Lys	Ser	Glu	Ala
				215					220					225
Leu	Leu	Leu	Lys	Ile	Arg	Gly	Val	Ile	Asn	Gln	Leu	Ala	Phe	Gly
				230					235					240
Ile	Asp	Lys	Ser	Lys	Ser	Ile	Cys	Val	Asp	Gln	Asn	Lys	Pro	Leu
				245					250					255
Phe	Ile	Thr	Ala	Gly	Leu	Asp	Ser	Phe	Gly	Gln	Ile	Gly	Ser	Pro
				260					265					270
Pro	Val	Pro	Glu	Asn	Asp	Ile	Gly	Lys	Leu	His	Ala	His	Ser	Pro
				275					280					285
Met	Glu	Leu	Trp	Lys	Lys	Val	Tyr	Glu	Lys	Leu	Phe	Pro	Pro	Lys
				290					295					300
Ser	Ile	Asn	Thr	Leu	Lys	Asp	Ile	Lys	Asp	Pro	Ala	Arg	Asp	Pro
				305					310					315
Gln	Tyr	Ala	Glu	Asn	Glu	Val	Asp	Glu	Met	Arg	Ile	Gln	Lys	Asp
				320					325					330
Leu	Glu	Leu	Glu	Gln	Tyr	Lys	Arg	Ser	Ser	Ser	Lys	Ser	Trp	Lys
				335					340					345
Gln	Ile	Glu	Leu	Asp	Ser									
				350										

&lt;210&gt; 29

&lt;211&gt; 856

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5982278CD1

&lt;400&gt; 29

Met	Lys	Ser	Ala	Arg	Ala	Lys	Thr	Pro	Arg	Lys	Pro	Thr	Val	Lys
1				5					10					15
Lys	Gly	Ser	Gln	Thr	Asn	Leu	Lys	Asp	Pro	Val	Gly	Val	Tyr	Cys
				20					25					30
Arg	Val	Arg	Pro	Leu	Gly	Phe	Pro	Asp	Gln	Glu	Cys	Cys	Ile	Glu
				35					40					45
Val	Ile	Asn	Asn	Thr	Thr	Val	Gln	Leu	His	Thr	Pro	Glu	Gly	Tyr
				50					55					60
Arg	Leu	Asn	Arg	Asn	Gly	Asp	Tyr	Lys	Glu	Thr	Gln	Tyr	Ser	Phe
				65					70					75
Lys	Gln	Val	Phe	Gly	Thr	His	Thr	Thr	Gln	Lys	Glu	Leu	Phe	Asp
				80					85					90

Val	Val	Ala	Asn	Pro	Leu	Val	Asn	Asp	Leu	Ile	His	Gly	Lys	Asn
				95					100					105
Gly	Leu	Leu	Phe	Thr	Tyr	Gly	Val	Thr	Gly	Ser	Gly	Lys	Thr	His
				110					115					120
Thr	Met	Thr	Gly	Ser	Pro	Gly	Glu	Gly	Gly	Leu	Leu	Pro	Arg	Cys
				125					130					135
Leu	Asp	Met	Ile	Phe	Asn	Ser	Ile	Gly	Ser	Phe	Gln	Ala	Lys	Arg
				140					145					150
Tyr	Val	Phe	Lys	Ser	Asn	Asp	Arg	Asn	Ser	Met	Asp	Ile	Gln	Cys
				155					160					165
Glu	Val	Asp	Ala	Leu	Leu	Glu	Arg	Gln	Lys	Arg	Glu	Ala	Met	Pro
				170					175					180
Asn	Pro	Lys	Thr	Ser	Ser	Ser	Lys	Arg	Gln	Val	Asp	Pro	Glu	Phe
				185					190					195
Ala	Asp	Met	Ile	Thr	Val	Gln	Glu	Phe	Cys	Lys	Ala	Glu	Glu	Val
				200					205					210
Asp	Glu	Asp	Ser	Val	Tyr	Gly	Val	Phe	Val	Ser	Tyr	Ile	Glu	Ile
				215					220					225
Tyr	Asn	Asn	Tyr	Ile	Tyr	Asp	Leu	Leu	Glu	Glu	Val	Pro	Phe	Asp
				230					235					240
Pro	Ile	Asn	Pro	Asn	Leu	His	Asn	Leu	Asn	Cys	Phe	Val	Lys	Ile
				245					250					255
Lys	Asn	His	Asn	Met	Tyr	Val	Ala	Gly	Cys	Thr	Glu	Val	Glu	Val
				260					265					270
Lys	Ser	Thr	Glu	Glu	Ala	Phe	Glu	Val	Phe	Trp	Arg	Gly	Gln	Lys
				275					280					285
Lys	Arg	Arg	Ile	Ala	Asn	Thr	His	Leu	Asn	Arg	Glu	Ser	Ser	Arg
				290					295					300
Ser	His	Ser	Val	Phe	Asn	Ile	Lys	Leu	Val	Gln	Ala	Pro	Leu	Asp
				305					310					315
Ala	Asp	Gly	Asp	Asn	Val	Leu	Gln	Glu	Lys	Glu	Gln	Ile	Thr	Ile
				320					325					330
Ser	Gln	Leu	Ser	Leu	Val	Asp	Leu	Ala	Gly	Ser	Glu	Arg	Thr	Asn
				335					340					345
Arg	Thr	Arg	Ala	Glu	Gly	Asn	Arg	Leu	Arg	Glu	Ala	Gly	Asn	Ile
				350					355					360
Asn	Gln	Ser	Leu	Met	Thr	Leu	Arg	Thr	Cys	Met	Asp	Val	Leu	Arg
				365					370					375
Glu	Asn	Gln	Met	Tyr	Gly	Thr	Asn	Lys	Met	Val	Pro	Tyr	Arg	Asp
				380					385					390
Ser	Lys	Leu	Thr	His	Leu	Phe	Lys	Asn	Tyr	Phe	Asp	Gly	Glu	Gly
				395					400					405
Lys	Val	Arg	Met	Ile	Val	Cys	Val	Asn	Pro	Lys	Ala	Glu	Asp	Tyr
				410					415					420
Glu	Glu	Asn	Leu	Gln	Val	Met	Arg	Phe	Ala	Glu	Val	Thr	Gln	Glu
				425					430					435
Val	Glu	Val	Ala	Arg	Pro	Val	Asp	Lys	Ala	Ile	Cys	Gly	Leu	Thr
				440					445					450
Pro	Gly	Arg	Arg	Tyr	Arg	Asn	Gln	Pro	Arg	Gly	Pro	Val	Gly	Asn
				455					460					465
Glu	Pro	Leu	Val	Thr	Asp	Val	Val	Leu	Gln	Ser	Phe	Pro	Pro	Leu
				470					475					480
Pro	Ser	Cys	Glu	Ile	Leu	Asp	Ile	Asn	Asp	Glu	Gln	Thr	Leu	Pro
				485					490					495
Arg	Leu	Ile	Glu	Ala	Leu	Glu	Lys	Arg	His	Asn	Leu	Arg	Gln	Met
				500					505					510
Met	Ile	Asp	Glu	Phe	Asn	Lys	Gln	Ser	Asn	Ala	Phe	Lys	Ala	Leu
				515					520					525
Leu	Gln	Glu	Phe	Asp	Asn	Ala	Val	Leu	Ser	Lys	Glu	Asn	His	Met
				530					535					540
Gln	Gly	Lys	Leu	Asn	Glu	Lys	Glu	Lys	Met	Ile	Ser	Gly	Gln	Lys
				545					550					555
Leu	Glu	Ile	Glu	Arg	Leu	Glu	Lys	Lys	Asn	Lys	Thr	Leu	Glu	Tyr
				560					565					570
Lys	Ile	Glu	Ile	Leu	Glu	Lys	Thr	Thr	Thr	Ile	Tyr	Glu	Glu	Asp
				575					580					585
Lys	Arg	Asn	Leu	Gln	Gln	Glu	Leu	Glu	Thr	Gln	Asn	Gln	Lys	Leu

Gln	Arg	Gln	Phe	Ser	Asp	Lys	Arg	Arg	Leu	Glu	Ala	Arg	Leu	Gln
				605					595					600
Gly	Met	Val	Thr	Glu	Thr	Thr	Met	Lys	Trp	Glu	Lys	Glu	Cys	Glu
				620					610					615
Arg	Arg	Val	Ala	Ala	Lys	Gln	Leu	Glu	Met	Gln	Asn	Lys	Leu	Trp
				635					625					630
Val	Lys	Asp	Glu	Lys	Leu	Lys	Gln	Leu	Lys	Ala	Ile	Val	Thr	Glu
				650					640					645
Pro	Lys	Thr	Glu	Lys	Pro	Glu	Arg	Pro	Ser	Arg	Glu	Arg	Asp	Arg
				665					655					660
Glu	Lys	Val	Thr	Gln	Arg	Ser	Val	Ser	Pro	Ser	Pro	Val	Pro	Leu
				680					670					675
Leu	Phe	Gln	Pro	Asp	Gln	Asn	Ala	Pro	Pro	Ile	Arg	Leu	Arg	His
				695					685					690
Arg	Arg	Ser	Arg	Ser	Ala	Gly	Asp	Arg	Trp	Val	Asp	His	Lys	Pro
				710					700					705
Ala	Ser	Asn	Met	Gln	Thr	Glu	Thr	Val	Met	Gln	Pro	His	Val	Pro
				725					715					720
His	Ala	Ile	Thr	Val	Ser	Val	Ala	Asn	Glu	Lys	Ala	Leu	Ala	Lys
				740					730					735
Cys	Glu	Lys	Tyr	Met	Leu	Thr	His	Gln	Glu	Leu	Ala	Ser	Asp	Gly
				755					745					750
Glu	Ile	Glu	Thr	Lys	Leu	Ile	Lys	Gly	Asp	Ile	Tyr	Lys	Thr	Arg
				770					760					765
Gly	Gly	Gly	Gln	Ser	Val	Gln	Phe	Thr	Asp	Ile	Glu	Thr	Leu	Lys
				785					775					780
Gln	Glu	Ser	Pro	Asn	Gly	Ser	Arg	Lys	Arg	Arg	Ser	Ser	Thr	Val
				800					790					795
Ala	Pro	Ala	Gln	Pro	Asp	Gly	Ala	Glu	Ser	Glu	Trp	Thr	Asp	Val
				815					805					810
Glu	Thr	Arg	Cys	Ser	Val	Ala	Val	Glu	Met	Arg	Ala	Gly	Ser	Gln
				830					820					825
Leu	Gly	Pro	Gly	Tyr	Gln	His	His	Ala	Gln	Pro	Lys	Arg	Lys	Lys
				845					835					840
Pro									850					855

&lt;210&gt; 30

&lt;211&gt; 1056

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 6437362CD1

&lt;400&gt; 30

Met	Ala	Cys	Pro	Ala	Leu	Gly	Leu	Glu	Ala	Leu	Gln	Pro	Leu	Gln
1				5					10					15
Pro	Glu	Pro	Pro	Pro	Glu	Pro	Ala	Phe	Ser	Glu	Ala	Gln	Lys	Trp
				20					25					30
Ile	Glu	Gln	Val	Thr	Gly	Arg	Ser	Phe	Gly	Asp	Lys	Asp	Phe	Arg
				35					40					45
Thr	Gly	Leu	Glu	Asn	Gly	Ile	Leu	Leu	Cys	Glu	Leu	Leu	Asn	Ala
				50					55					60
Ile	Lys	Pro	Gly	Leu	Val	Lys	Lys	Ile	Asn	Arg	Leu	Pro	Thr	Pro
				65					70					75
Ile	Ala	Gly	Leu	Asp	Asn	Ile	Ile	Leu	Phe	Leu	Arg	Gly	Cys	Lys
				80					85					90
Glu	Leu	Gly	Leu	Lys	Glu	Ser	Gln	Leu	Phe	Asp	Pro	Ser	Asp	Leu
				95					100					105
Gln	Asp	Thr	Ser	Asn	Arg	Val	Thr	Val	Lys	Ser	Leu	Asp	Tyr	Ser
				110					115					120
Arg	Lys	Leu	Lys	Asn	Val	Leu	Val	Thr	Ile	Tyr	Trp	Leu	Gly	Lys
				125					130					135
Ala	Ala	Asn	Ser	Cys	Thr	Ser	Tyr	Ser	Gly	Thr	Thr	Leu	Asn	Leu

Lys	Glu	Phe	Glu	140	Gly	Leu	Leu	Ala	Gln	145	Met	Arg	Lys	Asp	Thr	150	Asp
Asp	Ile	Glu	Ser	155	Pro	Lys	Arg	Ser	Ile	160	Arg	Asp	Ser	Gly	Tyr	165	Ile
Asp	Cys	Trp	Asp	170	Ser	Glu	Arg	Ser	Asp	175	Ser	Leu	Ser	Pro	Pro	180	Arg
His	Gly	Arg	Asp	185	Asp	Ser	Phe	Asp	Ser	190	Leu	Asp	Ser	Phe	Gly	195	Ser
Arg	Ser	Arg	Gln	200	Thr	Pro	Ser	Pro	Asp	205	Val	Val	Leu	Arg	Gly	210	Ser
Ser	Asp	Gly	Arg	215	Gly	Ser	Asp	Ser	Glu	220	Ser	Asp	Leu	Pro	His	225	Arg
Lys	Leu	Pro	Asp	230	Val	Lys	Lys	Asp	Asp	235	Met	Ser	Ala	Arg	Arg	240	Thr
Ser	His	Gly	Glu	245	Pro	Lys	Ser	Ala	Val	250	Pro	Phe	Asn	Gln	Tyr	255	Leu
Pro	Asn	Lys	Ser	260	Asn	Gln	Thr	Ala	Tyr	265	Val	Pro	Ala	Pro	Leu	270	Arg
Lys	Lys	Lys	Ala	275	Glu	Arg	Glu	Glu	Tyr	280	Arg	Lys	Ser	Trp	Ser	285	Thr
Ala	Thr	Ser	Pro	290	Leu	Gly	Gly	Glu	Arg	295	Pro	Phe	Arg	Tyr	Gly	300	Pro
Arg	Thr	Pro	Val	305	Ser	Asp	Asp	Ala	Glu	310	Ser	Thr	Ser	Met	Phe	315	Asp
Met	Arg	Cys	Glu	320	Glu	Glu	Ala	Ala	Val	325	Gln	Pro	His	Ser	Arg	330	Ala
Arg	Gln	Glu	Gln	335	Leu	Gln	Leu	Ile	Asn	340	Asn	Gln	Leu	Arg	Glu	345	Glu
Asp	Asp	Lys	Trp	350	Gln	Asp	Asp	Leu	Ala	355	Arg	Trp	Lys	Ser	Arg	360	Arg
Arg	Ser	Val	Ser	365	Gln	Asp	Leu	Ile	Lys	370	Lys	Glu	Glu	Glu	Arg	375	Lys
Lys	Met	Glu	Lys	380	Leu	Leu	Ala	Gly	Glu	385	Asp	Gly	Thr	Ser	Glu	390	Arg
Arg	Lys	Ser	Ile	395	Lys	Thr	Tyr	Arg	Glu	400	Ile	Val	Gln	Glu	Lys	405	Glu
Arg	Arg	Glu	Arg	410	Glu	Leu	His	Glu	Ala	415	Tyr	Lys	Asn	Ala	Arg	420	Ser
Gln	Glu	Glu	Ala	425	Glu	Gly	Ile	Leu	Gln	430	Gln	Tyr	Ile	Glu	Arg	435	Phe
Thr	Ile	Ser	Glu	440	Ala	Val	Leu	Glu	Arg	445	Leu	Glu	Met	Pro	Lys	450	Ile
Leu	Glu	Arg	Ser	455	His	Ser	Thr	Glu	Pro	460	Asn	Leu	Ser	Ser	Phe	465	Leu
Asn	Asp	Pro	Asn	470	Pro	Met	Lys	Tyr	Leu	475	Arg	Gln	Gln	Ser	Leu	480	Pro
Pro	Pro	Lys	Phe	485	Thr	Ala	Thr	Val	Glu	490	Thr	Thr	Ile	Ala	Arg	495	Ala
Ser	Val	Leu	Asp	500	Thr	Ser	Met	Ser	Ala	505	Gly	Ser	Gly	Ser	Pro	510	Ser
Lys	Thr	Val	Thr	515	Pro	Lys	Ala	Val	Pro	520	Met	Leu	Thr	Pro	Lys	525	Pro
Tyr	Ser	Gln	Pro	530	Lys	Asn	Ser	Gln	Asp	535	Val	Leu	Lys	Thr	Phe	540	Lys
Val	Asp	Gly	Lys	545	Val	Ser	Val	Asn	Gly	550	Glu	Thr	Val	His	Arg	555	Glu
Glu	Glu	Lys	Glu	560	Arg	Glu	Cys	Pro	Thr	565	Val	Ala	Pro	Ala	His	570	Ser
Leu	Thr	Lys	Ser	575	Gln	Met	Phe	Glu	Gly	580	Val	Ala	Arg	Val	His	585	Gly
Ser	Pro	Leu	Glu	590	Leu	Lys	Gln	Asp	Asn	595	Gly	Ser	Ile	Glu	Ile	600	Asn
Ile	Lys	Lys	Pro	605	Asn	Ser	Val	Pro	Gln	610	Glu	Leu	Ala	Ala	Thr	615	Thr
Glu	Lys	Thr	Glu	620	Pro	Asn	Ser	Gln	Glu	625	Asp	Lys	Asn	Asp	Gly	630	Gly
				635						640						645	

Lys	Ser	Arg	Lys	Gly	Asn	Ile	Glu	Leu	Ala	Ser	Ser	Glu	Pro	Gln
				650					655					660
His	Phe	Thr	Thr	Thr	Val	Thr	Arg	Cys	Ser	Pro	Thr	Val	Ala	Phe
				665					670					675
Val	Glu	Phe	Pro	Ser	Ser	Pro	Gln	Leu	Lys	Asn	Asp	Val	Ser	Glu
				680					685					690
Glu	Lys	Asp	Gln	Lys	Lys	Pro	Glu	Asn	Glu	Met	Ser	Gly	Lys	Val
				695					700					705
Glu	Leu	Val	Leu	Ser	Gln	Lys	Val	Val	Lys	Pro	Lys	Ser	Pro	Glu
				710					715					720
Pro	Glu	Ala	Thr	Leu	Thr	Phe	Pro	Phe	Leu	Asp	Lys	Met	Pro	Glu
				725					730					735
Ala	Asn	Gln	Leu	His	Leu	Pro	Asn	Leu	Asn	Ser	Gln	Val	Asp	Ser
				740					745					750
Pro	Ser	Ser	Glu	Lys	Ser	Pro	Val	Thr	Thr	Pro	Phe	Lys	Phe	Trp
				755					760					765
Ala	Trp	Asp	Pro	Glu	Glu	Glu	Arg	Arg	Arg	Gln	Glu	Lys	Trp	Gln
				770					775					780
Gln	Glu	Gln	Glu	Arg	Leu	Leu	Gln	Glu	Arg	Tyr	Gln	Lys	Glu	Gln
				785					790					795
Asp	Lys	Leu	Lys	Glu	Glu	Trp	Glu	Lys	Ala	Gln	Lys	Glu	Val	Glu
				800					805					810
Glu	Glu	Glu	Arg	Arg	Tyr	Tyr	Glu	Glu	Glu	Arg	Lys	Ile	Ile	Glu
				815					820					825
Asp	Thr	Val	Val	Pro	Phe	Thr	Val	Ser	Ser	Ser	Ser	Ala	Asp	Gln
				830					835					840
Leu	Ser	Thr	Ser	Ser	Ser	Met	Thr	Glu	Gly	Ser	Gly	Thr	Met	Asn
				845					850					855
Lys	Ile	Asp	Leu	Gly	Asn	Cys	Gln	Asp	Glu	Lys	Gln	Asp	Arg	Arg
				860					865					870
Trp	Lys	Lys	Ser	Phe	Gln	Gly	Asp	Asp	Ser	Asp	Leu	Leu	Leu	Lys
				875					880					885
Thr	Arg	Glu	Ser	Asp	Arg	Leu	Glu	Glu	Lys	Gly	Ser	Leu	Thr	Glu
				890					895					900
Gly	Ala	Leu	Ala	His	Ser	Gly	Asn	Pro	Val	Ser	Lys	Gly	Val	His
				905					910					915
Glu	Asp	His	Gln	Leu	Asp	Thr	Glu	Ala	Gly	Ala	Pro	His	Cys	Gly
				920					925					930
Thr	Asn	Pro	Gln	Leu	Ala	Gln	Asp	Pro	Ser	Gln	Asn	Gln	Gln	Thr
				935					940					945
Ser	Asn	Pro	Thr	His	Ser	Ser	Glu	Asp	Val	Lys	Pro	Lys	Thr	Leu
				950					955					960
Pro	Leu	Asp	Lys	Ser	Ile	Asn	His	Gln	Ile	Glu	Ser	Pro	Ser	Glu
				965					970					975
Arg	Arg	Lys	Ser	Ile	Ser	Gly	Lys	Lys	Leu	Cys	Ser	Ser	Cys	Gly
				980					985					990
Leu	Pro	Leu	Gly	Lys	Gly	Ala	Ala	Met	Ile	Ile	Glu	Thr	Leu	Asn
				995					1000					1005
Leu	Tyr	Phe	His	Ile	Gln	Cys	Phe	Arg	Cys	Gly	Ile	Cys	Lys	Gly
				1010					1015					1020
Gln	Leu	Gly	Asp	Ala	Val	Ser	Gly	Thr	Asp	Val	Arg	Ile	Arg	Asn
				1025					1030					1035
Gly	Leu	Leu	Asn	Cys	Asn	Asp	Cys	Tyr	Met	Arg	Ser	Arg	Ser	Ala
				1040					1045					1050
Gly	Gln	Pro	Thr	Thr	Leu									
				1055										

&lt;210&gt; 31

&lt;211&gt; 1569

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 4173970CD1

&lt;400&gt; 31

Met	Val	Val	Pro	Pro	Gln	Glu	Pro	Asp	Arg	Thr	Ser	Gln	Glu	Asn
1				5					10					15
Ser	Pro	Ala	Leu	Leu	Gly	Val	Gln	Lys	Ala	Val	Ser	Thr	Arg	Val
				20					25					30
Pro	Thr	Gly	Ser	Asn	Ser	Ser	Ser	Gln	Thr	Thr	Glu	Cys	Leu	Thr
				35					40					45
Pro	Glu	Ser	Cys	Ser	Gln	Thr	Thr	Ser	Asn	Val	Ala	Ser	Gln	Ser
				50					55					60
Met	Pro	Pro	Val	Tyr	Pro	Ser	Val	Asp	Ile	Asp	Ala	His	Thr	Glu
				65					70					75
Ser	Asn	His	Asp	Thr	Ala	Leu	Thr	Leu	Ala	Cys	Ala	Gly	Gly	His
				80					85					90
Glu	Glu	Leu	Val	Ser	Val	Leu	Ile	Ala	Arg	Asp	Ala	Lys	Ile	Glu
				95					100					105
His	Arg	Asp	Lys	Lys	Gly	Phe	Thr	Pro	Leu	Ile	Leu	Ala	Ala	Thr
				110					115					120
Ala	Gly	His	Val	Gly	Val	Val	Glu	Ile	Leu	Leu	Asp	Lys	Gly	Gly
				125					130					135
Asp	Ile	Glu	Ala	Gln	Ser	Glu	Arg	Thr	Lys	Asp	Thr	Pro	Leu	Ser
				140					145					150
Leu	Ala	Cys	Ser	Gly	Gly	Arg	Gln	Glu	Val	Val	Asp	Leu	Leu	Leu
				155					160					165
Ala	Arg	Gly	Ala	Asn	Lys	Glu	His	Arg	Asn	Val	Ser	Asp	Tyr	Thr
				170					175					180
Pro	Leu	Ser	Leu	Ala	Ala	Ser	Gly	Gly	Tyr	Val	Asn	Ile	Ile	Lys
				185					190					195
Ile	Leu	Leu	Asn	Ala	Gly	Ala	Glu	Ile	Asn	Ser	Arg	Thr	Gly	Ser
				200					205					210
Lys	Leu	Gly	Ile	Ser	Pro	Leu	Met	Leu	Ala	Ala	Met	Asn	Gly	His
				215					220					225
Val	Pro	Ala	Val	Lys	Leu	Leu	Leu	Asp	Met	Gly	Ser	Asp	Ile	Asn
				230					235					240
Ala	Gln	Ile	Glu	Thr	Asn	Arg	Asn	Thr	Ala	Leu	Thr	Leu	Ala	Cys
				245					250					255
Phe	Gln	Gly	Arg	Ala	Glu	Val	Val	Ser	Leu	Leu	Leu	Asp	Arg	Lys
				260					265					270
Ala	Asn	Val	Glu	His	Arg	Ala	Lys	Thr	Gly	Leu	Thr	Pro	Leu	Met
				275					280					285
Glu	Ala	Ala	Ser	Gly	Gly	Tyr	Ala	Glu	Val	Gly	Arg	Val	Leu	Leu
				290					295					300
Asp	Lys	Gly	Ala	Asp	Val	Asn	Ala	Pro	Pro	Val	Pro	Ser	Ser	Arg
				305					310					315
Asp	Thr	Ala	Leu	Thr	Ile	Ala	Ala	Asp	Lys	Gly	His	Tyr	Lys	Phe
				320					325					330
Cys	Glu	Leu	Leu	Ile	His	Arg	Gly	Ala	His	Ile	Asp	Val	Arg	Asn
				335					340					345
Lys	Lys	Gly	Asn	Thr	Pro	Leu	Trp	Leu	Ala	Ser	Asn	Gly	Gly	His
				350					355					360
Phe	Asp	Val	Val	Gln	Leu	Leu	Val	Gln	Ala	Gly	Ala	Asp	Val	Asp
				365					370					375
Ala	Ala	Asp	Asn	Arg	Lys	Ile	Thr	Pro	Leu	Met	Ser	Ala	Phe	Arg
				380					385					390
Lys	Gly	His	Val	Lys	Val	Val	Gln	Tyr	Leu	Val	Lys	Glu	Val	Asn
				395					400					405
Gln	Phe	Pro	Ser	Asp	Ile	Glu	Cys	Met	Arg	Tyr	Ile	Ala	Thr	Ile
				410					415					420
Thr	Asp	Lys	Glu	Leu	Leu	Lys	Lys	Cys	His	Gln	Cys	Val	Glu	Thr
				425					430					435
Ile	Val	Lys	Ala	Lys	Asp	Gln	Gln	Ala	Ala	Glu	Ala	Asn	Lys	Asn
				440					445					450
Ala	Ser	Ile	Leu	Leu	Lys	Glu	Leu	Asp	Leu	Glu	Lys	Ser	Arg	Glu
				455					460					465
Glu	Ser	Arg	Lys	Gln	Ala	Leu	Ala	Ala	Lys	Arg	Glu	Lys	Arg	Lys
				470					475					480
Glu	Lys	Arg	Lys	Lys	Lys	Lys	Glu	Glu	Gln	Lys	Arg	Lys	Gln	Glu
				485					490					495
Glu	Asp	Glu	Glu	Asn	Lys	Pro	Lys	Glu	Asn	Ser	Glu	Leu	Pro	Glu

Asp	Glu	Asp	Glu	500	Glu	Glu	Asn	Asp	Glu	505	Asp	Val	Glu	Gln	Glu	510
Pro	Ile	Glu	Pro	515	Pro	Ser	Ala	Thr	Thr	520	Thr	Thr	Ile	Gly	Ile	525
Ser	Ala	Thr	Ser	530	Ala	Thr	Phe	Thr	Asn	535	Val	Phe	Gly	Lys	Lys	540
Ala	Asn	Val	Val	545	Thr	Pro	Ser	Thr	Thr	550	Asn	Arg	Lys	Asn	Lys	555
Asn	Lys	Thr	Lys	560	Glu	Thr	Pro	Pro	Thr	565	Ala	His	Leu	Ile	Leu	570
Glu	Gln	His	Met	575	Ser	Leu	Ala	Gln	Gln	580	Lys	Ala	Asp	Lys	Asn	585
Ile	Asn	Gly	Glu	590	Pro	Arg	Gly	Gly	Gly	595	Ala	Gly	Gly	Asn	Ser	600
Ser	Asp	Asn	Leu	605	Asp	Ser	Thr	Asp	Cys	610	Asn	Ser	Glu	Ser	Ser	615
Gly	Gly	Lys	Ser	620	Gln	Glu	Leu	Asn	Phe	625	Val	Met	Asp	Val	Asn	630
Ser	Lys	Tyr	Pro	635	Ser	Leu	Leu	Leu	His	640	Ser	Gln	Glu	Glu	Lys	645
Ser	Thr	Ala	Thr	650	Ser	Lys	Thr	Gln	Thr	655	Arg	Tyr	Lys	Thr	Val	660
Leu	Pro	Leu	Ser	665	Ser	Pro	Asn	Ile	Lys	670	Leu	Asn	Leu	Thr	Ser	675
Lys	Arg	Gly	Gln	680	Lys	Arg	Glu	Glu	Gly	685	Trp	Lys	Glu	Val	Val	690
Arg	Ser	Lys	Lys	695	Leu	Ser	Val	Pro	Ala	700	Ser	Val	Val	Ser	Arg	705
Met	Gly	Arg	Gly	710	Gly	Cys	Asn	Ile	Thr	715	Ala	Ile	Gln	Asp	Val	720
Gly	Ala	His	Ile	725	Asp	Val	Asp	Lys	Gln	730	Asp	Lys	Asn	Gly	Glu	735
Arg	Met	Ile	Thr	740	Ile	Arg	Gly	Gly	Thr	745	Glu	Ser	Thr	Arg	Tyr	750
Val	Gln	Leu	Ile	755	Asn	Ala	Leu	Ile	Gln	760	Asp	Pro	Ala	Lys	Glu	765
Glu	Asp	Leu	Ile	770	Pro	Lys	Asn	His	Ile	775	Arg	Thr	Pro	Ala	Ser	780
Lys	Ser	Ile	His	785	Ala	Asn	Phe	Ser	Ser	790	Gly	Val	Gly	Thr	Thr	795
Ala	Ser	Ser	Lys	800	Asn	Ala	Phe	Pro	Leu	805	Gly	Ala	Pro	Thr	Leu	810
Thr	Ser	Gln	Ala	815	Thr	Thr	Leu	Ser	Thr	820	Phe	Gln	Pro	Ala	Asn	825
Leu	Asn	Lys	Asn	830	Val	Pro	Thr	Asn	Val	835	Arg	Ser	Ser	Phe	Pro	840
Ser	Leu	Pro	Leu	845	Ala	Tyr	Pro	His	Pro	850	His	Phe	Ala	Leu	Leu	855
Ala	Gln	Thr	Met	860	Gln	Gln	Ile	Arg	His	865	Pro	Arg	Leu	Pro	Met	870
Gln	Phe	Gly	Gly	875	Thr	Phe	Ser	Pro	Ser	880	Pro	Asn	Thr	Trp	Gly	885
Phe	Pro	Val	Arg	890	Pro	Val	Asn	Pro	Gly	895	Asn	Thr	Asn	Ser	Ser	900
Lys	His	Asn	Asn	905	Thr	Ser	Arg	Leu	Pro	910	Gln	Asn	Gly	Thr	Val	915
Leu	Pro	Ser	Glu	920	Ser	Ala	Gly	Leu	Ala	925	Thr	Ala	Ser	Cys	Pro	930
Thr	Val	Ser	Ser	935	Val	Val	Ala	Ala	Ser	940	Gln	Gln	Leu	Cys	Val	945
Asn	Thr	Arg	Thr	950	Pro	Ser	Ser	Val	Arg	955	Lys	Gln	Leu	Phe	Ala	960
Val	Pro	Lys	Thr	965	Ser	Pro	Pro	Ala	Thr	970	Val	Ile	Ser	Ser	Val	975
Ser	Thr	Cys	Ser	980	Ser	Leu	Pro	Ser	Val	985	Ser	Ser	Ala	Pro	Ile	990
				995						1000						1005

Ser	Gly	Gln	Ala	Pro	Thr	Thr	Phe	Leu	Pro	Ala	Ser	Thr	Ser	Gln
				1010					1015					1020
Ala	Gln	Leu	Ser	Ser	Gln	Lys	Met	Glu	Ser	Phe	Ser	Ala	Val	Pro
				1025					1030					1035
Pro	Thr	Lys	Glu	Lys	Val	Ser	Thr	Gln	Asp	Gln	Pro	Met	Ala	Asn
				1040					1045					1050
Leu	Cys	Thr	Pro	Ser	Ser	Thr	Ala	Asn	Ser	Cys	Ser	Ser	Ser	Ala
				1055					1060					1065
Ser	Asn	Thr	Pro	Gly	Ala	Pro	Glu	Thr	His	Pro	Ser	Ser	Ser	Pro
				1070					1075					1080
Thr	Pro	Thr	Ser	Ser	Asn	Thr	Gln	Glu	Glu	Ala	Gln	Pro	Ser	Ser
				1085					1090					1095
Val	Ser	Asp	Leu	Ser	Pro	Met	Ser	Met	Pro	Phe	Ala	Ser	Asn	Ser
				1100					1105					1110
Glu	Pro	Ala	Pro	Leu	Thr	Leu	Thr	Ser	Pro	Arg	Met	Val	Ala	Ala
				1115					1120					1125
Asp	Asn	Gln	Asp	Thr	Ser	Asn	Leu	Pro	Gln	Leu	Ala	Val	Pro	Ala
				1130					1135					1140
Pro	Arg	Val	Ser	His	Arg	Met	Gln	Pro	Arg	Gly	Ser	Phe	Tyr	Ser
				1145					1150					1155
Met	Val	Pro	Asn	Ala	Thr	Ile	His	Gln	Asp	Pro	Gln	Ser	Ile	Phe
				1160					1165					1170
Val	Thr	Asn	Pro	Val	Thr	Leu	Thr	Pro	Pro	Gln	Gly	Pro	Pro	Ala
				1175					1180					1185
Ala	Val	Gln	Leu	Ser	Ser	Ala	Val	Asn	Ile	Met	Asn	Gly	Ser	Gln
				1190					1195					1200
Met	His	Ile	Asn	Pro	Ala	Asn	Lys	Ser	Leu	Pro	Pro	Thr	Phe	Gly
				1205					1210					1215
Pro	Ala	Thr	Leu	Phe	Asn	His	Phe	Ser	Ser	Leu	Phe	Asp	Ser	Ser
				1220					1225					1230
Gln	Val	Pro	Ala	Asn	Gln	Gly	Trp	Gly	Asp	Gly	Pro	Leu	Ser	Ser
				1235					1240					1245
Arg	Val	Ala	Thr	Asp	Ala	Ser	Phe	Thr	Val	Gln	Ser	Ala	Phe	Leu
				1250					1255					1260
Gly	Asn	Ser	Val	Leu	Gly	His	Leu	Glu	Asn	Met	His	Pro	Asp	Asn
				1265					1270					1275
Ser	Lys	Ala	Pro	Gly	Phe	Arg	Pro	Pro	Ser	Gln	Arg	Val	Ser	Thr
				1280					1285					1290
Ser	Pro	Val	Gly	Leu	Pro	Ser	Ile	Asp	Pro	Ser	Gly	Ser	Ser	Pro
				1295					1300					1305
Ser	Ser	Ser	Ser	Ala	Pro	Leu	Ala	Ser	Phe	Ser	Gly	Ile	Pro	Gly
				1310					1315					1320
Thr	Arg	Val	Phe	Leu	Gln	Gly	Pro	Ala	Pro	Val	Gly	Thr	Pro	Ser
				1325					1330					1335
Phe	Asn	Arg	Gln	His	Phe	Ser	Pro	His	Pro	Trp	Thr	Ser	Ala	Ser
				1340					1345					1350
Asn	Ser	Cys	Asp	Ser	Pro	Ile	Pro	Ser	Val	Ser	Ser	Gly	Ser	Ser
				1355					1360					1365
Ser	Pro	Leu	Ser	Ala	Thr	Ser	Ala	Pro	Pro	Thr	Leu	Gly	Gln	Pro
				1370					1375					1380
Lys	Gly	Val	Ser	Ala	Ser	Gln	Asp	Arg	Lys	Ile	Pro	Pro	Pro	Ile
				1385					1390					1395
Gly	Thr	Glu	Arg	Leu	Ala	Arg	Ile	Arg	Gln	Gly	Gly	Ser	Val	Ala
				1400					1405					1410
Gln	Ala	Pro	Ala	Gly	Thr	Ser	Phe	Val	Ala	Pro	Val	Gly	His	Ser
				1415					1420					1425
Gly	Ile	Trp	Ser	Phe	Gly	Val	Asn	Ala	Val	Ser	Glu	Gly	Leu	Ser
				1430					1435					1440
Gly	Trp	Ser	Gln	Ser	Val	Met	Gly	Asn	His	Pro	Met	His	Gln	Gln
				1445					1450					1455
Leu	Ser	Asp	Pro	Ser	Thr	Phe	Ser	Gln	His	Gln	Pro	Met	Glu	Arg
				1460					1465					1470
Asp	Asp	Ser	Gly	Met	Val	Ala	Pro	Ser	Asn	Ile	Phe	His	Gln	Pro
				1475					1480					1485
Met	Ala	Ser	Gly	Phe	Val	Asp	Phe	Ser	Lys	Gly	Leu	Pro	Ile	Ser
				1490					1495					1500
Met	Tyr	Gly	Gly	Thr	Ile	Ile	Pro	Ser	His	Pro	Gln	Leu	Ala	Asp



Val	Pro	Gly	Gly	Pro	Leu	Phe	Asn	Gly	Leu	His	Asn	Pro	Asp	Pro	1505	1510	1515
Ala	Trp	Asn	Pro	Met	Ile	Lys	Val	Ile	Gln	Asn	Ser	Thr	Glu	Cys	1520	1525	1530
Thr	Asp	Ala	Gln	Gln	Ile	Trp	Pro	Gly	Thr	Trp	Ala	Pro	His	Ile	1535	1540	1545
Gly	Asn	Met	His	Leu	Lys	Tyr	Val	Asn							1550	1555	1560
															1565		

&lt;210&gt; 32

&lt;211&gt; 680

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2772751CD1

&lt;400&gt; 32

Met	Val	Phe	Leu	Leu	Leu	Thr	Thr	Leu	Leu	Leu	Leu	Ile	Gly	Val	1	5	10	15
Val	Cys	Ala	Phe	Val	Thr	Asn	Gln	Arg	Thr	His	Glu	Gln	Met	Gly	20	25	30	35
Pro	Ser	Ile	Glu	Ala	Met	Pro	Glu	Thr	Leu	Leu	Ser	Leu	Trp	Gly	40	45	50	55
Leu	Val	Ser	Asp	Val	Pro	Gln	Glu	Leu	Gln	Ala	Val	Ala	Gln	Gln	60	65	70	75
Phe	Ser	Leu	Pro	Gln	Glu	Gln	Val	Ser	Glu	Glu	Leu	Asp	Gly	Val	80	85	90	95
Gly	Val	Ser	Ile	Gly	Ser	Ala	Ile	His	Thr	Gln	Leu	Arg	Ser	Ser	100	105	110	115
Val	Tyr	Pro	Leu	Leu	Ala	Ala	Val	Gly	Ser	Leu	Gly	Gln	Val	Leu	120	125	130	135
Gln	Val	Ser	Val	His	His	Leu	Gln	Thr	Leu	Asn	Ala	Thr	Val	Val	140	145	150	155
Glu	Leu	Gln	Ala	Gly	Gln	Gln	Asp	Leu	Glu	Pro	Ala	Ile	Arg	Glu	160	165	170	175
His	Arg	Asp	Arg	Leu	Leu	Glu	Leu	Leu	Gln	Glu	Ala	Arg	Cys	Gln	180	185	190	195
Gly	Asp	Cys	Ala	Gly	Ala	Leu	Ser	Trp	Ala	Arg	Thr	Leu	Glu	Leu	200	205	210	215
Gly	Ala	Asp	Phe	Ser	Gln	Val	Pro	Ser	Val	Asp	His	Val	Leu	His	220	225	230	235
Gln	Leu	Lys	Gly	Val	Pro	Glu	Ala	Asn	Phe	Ser	Ser	Met	Val	Gln	240	245	250	255
Glu	Glu	Asn	Ser	Thr	Phe	Asn	Ala	Leu	Pro	Ala	Leu	Ala	Ala	Met	260	265	270	275
Gln	Thr	Ser	Ser	Val	Val	Gln	Glu	Leu	Lys	Lys	Ala	Val	Ala	Gln	280	285	290	295
Gln	Pro	Glu	Gly	Val	Arg	Thr	Leu	Ala	Glu	Gly	Phe	Pro	Gly	Leu	300	305	310	315
Glu	Ala	Ala	Ser	Arg	Trp	Ala	Gln	Ala	Leu	Gln	Glu	Val	Glu	Glu	320	325	330	335
Ser	Ser	Arg	Pro	Tyr	Leu	Gln	Glu	Val	Gln	Arg	Tyr	Glu	Thr	Tyr	340	345	350	355
Arg	Trp	Ile	Val	Gly	Cys	Val	Leu	Cys	Ser	Val	Val	Leu	Phe	Val	360	365	370	375
Val	Leu	Cys	Asn	Leu	Leu	Gly	Leu	Asn	Leu	Gly	Ile	Trp	Gly	Leu	380	385	390	395
Ser	Ala	Arg	Asp	Asp	Pro	Ser	His	Pro	Glu	Ala	Lys	Gly	Glu	Ala	400	405	410	415
Gly	Ala	Arg	Phe	Leu	Met	Ala	Gly	Val	Gly	Leu	Ser	Phe	Leu	Phe	420	425	430	435
Ala	Ala	Pro	Leu	Ile	Leu	Leu	Val	Phe	Ala	Thr	Phe	Leu	Val	Gly	440	445	450	455
Gly	Asn	Val	Gln	Thr	Leu	Val	Cys	Gln	Ser	Trp	Glu	Asn	Ser	Glu	460	465	470	475

Leu Phe Glu Phe	350	Ala Asp Thr Pro Gly	355	Asn Leu Pro Pro Ser	360
Asn Leu Ser Gln	365	Leu Gly Leu Arg	370	Lys Asn Ile Ser Ile	375
Gln Ala Tyr Gln	380	Gln Cys Lys Glu Gly	385	Ala Ala Leu Trp Thr	390
Leu Gln Leu Asn	395	Asp Ser Tyr Asp Leu	400	Glu His Leu Asp Ile	405
Asn Gln Tyr Thr	410	Asn Lys Leu Arg Gln	415	Glu Leu Gln Ser Leu	420
Val Asp Thr Gln	425	Ser Leu Asp Leu Leu	430	Ser Ser Ala Ala Arg	435
Asp Leu Glu Ala	440	Leu Gln Ser Ser Gly	445	Leu Gln Arg Ile His	450
Pro Asp Phe Leu	455	Val Gln Ile Gln Arg	460	Pro Val Val Lys Thr	465
Met Glu Gln Leu	470	Ala Gln Glu Leu Gln	475	Leu Ala Gln Ala Gln	480
Asp Asn Ser Val	485	Leu Gly Gln Arg Leu	490	Gln Glu Glu Ala Gln	495
Leu Arg Asn Leu	500	His Gln Glu Lys Val	505	Val Pro Gln Gln Ser	510
Val Ala Lys Leu	515	Asn Leu Ser Val Arg	520	Ala Leu Glu Ser Ser	525
Pro Asn Leu Gln	530	Leu Glu Thr Ser Asp	535	Val Leu Ala Asn Val	540
Tyr Leu Lys Gly	545	Glu Leu Pro Ala Trp	550	Ala Ala Arg Ile Leu	555
Asn Val Ser Glu	560	Cys Phe Leu Ala Arg	565	Glu Met Gly Tyr Phe	570
Gln Tyr Val Ala	575	Trp Val Arg Glu Glu	580	Val Thr Gln Arg Ile	585
Thr Cys Gln Pro	590	Leu Ser Gly Ala Leu	595	Asp Asn Ser Arg Val	600
Leu Cys Asp Met	605	Met Ala Asp Pro Trp	610	Asn Ala Phe Trp Phe	615
Leu Ala Trp Cys	620	Thr Phe Phe Leu Ile	625	Pro Ser Ile Ile Phe	630
Val Lys Thr Ser	635	Lys Tyr Phe Arg Pro	640	Ile Arg Lys Arg Leu	645
Ser Thr Ser Ser	650	Glu Thr Gln Leu	655	Phe His Ile Pro Arg	660
Thr Ser Leu Lys	665	Leu	670		675
	680				

&lt;210&gt; 33

&lt;211&gt; 590

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2793768CD1

&lt;400&gt; 33

Met Ser Ser Ala Cys	5	Asp Ala Gly Asp His	10	Tyr Pro Leu His Leu	15
Leu Val Trp Lys Asn	20	Asp Tyr Arg Gln Leu	25	Glu Lys Glu Leu Gln	30
Gly Gln Asn Val Glu	35	Ala Val Asp Pro Arg	40	Gly Arg Thr Leu Leu	45
His Leu Ala Val Ser	50	Leu Gly His Leu Glu	55	Ser Ala Arg Val Leu	60
Leu Arg His Lys Ala	65	Asp Val Thr Lys Glu	70	Asn Arg Gln Gly Trp	75
Thr Val Leu His Glu		Ala Val Ser Thr Gly		Asp Pro Glu Met Val	

				80					85					90
Tyr	Thr	Val	Leu	Gln	His	Arg	Asp	Tyr	His	Asn	Thr	Ser	Met	Ala
				95					100					105
Leu	Glu	Gly	Val	Pro	Glu	Leu	Leu	Gln	Lys	Ile	Leu	Glu	Ala	Pro
				110					115					120
Asp	Phe	Tyr	Val	Gln	Met	Lys	Trp	Glu	Phe	Thr	Ser	Trp	Val	Pro
				125					130					135
Leu	Val	Ser	Arg	Ile	Cys	Pro	Asn	Asp	Val	Cys	Arg	Ile	Trp	Lys
				140					145					150
Ser	Gly	Ala	Lys	Leu	Arg	Val	Asp	Ile	Thr	Leu	Leu	Gly	Phe	Glu
				155					160					165
Asn	Met	Ser	Trp	Ile	Arg	Gly	Arg	Arg	Ser	Phe	Ile	Phe	Lys	Gly
				170					175					180
Glu	Asp	Asn	Trp	Ala	Glu	Leu	Met	Glu	Val	Asn	His	Asp	Asp	Lys
				185					190					195
Val	Val	Thr	Thr	Glu	Arg	Phe	Asp	Leu	Ser	Gln	Glu	Met	Glu	Arg
				200					205					210
Leu	Thr	Leu	Asp	Leu	Met	Lys	Pro	Lys	Ser	Arg	Glu	Val	Glu	Arg
				215					220					225
Arg	Leu	Thr	Ser	Pro	Val	Ile	Asn	Thr	Ser	Leu	Asp	Thr	Lys	Asn
				230					235					240
Ile	Ala	Phe	Glu	Arg	Thr	Lys	Ser	Gly	Phe	Trp	Gly	Trp	Arg	Thr
				245					250					255
Asp	Lys	Ala	Glu	Val	Val	Asn	Gly	Tyr	Glu	Ala	Lys	Val	Tyr	Thr
				260					265					270
Val	Asn	Asn	Val	Asn	Val	Ile	Thr	Lys	Ile	Arg	Thr	Glu	His	Leu
				275					280					285
Thr	Glu	Glu	Glu	Lys	Lys	Arg	Tyr	Lys	Ala	Asp	Arg	Asn	Pro	Leu
				290					295					300
Glu	Ser	Leu	Leu	Gly	Thr	Val	Glu	His	Gln	Phe	Gly	Ala	Gln	Gly
				305					310					315
Asp	Leu	Thr	Thr	Glu	Cys	Ala	Thr	Ala	Asn	Asn	Pro	Thr	Ala	Ile
				320					325					330
Thr	Pro	Asp	Glu	Tyr	Phe	Asn	Glu	Glu	Phe	Asp	Leu	Lys	Asp	Arg
				335					340					345
Asp	Ile	Gly	Arg	Pro	Lys	Glu	Leu	Thr	Ile	Arg	Thr	Gln	Lys	Phe
				350					355					360
Lys	Ala	Met	Leu	Trp	Met	Cys	Glu	Glu	Phe	Pro	Leu	Ser	Leu	Val
				365					370					375
Glu	Gln	Val	Ile	Pro	Ile	Ile	Asp	Leu	Met	Ala	Arg	Thr	Ser	Ala
				380					385					390
His	Phe	Ala	Arg	Leu	Arg	Asp	Phe	Ile	Lys	Leu	Glu	Phe	Pro	Pro
				395					400					405
Gly	Phe	Pro	Val	Lys	Ile	Glu	Ile	Pro	Leu	Phe	His	Val	Leu	Asn
				410					415					420
Ala	Arg	Ile	Thr	Phe	Gly	Asn	Val	Asn	Gly	Cys	Ser	Thr	Ala	Glu
				425					430					435
Glu	Ser	Val	Ser	Gln	Asn	Val	Glu	Gly	Thr	Gln	Ala	Asp	Ser	Ala
				440					445					450
Ser	His	Ile	Thr	Asn	Phe	Glu	Val	Asp	Gln	Ser	Val	Phe	Glu	Ile
				455					460					465
Pro	Glu	Ser	Tyr	Tyr	Val	Gln	Asp	Asn	Gly	Arg	Asn	Val	His	Leu
				470					475					480
Gln	Asp	Glu	Asp	Tyr	Glu	Ile	Met	Gln	Phe	Ala	Ile	Gln	Gln	Ser
				485					490					495
Leu	Leu	Glu	Ser	Ser	Arg	Ser	Gln	Glu	Leu	Ser	Gly	Pro	Ala	Ser
				500					505					510
Asn	Gly	Gly	Ile	Ser	Gln	Thr	Asn	Thr	Tyr	Asp	Ala	Gln	Tyr	Glu
				515					520					525
Arg	Ala	Ile	Gln	Glu	Ser	Leu	Leu	Thr	Ser	Thr	Glu	Gly	Leu	Cys
				530					535					540
Pro	Ser	Ala	Leu	Ser	Glu	Thr	Ser	Arg	Phe	Asp	Asn	Asp	Leu	Gln
				545					550					555
Leu	Ala	Met	Glu	Leu	Ser	Ala	Lys	Glu	Leu	Glu	Glu	Trp	Glu	Leu
				560					565					570
Arg	Leu	Gln	Glu	Glu	Glu	Ala	Glu	Leu	Gln	Gln	Val	Leu	Gln	Leu
				575					580					585

Ser Leu Thr Asp Lys  
590

<210> 34  
<211> 315  
<212> PRT  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 3035248CD1

<400> 34  
Met Val Gly Pro Trp Val Tyr Leu Val Ala Ala Val Leu Leu Ile  
1 5 10 15  
Gly Leu Ile Leu Phe Leu Thr Arg Ser Arg Gly Arg Ala Ala Ala  
20 25 30  
Ala Asp Gly Glu Pro Leu His Asn Glu Glu Arg Ala Gly Ala  
35 40 45  
Gly Gln Val Gly Arg Ser Leu Pro Gln Glu Ser Glu Glu Gln Arg  
50 55 60  
Thr Gly Ser Arg Pro Arg Arg Arg Arg Asp Leu Gly Ser Arg Leu  
65 70 75  
Gln Ala Gln Arg Arg Ala Gln Arg Val Ala Trp Glu Asp Gly Asp  
80 85 90  
Glu Asn Val Gly Gln Thr Val Ile Pro Ala Gln Glu Glu Glu Gly  
95 100 105  
Ile Glu Lys Pro Ala Glu Val His Pro Thr Gly Lys Ile Gly Ala  
110 115 120  
Lys Lys Leu Arg Lys Leu Glu Glu Lys Gln Ala Arg Lys Ala Gln  
125 130 135  
Arg Glu Ala Glu Glu Ala Glu Arg Glu Glu Arg Lys Arg Leu Glu  
140 145 150  
Ser Gln Arg Glu Ala Glu Trp Lys Lys Glu Glu Glu Arg Leu Arg  
155 160 165  
Leu Lys Glu Glu Gln Lys Glu Glu Glu Glu Arg Lys Ala Gln Glu  
170 175 180  
Glu Gln Ala Arg Arg Asp His Glu Glu Tyr Leu Lys Leu Lys Glu  
185 190 195  
Ala Phe Val Val Glu Glu Glu Gly Val Ser Glu Thr Met Thr Glu  
200 205 210  
Glu Gln Ser His Ser Phe Leu Thr Glu Phe Ile Asn Tyr Ile Lys  
215 220 225  
Lys Ser Lys Val Val Leu Leu Glu Asp Leu Ala Phe Gln Met Gly  
230 235 240  
Leu Arg Thr Gln Asp Ala Ile Asn Arg Ile Gln Asp Leu Leu Thr  
245 250 255  
Glu Gly Thr Leu Thr Gly Val Ile Asp Asp Arg Gly Lys Phe Ile  
260 265 270  
Tyr Ile Thr Pro Glu Glu Leu Ala Ala Val Ala Asn Phe Ile Arg  
275 280 285  
Gln Arg Gly Arg Val Ser Ile Thr Glu Leu Ala Gln Ala Ser Asn  
290 295 300  
Ser Leu Ile Ser Trp Gly Gln Asp Leu Pro Ala Gln Ala Ser Ala  
305 310 315

<210> 35  
<211> 2345  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 1889577CB1

<400> 35

```

ggagccggggc cagagcacca gggcagggcc cggcgcccg cgtccccgcac cctcgtcctc 60
acagacgcca cagccatggc catgatgggtg tttccgcggg aggagaagct gagccaggat 120
gagatcgtgc tgggcaccaa ggctgtcatc cagggactgg agactctgcg tggggagcat 180
cgtgccctgc tggctcctct ggttgcaact gaggccggcg aacccgagcc tggctcgagc 240
gagcgctgca tctctctgcg tgcctccctg gaagccattg agcttgggct gggggaggcc 300
caggtgatct tggcattgtc gagccacctg ggggctgtag aatcagagaa gcagaagctg 360
cgggcgcagg tgcggcgtct ggtgcaggag aaccagtggc tgcgtgagga gctggcgggg 420
acacagcaga agctgcagcg cagtgcagcg gccgtggccc agctcgagga ggagaagcag 480
cacttgctgt tcatgagcca gatccgcaag ttggatgaag acgcctcccc taacgaggag 540
aaggggggacg tccccaaaga cacactggat gacctgttcc ccaatgagga tgagcagagc 600
ccagcccccta gcccaggagg aggggatgtg tctggtcagc atggggggcta cgagatcccc 660
gccccggctcc gcaccttgca caacctgggtg atccaatacg cctcacaggg ccgctacgag 720
gtagctgtgc cactctgcaa gcaggcactc gagacctgg agaagacgtc aggccacgac 780
caccctgacg ttgccaccat gctgaacatc ctggcactgg totatcgga tcagaacaag 840
tacaaggagg ctgcccacct gctcaatgat gctctggcca tccgggagaa aacactgggc 900
aaggaccacc cagcctgtggc tgcgacacta aacaacctgg cagtcctgta tggcaagagg 960
ggcaagtaca aggaggctga gccattgtgc aagcgggcac tggagatccg ggagaaggtc 1020
ctgggcaagt ttcacccaga tgtggccaag cagctcagca acctggccct cctgtgccag 1080
aaccagggca aagctgagga ggtggaatat tactatcggc gggcactgga gatctatgct 1140
acacgcctcg ggcccgatga tcccaatgtg gccaaagacca agaacaacct ggcttccctg 1200
tacctgaagc agggcaagta ccaggatgcy gagaccttgt acaaggagat cctcacccgc 1260
gctcatgaga aagagtttgg ctctgtcaat ggggacaaca agcccatctg gatgcacgca 1320
gaggagcggg aggaaagcaa ggataagcgc cgggacagcg ccccttatgg ggaatacggc 1380
agctggtaca aggcctgtaa agtagacagc cccacagtca acaccaccct gcgcacgttg 1440
ggggccctat accggcgcca gggcaagctg gaagccgcgc acacactaga ggactgtgcc 1500
agccgtaacc gcaagcaggg tttggacccc gcaagccaga ccaagggtgg agaactgctg 1560
aaagatggca gtggcaggcg gggagacccc cgcagcagcc gagacatggc tgggggtgcc 1620
gggcctcggt ctgagctgta cctcgaggac gtgggaccta cagctgagtg gaatggggat 1680
ggcagtggct ccttgaggcg cagcggttcc tttgggaaac tccgggatgc cctgaggcgc 1740
agcagtgaga tgctggtaaa gaagctgcag gggggcacc cccaggagcc ccctaacccc 1800
aggtgaagc gggccagttc cctcaacttc ctcaacaaga gcgtggaaga gccgaccccag 1860
cctggaggca caggtctctc tgacagccgc actctcagct ccagctccat ggacctctcc 1920
cgacgaagct ccctggtggg ctaatgctga aggggcagcc agtcaccaga gcgcccacct 1980
ggcacacccc cctcacccca gccctgcgca tgggcctgct gcttgctccc cctgtctctc 2040
ccacagcccc tgtcttttct gttcaatctc agggtaacct tctcccttgt catctcagcc 2100
tgagccctgg aggtggggcc tgcccactcc agctccatcc cttattttat ccttcagca 2160
gggcctctct ccttaggttc gggccagcag ggggtgccgg ctggagtctc caccatagac 2220
tcagtggcct ggctcccca gaccccagag ccaagaacac taagcactcg ccggcccttc 2280
ggcacccctc ccctccctcc cgactcaacc cggccgttgc ttctgtatat agagaaataa 2340
gttat 2345

```

<210> 36  
 <211> 709  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2427982CB1

```

<400> 36
gtgacagtag caaccgcccg aatggcgaaa gcaacaacaa tcaaagaagc cttagcgaga 60
tgggaagaga aaactggcca gaggccatct gaagccaaag agataaaact ttatgcccgag 120
attcccccta tagagaagat ggatgcatcc ttgtccatgc ttgctaattg cgagaagctt 180
tactgttcta caaactgcat tgaaaaaatt gccaacctga atggcttaaa aaacttgagg 240
atattatctt taggaagaaa caacataaag aacttaaatg gactggaggc agtaggggac 300
acattagaag aactgtggat ctctacaaat tttattgaga agttgaaagg gatccacata 360
atgaagaaat tgaagattct ctacatgtct aataacctgg taaagactg ggctgagttt 420
gtgaagctgg cagaactgcc atgcctcgaa gacctggtgt ttgtaggcaa tcccttgga 480
gagaaacatt ctgctgagaa taactggatt gaagaagcaa ccaagagagt gcccactg 540
aaaaagctgg atggtactcc agtaattaaa ggggatgagg aagaagacaa ctaatgccac 600
gctttccact gtgtgttaac ttattttaat gtcataagaa caatagataa attttatata 660
attgtctatt ttaaagattc tgtatgggac aaaagtttct taagataaa 709

```

<210> 37  
 <211> 1569  
 <212> DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2470833CB1

&lt;400&gt; 37

ctcgaccgct	cgagccgaat	tgggctcgag	attgagttta	agtggcctgc	agaaccccg	60
gggcccaggc	ggcacggcgg	gcgctggggc	tccgggcaga	gctttcagga	ggtttatggc	120
agcttcactt	tcacggcatc	cacccccctc	gggcccctgc	gcagagagga	ggaagctcct	180
gccggctgag	cgggcctgga	ggaagtgaac	agcggggctc	ctgcctcccg	gcctgggtcc	240
cgaagacccc	agaagaaccc	ggaacttgct	tccattcggg	atccagggac	caccctttgc	300
actcagtagg	cctttgtttt	cctgcgtgga	aagcggttgg	gcttgggagg	cgatggagcc	360
ggagttcttg	tacgacctgc	tgcagctccc	caagggggtg	gagccccag	cggaggagga	420
gctctcaaaa	ggaggaaaga	agaaatacct	gccacccact	tcccgggaag	accccaaatt	480
tgaagaactg	cagaaggtgt	tgatggagtg	gatcaatgcc	actcttctcc	ccgagcacat	540
tgtggtccgc	agcctggagg	aggacatggt	cgacgggctc	atcctacacc	acctattcca	600
gaggctgggc	gcgctcaagc	tggaaagcaga	ggacatcgcc	ctgacagcca	caagccagaa	660
gcacaagctc	acagtgggtg	tggaggccgt	gaaccggagt	ctgcagctgg	aggagtggca	720
ggccaagtgg	agcgtggaga	gcattctcaa	caaggacctg	ttgtctaccc	tgcacctcct	780
tgtggccctg	gccaaagcgt	tccagcccga	cctctccctc	ccaaccaacg	tccaggtgga	840
ggtcatcact	atcgagagca	ccaaaagtgg	tctgaagtca	gagaagtgtg	tggaacagct	900
cactgaatac	agcacagaca	aggacgagcc	tccaaaggac	gtctttgatg	aattatttaa	960
gctggctccg	gagaaagtga	acgcagtgaa	agaggccatc	gtgaactttg	tcaaccagaa	1020
gctggaccgc	ctgggcctgt	ctgtgcagaa	tctggacacc	cagtttgtag	atgggggtcat	1080
cttactcttg	ctgattggac	aacttgaagg	cttcttccctg	cacttaaagg	aattctacct	1140
cactcccaac	tctcctgcag	aaatgctgca	caacgtcacc	ctggcgctgg	agctgctgaa	1200
ggacgagggc	ctgctcagct	gccctgtcag	ccttgaagat	atcgtgaaca	aggatgccaa	1260
gagcacactg	cgggtgtctc	atggtctgtt	ctgcaagcac	acgcagaagg	cacacagggg	1320
caggacgccc	catggagccc	cgaattgacc	ctcactgcct	ccaaagccca	gagcctgcct	1380
gtcagcccag	ctggagggcc	cgaggctgca	gggtgtcctc	ccacagtccc	gctgtttcct	1440
gtgcattcgt	gaccgccttc	cctcccaccc	tgtctcctgt	ctccatcggt	ggattatctt	1500
tgaacccctt	tgtgtggatc	attttgagcc	gcctggcctt	gctcagttta	ttttaataaa	1560
agtattttct						1569

&lt;210&gt; 38

&lt;211&gt; 1172

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2080579CB1

&lt;400&gt; 38

cgacgggctg	cggcctgcgg	aacctgaggg	agctggggag	ggccggggcg	gccggccgga	60
tagcgagccg	cgctggcggc	ggcggtggcc	gcgatgatgg	agatccagat	ggacgagggc	120
ggcggcgctg	tgggtgtacca	ggacgactac	tgctccggct	cgggtgatgtc	ggagcgggtg	180
tggggcctgg	cgggctccat	ctaccgcgag	tccgagcgcc	tcatccactg	ctacgacgag	240
gaggtgggtca	aggagctcat	gccgctgggtg	gtgaacgtgc	tggagaacct	agactcgggtg	300
ctcagcgaga	accaggagca	cgagggtggag	ctggagctgc	tgcgcgagga	caacgagcag	360
ctgctcaccc	agtacgagcg	tgagaaggcg	ctgcgcaggg	aggcggagga	gaaattcatt	420
gagtttgaag	atgctctgga	acaagagaag	aaagagctgc	aaatccaggt	ggagcactac	480
gagttccaga	cgcgccagct	ggagctgaag	gccaagaact	atgccgatca	gatttcccg	540
ttggaggagc	gggagtcgga	gatgaagaag	gagtacaatg	ccctgcacca	gcggcacaca	600
gagatgatac	agacctacgt	ggagcacatt	gagaggtcca	agatgcagca	ggtcggagga	660
aacagccaga	ccgagagcag	cctgccgggg	cggaggtacg	cggggcgcg	cgggggtggag	720
gtacgcgggg	cgcggcgggg	cggaggtacg	caggacgcgg	cacatgccag	ggtcgtagt	780
cttgttatgg	cccgcgctct	gggctcagga	tgacagggaa	cactctggag	accagggagg	840
agggagctgg	ttggagcgtg	gctgagcagg	gatgtggggg	ggcggccctg	ggcgacggac	900
atgtgtcgag	ccagtgtgtc	ccttctgtct	gggtcccaat	tcagtcaggg	aggccccggg	960
cagcttgcgc	cttacttaacc	cttactatcc	acagattctg	tgtttgcaaa	tgatctact	1020
cgctaaaata	tatctgtgac	ctcagccggg	cacagtgact	tatgcctgta	atcctagaac	1080
tttgggaggg	caaggcagga	ggattacttg	agcccaggag	ttcgagctca	gcttggggca	1140
ccagcaagat	cttgtctcta	caaaaaaaaa	aa			1172

&lt;210&gt; 39

<211> 2380  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2156553CB1

<400> 39  
 caggggttttca ctgtcttagc caggatgggc tgcattctct gacctcatga tccatccacc 60  
 tcagcctccc aaagtgtctg gattacaggg gtcagccacc gtgaccagcc tgaacacagga 120  
 gcaagttctta aactcaggct ctagagtcag aaaaggtaga gtcagggttct ggatccaaaa 180  
 tggggcaagt catgatcagg ttctggaacc agaacaggcc tcaagcctag gggctgagca 240  
 ggggtatcccc tggcctggga gcagaggact tctggctgac tgctgcccgc aacgttctca 300  
 agctgggtggt gaagtctgag tggaaagtcac accctattca ggcagtagag gaagaggcct 360  
 caggagacaa gcagcccaag aaacaggaga aaaaccaggt gttgggtgtcc ccagagtttg 420  
 tggatgaagc tctgtgtgcg tgcgaggagt accttagcaa cttggcccac atggacatcg 480  
 acaaggacct ggaggccccg ctgtacctca cccccagggg ctgggtccctc ttctccaccg 540  
 gctactacca agtgggtccac gaagggggcga aactcaggca cctcgacact caggtccagc 600  
 gctgtgagga catcctgcag cagctgcagg ccgtggtacc ccagatagac atggaagggg 660  
 atcgcaacat ctggatcgtg aagccaggag ccaagtcctg tggacgaggg atcatgtgca 720  
 tggaccacct ggaggagatg ctgaagctgg tgaacggcaa ccccggtggg atgaaggacg 780  
 gcaagtgggt ggtgcagaag tatattgagc ggcctctcct catctttggc accaagtttg 840  
 acctcagaca gtgggttcctg gtaactgact ggaacccact taccgtgtgg ttctaccgcg 900  
 acagctatat cgccttttcc acgcagccct tctccctgaa gaacctggac aactcagtgc 960  
 acctgtgcaa caactccatc cagaagcacc tggagaactc atgccatcgg catccactgc 1020  
 ttccgccaga caacatgtgg tctagccaga ggttccaggc ccacctgcag gagatgggtg 1080  
 ccccaaatgc ttgggtccacc atcatcgtgc ctggcatgaa ggatgctgtg atccacgcac 1140  
 ttccagacct ccaggacacc gtgcagtgtc ggaaggccag ctttgagctc tatggcgctg 1200  
 acttcgtgtt cggggaggac ttccagccct ggctgattga gatcaacgcc agccccacga 1260  
 tggcaccctc cacagcagtc actgcccggc tctgtgctgg cgtgcaagct gacaccctgc 1320  
 gcgtgggtcat tgaccggatg ctggaccgca actgtgacac aggagccttt gagctcatct 1380  
 ataagcagcc tgctgtggag gtgcctcaat atgtgggcat cgggtcctg gttaggggct 1440  
 tcaccatcaa gaagcccatg gcgatgtgtc atcgccgat ggggggtccg ccagcagtc 1500  
 ctctgctgac ccagcgaggc tctggggaag gcaaggactc ggggatccct acccacaggt 1560  
 cagcttctag gaaaggcact ggggccaggga gcctggggca cagtgagaag ccagtctcca 1620  
 ctgccaccac ttccagcccc ggaaagggga agaaaggcaa ggcgaaaagg gccacagccc 1680  
 tggctgccc caatctctgg gactgggatg cccccagcac caggatgggc tgcattttca 1740  
 ccactgacct ttctagtggg gacaggcaac cccaccactt gaacagattg ccactgagtc 1800  
 cgaagaacct ccaggccctg ggtaagacca ttcccccaa acacccgagt gttccaaggc 1860  
 gatttattcc tgctctccag gccctccca accacctgga tcagccacct caccaaagag 1920  
 ccaccagtag caagtaaaag ccactactca caaagtattg tttaaaaata cacagccaaa 1980  
 ttactgggg acggtgtgtg gagcctgtgg tccaggctac tcgcgaggct aatgaggatc 2040  
 gcttgagccc gggaggtcaa ggctgcagtg agttatgac gcacctctgc actccagcct 2100  
 gggagacaga gcgagatgct gtcttaaaaa caaaaacaca aaaaaagcac tttgggaggc 2160  
 cgaggcgggg agatcacttg aggtcaggag ttcgagacca gcctggccaa catggtgaaa 2220  
 ccccatctct actaaaaaat acaaaaatta gccgagtgtg ggggcagggt cctataatcc 2280  
 cagctacttg gcacgactgag cgcttgacc caggaagcgg aggttgagtc 2340  
 gagccgagat tgcaccattg cactccagca tgggcagcat 2380

<210> 40  
 <211> 4396  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2182855CB1

<400> 40  
 tgccgctgca gggggctaga gatgcctcgt ggggttttcc agcagctttc caacttgggt 60  
 ttgcaggaaac ttaacgccaa cctgagcaac ctaacctcag cgtttgaaaa agcaacagct 120  
 gagaaaaatca agtgtcagca agaggccgat gccacgaaca gggatgatctt actggcgaa 180  
 aggctggctg ggggattagc atcggaaaac atccgctggg ctgagctctgt ggagaacttc 240  
 aggagccagg gggtcacgct gtgtggggac gtcctgctca tctctgcctt cgtgtccctac 300  
 gtgggctact tcaccaagaa ataccggaat gagctgatgg agaaaattctg gatcccttac 360  
 atacataact taaagggtccc catcccgatc acgaattggcc tggatccctt gaggctgctg 420

acagatgacg	cggacgtggc	cacctggaac	aaccagggcc	tccccagcga	ccgcatgtcc	480
accgagaatg	ccaccatcct	gggcaacacc	gagcgggtggc	cgctgatcgt	ggacgcccag	540
ctccaaggaa	tcaagtggat	caaaaacaaa	tacaggagtgt	aactgaaagc	catccgcctg	600
ggacagaaga	gtacacctga	tgatcatcgag	caggccatct	cggaagggga	cacgttgctc	660
attgagaaca	tcggcgaaac	cgtggacccc	gtgctggacc	ctctactggg	caggaacacg	720
attaaaaagg	gaaagtacat	taagatcggg	gacaaggagg	tggagtacca	ccccaaagtc	780
cgcttgatcc	tacacaccaa	gtacttcaac	ccacactaca	agccagagat	gcaggctcag	840
tgcaccctca	tcaacttcct	ggtcaccagg	gatggactcg	aggaccaact	cttggccgct	900
gtggtggcca	aagagcgccc	agatctggaa	cagctgaagg	caaacctcac	caagtcccaa	960
aacgaattta	agattgttct	gaaagagctg	gaagactcgc	tcctggcccg	tctgtcggct	1020
gcgtcgggga	actttctggg	agacacggcc	ttggtggaga	atctggagac	caccaagcac	1080
acagccagcg	agatcgagga	gaagggtggg	gaggcaaaaa	tcacagaagt	taaaatcaac	1140
gaagcgagag	agaactaccg	cccggctgcg	gagagggcat	ctctgctcta	cttcatactg	1200
aacgatctca	acaaaatcaa	cccgtctac	cagttctccc	tcaaggcctt	caactgtggtg	1260
tttgagaaag	ccatccagag	gaccacccct	gccaacgagg	tgaagcagcg	ggtgatcaac	1320
ctgacgggacg	agatcaccta	ctccgtctac	atgtacacgg	cccggggact	cttcgagagg	1380
gacaaactca	ttttcctggc	acaagttacg	tttcagggtcc	tgatccatgaa	gaaggagctg	1440
aaccagtggt	agctggattt	cctcctggcg	ttccctttta	aggccggagt	ggtctcacca	1500
gtggacttcc	tcagcatca	aggctggggc	gggatacaagg	ccctctcgga	gatggatgag	1560
ttcaaaaatc	tggacagtga	catcgaagga	tctgccaaagc	gctggaaaaa	gctgggtggag	1620
tcggaagccc	ccgagaagga	gatcttcccc	aaggagtggg	agaacaagac	ggccctgcag	1680
aagctgtgca	tgggtgcgctg	cctgcggcca	gatcgcatga	cctacgctat	caagaacttc	1740
ttggaggaaa	agatgggcag	caagttctgtg	gaaggccgga	gtgttgagtt	ttctaagttc	1800
tacgaggaga	gcagcccttc	caogtcaatc	ttctcatccc	tctccccggg	ggttgacccc	1860
ttgaaagacg	tggaaagccct	gggaaaaaaa	ctagggttta	ccatagacaa	tggaaaactc	1920
cataatgtgt	ccctggggca	gggacaagag	gtgggtggctg	agaacgccc	ggacgtggct	1980
gcagagaaaag	gacactgggt	cattctgcag	aatatccacc	tgggtggccc	gtggctggga	2040
acactggaca	agaagctgga	gcgtacacgc	acgggcagcc	atgaggacta	ccgggtgttc	2100
atcagcgcg	agcctgcccc	cagtcccgag	accacatca	ccccccagg	cattctggag	2160
aacgccatca	agatcaccaa	cgagccccc	acgggcatgt	acgccaaact	gcacaaggcc	2220
ctggacctgt	tcaccagga	caccctggag	atgtgcacca	aggagatgga	gttcaagtgc	2280
atgctctctg	ccctgtgcta	cttccacgct	gtgggtggcag	agaggcgcaa	gttcggcgcc	2340
cagggctgga	accggtcgta	ccccttcaac	aacggggacc	tcaccatctc	catcaacgtg	2400
ctctacaact	acctggaggc	caaccccaag	gtgccttggg	acgatctccg	ctaccttttt	2460
ggtgaaatca	tgtatggcgg	ccacatcaca	gatgactggg	accgtcggct	gtgcaggacc	2520
tacctggctg	aatacatccg	gacggagatg	ctggaggggag	acgtcctgct	ggcccccg	2580
tttcagatcg	ccccacact	ggactacaag	ggttacacg	aatacatcga	tgagaacctg	2640
ccccctgaga	gtccctatct	gtatggcctg	caccccacg	cagagattgg	ctttctgacg	2700
gtcacctcag	agaagctggt	cgcactgtc	ctggaaatgc	agccaaaaga	gacggactcg	2760
ggggcaggca	cgggagtgtc	ccgcgaggag	aagggtgaagg	ccgtgctgga	cgacatcctg	2820
gagaagattc	cggagacttt	caacatggct	gagatcatgg	caaaggcagc	ggaaaagacc	2880
cctcagtggt	tagtcgcctt	tcaagaatga	gaaagaatga	acatcctgac	caacgaaatg	2940
cgccgttcgc	tcaaggagct	gaacctgggt	ctgaaggagg	aactgaccat	cacgaccgac	3000
gtggaagatc	tgtccacggc	tctcttctat	gacaccgtgc	ctgatacgtg	ggtggcccgg	3060
gcctacccct	ccatgatggg	cctggcggcc	tggtagcgag	acctgctgct	ccgcatcagg	3120
gaaactcgagg	ctgggacgac	agaactttgc	ctgcccacca	ccgtgtggtt	ggccggcttc	3180
ttcaaccccc	agtcgttctt	cacggcctcc	atgcagtcga	tggccaggga	gaacgagtgg	3240
ccccctggaca	agatgtgtct	gtctgtcgag	gtgaccaaga	aaaaccgaga	ggacatgacc	3300
gtccccccac	gagagggtct	ctacgtgtac	ggactcttca	tggaaagggc	tcgctgggac	3360
accagactgt	gagtcatcgc	tgaagcgcg	ctgaaagagc	tgaccccgcc	catgcctgtc	3420
atcttcatca	aggccatttc	tgtggaccgc	atggagacca	agaacatcta	tgagtgtccc	3480
gtgtacaaaa	cacgcatccg	cgccccacc	tatgtctgga	cctttaactt	gaagacaaa	3540
gagaaggcag	cgaagtggat	cctggcagcc	gtggcgctgc	tcctacaggt	ttagctcgct	3600
cctgcctcac	agcccacact	ccctggggct	ggaccacaac	tcagcccttc	acctgtgcac	3660
ctgtgactta	ttctttacag	gaactgggtg	tggtttttcg	ttctcttaaa	taatcaggtg	3720
ctttgtaacc	aagcacatcg	gaaccagagg	gtggaggttg	gtgtggaaga	ggtggggcag	3780
attaaagcca	gtggagccac	tcagctgtgc	ccatccattc	tgtgcctgat	ggccactgtg	3840
aggcctgggt	caggcttttg	ggaaaggccc	caattcccag	cagccagagg	caagcatttc	3900
aggaagtaaa	tcccaaatcc	tgacttcccc	ggggggtcag	tggagtgttc	ttttaacagg	3960
agcagcatgt	ggggtgaggg	cttggcactc	acaccatctg	gcctgatgct	tcatgcaaac	4020
cacttgccaa	tgaggggaca	gggcgttacc	ccataccagg	agctgcgcgc	tccacacagc	4080
gtctcatccg	ctgctcccaa	ctgctcccaa	aaacaacagg	accttttaca	agtaaggaaa	4140
cacacccaag	ccccgccttc	cagagtctag	aagaacaagc	cgttcccaca	agcaagcccc	4200
ggacctgtag	gactctgtgg	ctgggctgat	gcaggacacg	gcagtgaagt	ggccaggcta	4260
gctagggcag	caggggtgcag	gcacgagggg	agagaattaa	ctagcaacct	gggggatgag	4320
gggagacagc	ggcaaacgtc	agaaaagacg	gatcccaggg	ggaccaagac	aggatcaaca	4380
tttttttttt	tttttcg					4396



<210> 41  
 <211> 1831  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2242106CB1

<400> 41  
 gtgagctgcc tgctgtgtcc gcctcctctc ctctcaggac tcagcttccc gccctgcaga 60  
 gggcacaagg gagctttcca tggggcccttt gggcccgaga caggaaggcg ggccctgacc 120  
 ctctcagctc ggagcccagg tgaaggtctg cagtgaagccc tgaggggtgg aactcggggg 180  
 aggagtcttg aagggggcca gggctggggc caggactggg ggctgtggag aggcagaaag 240  
 aggcaggtat ccttggaag cgcagctgg tagggggcag gcaacgctgg agggagctgc 300  
 caggtgcagg gcaggggtgt ccggcaaac tccctggggg acaacagggg agctgggggg 360  
 taggactctc cctctgtggg gatgacttcc agccaagtgg ctctgccgcc tggccctggc 420  
 cctggccctc taggtgggct ggggcgggtg tttggccagg aagtcaggct gggggtgggg 480  
 ctgagcctct gccagggtcc actgtccctg caggagggt gtacagtggg gaggtgttgg 540  
 tgtgggacct gagccgtctt gaggaccgc tgctgtggcg cacaggcctg acggatgaca 600  
 cccacacaga ccctgtgtcc caggtcaggg caggggctgc cgggcggggg gggccctgag 660  
 gcagagctgg ggagagggcg cccactctcc taggaaggct acccgggcgt tggcagttgg 720  
 acagctgccc cctgcattct cggagctccc gcagcatgag tcacccccct caggagccac 780  
 tcccagggcc ctgtggccat tcccagcctc ctctgcaggc accgagtgg agcgctggg 840  
 attctggggc tggagagggc cctgggaacc ctccaggcct gcccctggg ggggtgagcct 900  
 gggtctgggg cctcccgagg aatttttttt tttcctggaa gagagggagg gtaggggtgg 960  
 agcgtgacac ctgggcagggt gtccctgtgc tccatcctgg ccttgcattg tgtaaactca 1020  
 ggtggtgtgg ctgcccgagc ctgggcacag ccaccgcttc caggtgctga gtgtggccac 1080  
 tgacgggaag gtgctactct ggcagggcat cggggtaggc cagctgcagc tcacagaggg 1140  
 ctctgccctg gtcatgcagc agctgccacg gagcaccaag ctcaagaagc atcccccgcg 1200  
 ggagaccgag gtgggcgcca cggcagtggt cttctccagc tttgacccta ggctgttcat 1260  
 tctgggcacg gaaggcggct tcccgcctca gtgttccctg gcagctggag aggcagccct 1320  
 cactcggtat cccagctccg tgcccctgcy ggcccagca cagtttacct tctcccccca 1380  
 cggcggtccc atctactctg tgagctgttc ccccttccac aggaatctct tctgagcgc 1440  
 tgggactgac gggcatgtcc acctgtactc catgtgcag gccctccct tgacttgcct 1500  
 gcagctctcc ctcaagtatc tgtttgtgtg gcgctgggtc ccagtgccgg ccttgggttt 1560  
 tggcagctgc ctgggaaag gtgacgtgca gctgtttgat ctccagaaaa gctccagaa 1620  
 accacagatt ttgatcaagc aaaccagga tgaagccct gtctactgtc tggagtcaaa 1680  
 cagccagcag actcagctct tggctgcggg cgatgcccg ggcacagtga aggtgtggca 1740  
 gctgagcaca gagttcacgg aacaagggcc cggggaagct gaggacctgg actgcctggc 1800  
 agcagaggtg gcggcctgag ggggtcccgga a 1831

<210> 42  
 <211> 3249  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2726877CB1

<400> 42  
 tttttatcga actcttatca ccttgttggc ccatagtagt ttaactgtgg ttgtgtttgc 60  
 actttcaata ttatccagtt tgacattaaa tgaagaggtg ggggaaaagc tattccatgc 120  
 tcgaaacatt catcagactt ttcaactaat atttaatat ctcataaacg gtgatggcac 180  
 tctaactaga aagtattcag ttgacctact gatggatctc cttagaatc ctaaaattgc 240  
 tgattatctc accagatatg agcacttttc ttcatgtctt caccaagtat taggtcttct 300  
 taatggaaag gatcctgatt cctcttcaaa ggttttagaa ttacttcttg ccttctgttc 360  
 agtgactcag ctgcgccata tgctcactca gatgatgttt gaacagtctc cacctggcag 420  
 gcgccactct ggaagccata ctaaatgttt agaacctact gtggctctac tgcgctgggt 480  
 aagccaacct ttggacggat cagaaaactg ttctgtttta gcattggagt tgttcaagga 540  
 aatattttgag gatgtcatag atgtgtctaa ctgttccctg gctgatcggt ttgtgacct 600  
 tctgtgcct acaatccttg atcaacttca gttcacagaa caaaatctag atgaggcttt 660  
 aacaagacaa aaatgtgaaa ggattgccaa ggcctttgaa gttttgttaa ctctctgtgg 720  
 agatgatata ctaaaaatgc atattgcaaa aatcttgaca actgtcaagt gtaccactct 780  
 tatagaacaa caatttacat atggcaagat tgacctggga tttggaacaa aggttgacaa 840  
 ttctgaatta tgcaaacttg ctgctgatgt aattttgaaa actcttgatt tgattaacaa 900

acttaaacca	ttggttcctg	gtatggaagt	aagcttctac	aaaatacttc	aggacccacg	960
tttgattact	ccttttgctt	ttgctttaac	gtcagataat	agagaacaag	tacagtctgg	1020
actgagaata	ttattggagg	ctgctccact	gccagatfff	cctgcttttag	tactttggaga	1080
aagtatagca	gcaaacaatg	cctatagaca	acaggaaca	gaacatatat	ccagaaaaat	1140
gccctggcaa	tcatcaaate	acagttttcc	aacatcaata	aagtgtttta	ctcctcattt	1200
gaaagatggt	gttcctggat	tgaatattga	agaattaata	gagaaacttc	agtctggaat	1260
ggtggtaaa	gatcagatff	gtgatgtgag	aatatctgac	ataatggatg	tatatgaaat	1320
gaaactatcc	acattagctt	ccaaagaaag	caggctacaa	gatctttttg	aaacaaaagc	1380
tctagccctt	gcacaggctg	atagactgat	tgctcagcat	cgctgtcaaa	gaactcaagc	1440
tgaacacag	gcacggacac	ttgctagtat	gttgagagaa	gttgagagaa	aaaatgaaga	1500
gcttagtgtg	ttgctgaagg	cgcagcaagt	tgaatcagaa	agagcgcaga	gtgatattga	1560
gcatctcttt	caacataata	ggaagttaga	gtctgtggct	gaagaacatg	aaatactgac	1620
aaaaatctac	atggaacttc	ttcagagaaa	tgaagtact	gaaaagaaga	ataaagattt	1680
acagatcaca	tgtgattctc	tgaataaaca	aattgagaca	gtgaaaaagt	tgaatgagtc	1740
actcaaggaa	caaatgaaa	aaagtattgc	ccaattaata	gagaaagaag	aacagagaaa	1800
agaagtacag	aatcagctag	tagacagaga	acataagcta	gcaaatttgc	atcaaaaaac	1860
aaaagtacaa	gaagaaaaga	ttaaaaccct	acaaaaggaa	aggggaagata	aggaagaaac	1920
cattgatata	cttagaaaag	aattaagcag	aacagaacag	ataagaaaag	agttgagcat	1980
taagccttcc	tccctagagg	ttcaaaaggc	acaattagaa	ggctcgtttg	aagagaaaaga	2040
gtccttggtg	aaacttcagc	aagaggaatt	gaacaaacac	tcccacatga	tagcaatgat	2100
ccacagttta	agtggtggaa	aaataaatcc	agaaactgtg	aatctcagta	tatagacatt	2160
atggcatttt	ggaatttgta	atctcatgat	atttttgatg	tatttatcta	ttggaggggg	2220
gggtggtagt	ggagtttaatt	tgtgacttcg	taacaataag	aagttattat	ctaatttagt	2280
aaagaccctg	atctgttgca	tgttttttat	ttaatagttt	gaatagaaat	ttaattttct	2340
aagttttact	ttttgtttct	ggcttttatg	gcttaagggt	ttctttgggt	cttacattag	2400
aaaatcattt	ttaacctcca	ttatcatttt	tctaagggtc	tttctttttt	cttagttgct	2460
ttctattctg	ttttgcctgt	cttattttat	ctcattttgt	attatttaga	tcttaagacc	2520
aaactttctt	ggtataacag	tctataacag	tacaaaataa	aaatatagag	agagttaaag	2580
taaaaagtaa	agtaaaaaga	gagaaggat	gattttactg	ctagagaagt	ttgtctctga	2640
agaagcacaa	aagaaaatat	tagtgaattt	aaaataattt	ttatactgct	gtagcataat	2700
ttctaaattt	gaaaaaaatg	caatggtaat	aaaatgtata	aaaattagaa	aactgtcatt	2760
gtgttaaact	attacattta	aatgattaca	tttaacacaa	tagctgtctc	ataaaaaatc	2820
taagaacttg	tagaatttatt	tgtaaaggtat	attagtgttt	ttttctcttt	tttcaaatgt	2880
tgcatagtgg	tccagatcat	ctataattaa	ttataagttt	cttatgattt	acataaggat	2940
aattttggatt	ctactcttgc	tgcataattca	gaaaaatat	atttactgtt	aattctgata	3000
tgaattgtat	ttaatattgt	gtcctaata	tttttaattg	gttcatttaa	ttttaagca	3060
taggttaaat	atctttattt	tttttaaaat	acaagggctc	ttctgtactt	ttcatttgaa	3120
tcatatcctc	taactactct	gtaagaaacg	tttttgcata	aatattttga	ttgatattat	3180
tgcataacca	atacagtcatt	taatttttact	tggtgttttc	ttcacacata	tgtctgtttg	3240
atttaagtgt						3249

<210> 43  
 <211> 4133  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2738233CB1

<400> 43						
gaaaaatgtg	agaaagggaat	agcagattcc	ctggagaaac	tacgaacttt	caaaaagaag	60
ctttcgcagt	ctctcccgga	tcaccatgaa	gagctccatg	cagaacaaat	gcgttgcaag	120
gaattagaaa	atgcagttgg	gagctggaca	gatgacttga	cccagttgag	cctgctgaag	180
gacaccctct	ctgcctatat	cagtgtctgat	gatattctcca	ttcttaatat	acgcgtagag	240
cttctgcaaa	ggcagtggga	agaactatgc	caccagctct	ccttaaggcg	gcagcaata	300
ggtgaaagat	tgaatgaatg	ggcagtcctc	agtgaagaaga	acaagggaact	ctgtgagtg	360
ttgactcaaa	tggaaagcaa	agtttctcag	aatggagaca	ttctcattga	agaaatgata	420
gagaagctca	agaaggatta	tcaagaggaa	attgctattg	ctcaagagaa	caaaatacag	480
ctccaacaaa	tgggagaacg	acttgctaaa	gccagccatg	aaagcaaagc	atctgagatt	540
gaatacaagc	tgggaaagggt	caacgaccgg	tggcagcatc	tcctggacct	cattgcagcc	600
aggggtgaaga	agctgaagga	gacctgggta	gccgtgcagc	agcttgataa	gaacatgagc	660
agcctgagga	cctggctcgc	tcacatcgag	tcagagctgg	ccaagccaat	agtctacggg	720
tcctgtaaact	cggaagaaat	acagagaaag	cttaatgagc	agcaggagct	tcagagagac	780
atagagaagc	acagtacagg	tgttgcattc	gtcctcaacc	tgtgtgaagt	cctgctgcac	840
gactgtgacg	cctgtgccac	tgatgccgag	tgtgactcta	tacagcaggc	tacagaaaac	900
ctggaccggc	ggtggagaaa	catttctgct	atgtccatgg	aaaggaggct	gaaaatcgaa	960

gagacgtggc	gattgtggca	gaaatttctg	gatgactatt	cacgttttga	agattggctg	1020
aagtcttcag	aaaggacagc	tgcttttccc	agctcttctg	gggtgatcta	tacagttagcc	1080
aaggaagaac	taaagaaatt	tgaggctttc	cagcgacagg	tccacgagtg	cctgacgcag	1140
ctggaaactga	tcaacaagca	gtaccgcgcg	ctggccaggg	agaaccgcac	tgattcagca	1200
tgtagcctca	aacagatggt	tcacgaaggc	aaccagagat	gggacaacct	gcaaaaagcgt	1260
gtcacctcca	tcttgccgag	actcaagcat	tttattggcc	agcgtgagga	gtttgagact	1320
gcgcggggaca	gcattctggt	ctggctcaca	gagatggatc	tgacgtcac	taatatggaa	1380
catttttctg	agtgtgatgt	tcaagctaaa	ataaagcaac	tcaaggcctt	ccagcaggaa	1440
atttcactga	accacaataa	gattgagcag	ataattgccc	aaggagaaca	gctgatagaa	1500
aagagtgagc	ccttggtatgc	agcgatcatc	gaggaggaaac	tagatgagct	ccgacgggtac	1560
tgccaggagg	tcttcggggcg	tgtggaaaga	taccataaga	aactgatccg	cctgcctctc	1620
ccagacgatg	agcacgacct	ctcagacagg	gagctggagc	tggaagactc	tgcagctctg	1680
tcggacctgc	actggcacga	ccgctctgca	gacagcctgc	tttctccaca	gccttctctc	1740
aatctctccc	tctcgctcgc	tcagccccctc	cggagcgagc	ggtcaggagc	agacaccccc	1800
gctagtgtgg	actccatccc	cctggagtgg	gatcacgact	atgacctcag	tcgggacctg	1860
gagtcctgcaa	tgtccagagc	tctgccccctc	gaggatgaag	aagggtcagga	tgacaaagat	1920
ttctacctcc	ggggagctgt	tggtttatca	ggggaccaca	gtgccctaga	gtcacagatc	1980
cgacaactgg	gcaaagccct	ggatgatagc	cgctttcaga	tacagcaaac	cgaaaatatc	2040
attcgacagca	aaactcccac	ggggccggag	ctagacacca	gctacaaagg	ctacatgaaa	2100
ctgctggggcg	aatgcagtag	cagtatagac	tccgtgaaga	gactggagca	caaactgaag	2160
gaggaagagg	agagccttcc	tggttttggt	aacctgcata	gtaccgaaac	ccaaacggct	2220
ggtgtgattg	accgatggga	gcttctccag	gcccaggcat	tgagcaagga	gttgaggatg	2280
aagcagaacc	tccagaagtg	gcagcagttt	aactcagact	tgaacagcat	ctgggcctgg	2340
ctgggggaca	cggaggagga	gttgggaacag	ctccagcgtc	tggaactcag	cactgacatc	2400
cagaccatcg	agctccagat	caaaaagctc	aaggagctcc	agaaagctgt	ggaccaccgc	2460
aaagccatca	tcctctccat	caatctctgc	agccctgagt	tcacccaggc	tgacagcaag	2520
gagagccggg	acctgcagga	tcgcttgctg	cagatgaatg	ggcgctggga	ccgagtgtgc	2580
tctctgctgg	aggagtggcg	gggcctgctg	caggatgcc	tgatgcagtg	ccagggtttc	2640
catgaaatga	gccatgggtt	gcttcttatg	ctggagaaca	ttgacagaag	gaaaaatgaa	2700
attgtcccta	ttgattctaa	ccttgatgca	gagatacttc	aggaccatca	caaacagctt	2760
atgcaataaa	agcatgagct	gttgggaatcc	caactcagag	tagcctcttt	gcaagacatg	2820
tcttgccaac	tactggtgaa	tgtgaagga	acagactggt	tagaagccaa	agaaaaagtc	2880
catgttattg	gaaactcggc	caaacttctc	ttgaaggagg	tcagtcgtca	tatcaaggaa	2940
ctggagaagt	tatttagacgt	gtcaagtagt	cagcaggatt	tgtcttctctg	gtcttctgct	3000
gatgaactgg	acacctcagg	gtctgtgagt	cccacatcag	gaaggagcac	cccaaacaga	3060
cagaaaaacgc	cacgaggcaa	gtgtagtctc	tcacagcctg	gacctctgt	cagcagtcca	3120
catagcaggt	ccacaaaagg	tggtctccgat	tcctcccttt	ctgagccagg	gccagggtcgg	3180
tcgggcgcgcg	gcttctgttt	cagagtctctc	cgagcagctc	ttccccctca	gcttctcctg	3240
ctctctctca	tcgggcttgc	ctgccttgta	ccaatgtcag	aggaagacta	cagctgtgcc	3300
ctctccaaca	actttgcccg	gtcattccac	cccattgctca	gatacacgaa	tggtccctcct	3360
ccactctgaa	ctaagcagat	gccatctgca	gaagtgtctg	tagcataagg	aggatcgggt	3420
caataagcaat	cccaaaactac	caacaagagg	accttgatct	tggtcgaaagc	cctcggtgtg	3480
gcagcttttag	ccctcctcca	gatcacatgt	gtgcaaatta	tggttccaga	ggtggaagat	3540
aaacagtgc	aggggaacaa	acagacaaca	agaaggtttg	gaagaaatct	ggtttgagac	3600
tctgaacctt	agcactaagg	agattgagta	aggacctcca	aagttccccg	gactcatgaa	3660
ttctggggccc	ttggcccatt	ctgtgcacag	ccaaggactt	cagtagacca	cttggggcagc	3720
tttcccatgg	tgctgctcca	accatcagat	aaatgacct	cccaagcacc	atgtcagtg	3780
cgtacaatat	accaaccaac	cagtgtctgaa	gagatttttag	aaccttgtaa	catacaattt	3840
ttaagagctt	atatggcagc	ttccttttta	ccttggtttc	ctttggggca	tgatgtttta	3900
acctttgctt	tagaagcaca	agctgtaaat	ctaaaaggca	ctttttttta	gaggtataaa	3960
gaaaaactag	atgtaataaa	taagatcatg	gaaggcttta	tgtgaaaaaa	gttgaatgtt	4020
atagtaaaaa	aaaaaaagat	atttatgtat	gtacagtttg	ctaaagccaa	gttttgtttg	4080
tattgatttc	tttgcattta	ttatagatat	tataaaataa	aaaaaaaaaa	aaa	4133

&lt;210&gt; 44

&lt;211&gt; 1754

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1833116CB1

&lt;400&gt; 44

cggaagggga	aaacaactac	ggctgcgggtg	tggttggtgg	tgagatgacg	accttagtgc	60
tggataatgg	agcttacaac	gocaaaatcg	gttacagcca	tgaaaatgtg	tcggttatcc	120
ctaattgtca	gttccgggtca	aaaacagcac	gtcttaaaac	ttttactgcc	aaccagatag	180

atgaaataaaa	agacccttct	ggactctttt	acatcctccc	ttttcaaaa	ggctacttgg	240
tgaattggga	tggtcagaga	caagtttggg	attacctttt	tggaaaagaa	atgtatcagg	300
ttgatttttt	agatacta	attattatca	ctgaaccata	ctttaacttc	acttcaattc	360
aagaatcaat	gaatgaaatt	ctatgtgaag	aataccagtt	tcaagcagta	ttaagagtaa	420
atgctggggc	tctcagtgca	cataggtatt	tccgagataa	tccttccgaa	ttatgctgta	480
tcattgttga	tagtggatat	tcctttacac	atatagttcc	ttattgtaga	agtaaaaaga	540
aaaaagaagc	aattattcgg	ataaatgtgg	gaggaaaact	cttaaccaat	catctaaagg	600
agatcatatc	ttacaggcag	ctacatgtta	tggatgaaac	acatgtgatt	aatcaagtga	660
aagaagatgt	atgctatgtg	tctcaggatt	tttatagaga	catggatatt	gcaaagttga	720
aaggagaaga	aaatacagta	atgatagact	atgtcttgcc	tgacttcagt	acaattaaaa	780
agggcttttg	taagccaagg	gaagagatgg	tgttgagtgg	aaaatacaaa	tctggggaac	840
aaattcttcg	tttggccaat	gagagatttg	ctgttccgga	aatactcttt	aatccttctg	900
atataggcat	tcaagaaatg	ggaattccag	aagctattgt	ctattcaatt	caaaatctac	960
ctgaagaaat	gcagccgcat	ttttttaaga	acattgtctt	gacaggagga	aattcccttt	1020
tcccaggatt	tagggatcgg	gtttactcag	aagttcgatg	tcttactcca	acagattatg	1080
atgtttctgt	tgtgctgcct	gaaaacccta	ttacttatgc	ctgggaagg	ggaaaattga	1140
tatcagagaa	tgatgatatt	gaagatatgg	tggtacaacg	agaagattac	gaagaaaatg	1200
gacatagcgt	ctgtgaagag	aaatttgata	tttaagcaac	atttttgaat	gaaagtttgt	1260
accataaggt	ttaaatttcaa	agttcctttt	aaaagaggtt	aaggaaactgt	gttacctttt	1320
gtcctaagaa	aaaggcttga	atztatgtaa	atactttgat	cgattgctaa	ttttcaaagg	1380
cttcttaggt	aggttactac	agtaaaactgt	aactcagtc	acattttcat	ttaggagcta	1440
gactaccata	acaatgccta	tgtctgttcc	aagggttaggt	tatttttcat	taaaagaaga	1500
atgaatgcct	tttaagttta	attccttcata	gctgaaagca	caaatttaac	ggcttcactg	1560
gacagttttc	cttagaagg	agttttgtgt	gactgtgact	aaactatatt	attttaaaat	1620
gtcattctta	tttatacatt	ctaaagttgg	aaagactgat	cttatatgtg	tataatgttt	1680
attttgtacc	tagagtacat	ttaaaagggt	ggagactaag	ctaataaagt	ttttttggcc	1740
actaaaaaaa	aaaa					1754

&lt;210&gt; 45

&lt;211&gt; 2713

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 001799CB1

&lt;400&gt; 45

ttgaatggcc	ctgagtggga	ctcggcccag	aagccgaggg	actctctagg	ctgccggggcg	60
ctgggtcgtca	gcgcccaggc	tgggctgagg	cgccgcggta	ccatgaggcg	ccgggtactta	120
agagattatg	gcatacagaa	cccacaatgt	taaaaaacgg	aactttttgta	ataagattga	180
ggatcatttt	attgatcttc	ctagaaaaaa	gatctcta	ttcactaata	agaacatgaa	240
ggagggttaag	aaatctccaa	aacagttggc	tgcttacata	aatagaacag	ttggacaaac	300
tgtgaaaagc	ccagataaac	ttcgtaaaag	gatctatcgc	agaaagaaaag	ttcatcatcc	360
ctttccaaat	ccttgttaca	gaaaaaaaaca	gtcccctgga	agtggggggct	gtgacatggc	420
aaataaagaa	aatgaactgg	cttgtgcagg	ccacctgcct	gaaaaattac	accatgatag	480
tcgaacatat	ttggttaact	ccagtgatct	tggttcttca	cagacagaaa	gcccatcatc	540
aaaatatagt	gggttttttt	ctgaggtttc	tcaggaccat	gaaacaatgg	cccaagtttt	600
gttcagcagg	aatatgagat	tgaatgtagc	tttaactttc	tggagaaaaga	gaagtataag	660
tgaacttgta	gcttattttg	tgaggataga	agatccttgg	gttggtggtag	attgccttcc	720
tgtgctcacc	aattgtttac	aggaagaaaa	acaatatatc	tcacttggct	gctgtgttga	780
cttggtgcct	ctagtaaagt	cactacttaa	aagcaaat	gaagaatatg	ttatagttgg	840
tttaaaactgg	cttcaagcag	tcattaaaag	gtgggtggta	gaactatcat	ccaaaacaga	900
aattataaat	gatggaaata	ttcaaat	aaaacaacaa	ttaagtggat	tatgggaaca	960
ggaaaaccat	cttacttttg	ttccaggata	tactggta	atagctaagg	atgtagatgc	1020
ttatttatta	cagttacatt	gagagatttc	atctactaaa	gagcattttg	tttttcaaaa	1080
catccctgaa	ctgtataatt	tacaaaaaaa	aaagtctcgt	ctgagaactg	tgaactgtgg	1140
aagaaatcaa	aactat	tcttttaaaa	agccacgtaa	tgaaccact	aatgaaatcc	1200
cagcaatctg	cttcacattg	aagtggaaaa	atatccaaaa	ggagcagctt	caatttcatt	1260
gagggtgaaag	tgactatga	agattgttca	cctttgctgc	atgtgggagt	tatatgtgta	1320
tttggttaaca	ttaaagaacta	ctggatttta	atgcaatcct	gcataaaaaa	ataatttata	1380
ctatgtgaaa	aaataagaca	ggacttacca	ctaggaacca	ccaagaccaa	tcatatttaa	1440
cttttttaag	attgtgtttt	attaaaaaaa	aaaaacactt	aaatgtgtgc	agctat	1500
ttatgttgaa	aagactgaaa	gttttaaaaca	tgaaaaaaat	caatatataa	cattttttgt	1560
tcacactgag	atactgtgta	tgtaaaaatgc	cttaattatt	aataagccaa	tgtgttatga	1620
taccaaatatc	tgtttttaaaa	aactaaaacc	aacctgctt	ctggcatgat	aaaatcatgg	1680
aattaaatca	ggggtttaca	ttcttgtaga	gtgttcttga	aacactctct	gcaccatttt	1740

taaaacttga	gaatagtttt	agtatctctg	atattttttg	ccagaatcat	catgtcatgt	1800
atgaatgtgt	tatccctatc	taaggaaaaa	ggtgaatatg	tttttgtatg	aatgttttaac	1860
tggaaatgtc	catggacttg	gctaattttat	atttactttt	tattgtacat	agattttctaa	1920
tattttttcat	tccctgtatca	tttaaacttc	cttcatttga	gtaaattcac	taaatatttc	1980
tatttttttgc	ttttttaaat	tctgattttta	tatgaattct	aattcttttt	cactacatat	2040
gttttaaaga	gttacatata	gtgattttaga	atgggtttaca	gttaatgctg	atcttgtatt	2100
ttaaattcca	acacttttgt	tcactacctc	ctctaattgg	tagtatgata	tgctagcaga	2160
ctgtatgagg	tcttttttta	aaataccact	tttagtgtca	gtgaacccaa	ttctggaatg	2220
tcttaacagc	tctaaatctt	acttgtcttg	aaaatgattg	gggtttaata	ccactgctgg	2280
tggttcacac	atcatcccat	ccttaatatg	cctgacaggc	atctgagcaa	agggtttttag	2340
taattgaatt	tctctgcagt	agtccttcaa	gcacttgaat	gtaaaccttt	agcattttatt	2400
cgtttaatat	ctactgatac	gaatctcaag	cagattttct	gctcttaaaa	gttatgtttc	2460
actgagttct	ggtttttgtgt	agctatatatt	tatatagcta	gatatctctc	acagtgaaca	2520
tgaattgttaa	taattgggtta	tttccttaag	tcttttagatt	ataataattt	cagattattg	2580
cacgtctgtg	atttgagagg	tgagttattt	aagaggccag	ttttcaggac	atgggaattt	2640
gaattgtaaa	cctgttatct	ctgtgaaact	tttaacatga	taaaatataa	cctttctttg	2700
tgtttaaaaa	aaa					2713

&lt;210&gt; 46

&lt;211&gt; 1768

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 119814CB1

&lt;400&gt; 46

tttttttttt	tttttttttg	ctcttttagaa	gaggttatat	ttttattatc	cttatttttgg	60
agaacttttt	cttataaaat	ttttttttcca	gattcccttat	gaactcaagt	tagtggttaa	120
gcttttgatt	ccactgttaa	cagttttatgt	aaaaacactt	aacaaattgc	cattttatatg	180
ccaaactata	gctcaagaac	actctgtttt	agaaaaatta	cgcattagat	caggaagcct	240
catatatatg	tgcctctggg	acttcatttg	cagtcacatt	tagccagaaa	agcaatgact	300
tctatatattc	ttatggaaac	caatgtaaca	taaattaatg	ttctaaatat	agaaattaag	360
agttcataaaa	gagactgagg	ttgcatgtaa	aagagttatg	gtttgagaca	gtctaaaaaat	420
actatgttaa	tttcaaggat	cttattttcca	atgtttttgt	taaaaaatta	taataacttt	480
tgagctcttg	ctttgcattt	caatcgcaaa	cccactcaga	tacgggaact	gtttaaattc	540
atatattggac	aaataggttt	cagtgatgca	atactttaaa	attctgccat	ctccttgtgt	600
ttttctttct	agggtgagtgg	actgccagct	cctgatgtgt	catggtatct	aaatggaaga	660
acagttcaat	cagatgattt	gcacaaaatg	atagtgctctg	agaagggtct	tcattcactc	720
atctttgaag	tagtcagagc	ttcagatgca	ggggccttatg	catgtgttgc	caagaataga	780
gcaggagaag	ccaccttcac	tgtgcagctg	gatgtccttg	caaaaagaaca	taaaagagca	840
ccaatgttta	tctacaaacc	acagagcaaa	aaagttttag	aggagagattc	agtgaacta	900
gaatgccaga	tctcggtctat	acctccacca	aagcttttct	ggaaaagaaa	taatgaaatg	960
gtacaattca	acactgaccg	aataagctta	tatcaagata	acactggaag	agttacttta	1020
ctgataaaa	atgtaaacaa	gaaagatgct	gggttggtata	ctgtgtcagc	agttaatgaa	1080
gctggagtga	ctacatgtaa	cacaagatta	gacgttacgg	cacgtccaaa	ccaaactctt	1140
ccagctccta	agcagttacg	ggttcgacca	acattcagca	aatatttagc	acttaatggg	1200
aaaggtttga	atgtaaaaca	agctttttaac	ccagaaggag	aatttcagcg	tttggcagct	1260
caatctggac	tctatgaaag	tgaagaactt	taataacttt	accaacattg	gaaaacagcc	1320
aactacacca	ttagtaatat	atttgattac	atttttttga	aattaatcca	tagctgtatt	1380
aacagattat	ggtttttaatt	aggtaatat	gttaatatat	atttataata	ttatttatcc	1440
tttgactctt	gcacattcta	tgtacccctc	cgatttgtga	agcctacagg	aaatctgggt	1500
atatggattt	gtaactgcag	aagactatct	taaaatacag	gatttttaaca	tttaagtcac	1560
gcacatttaa	caattacagg	ttataaaatta	gtatcaactt	tttaaacaca	tctaattgctt	1620
gtaataacgt	ttactgggtac	tgttttctaa	atactgtttt	accctgtttt	tcttgttagga	1680
atactaacat	ggtatagatt	atctgagtgt	tccacagttg	tatgtcaaaa	gaaaataaaa	1740
ttcaaatatt	taaaacggaa	aaaaaaaa				1768

&lt;210&gt; 47

&lt;211&gt; 3287

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1295420CB1

&lt;400&gt; 47

```

cgtccccctga caccgacacc ccattgctcc cacagtctcc ccagtctcca ctttgggtccc 60
cagcgctgtc tgcccagagga tttgctgaa ggctgcccc aactctgcac ccgccccccg 120
agggccaccg agggccatga ctaagacaga tcttgcccc atggccccgc caccocgagg 180
aggaggaggaa gaagaggagg agggaggatga acccgtcccc gagggcccca gccccacca 240
ggagcgccgg cagaagcctg ttgtgcaccc ctccggcacct gccccctcc ctaaggacta 300
cgctttcacc ttcttcgatc ccaatgaccc ggctgcccag gagatcctgt ttgaccctca 360
gaccaccatc cccgagctgt ttgccattgt gcgccagtgg gtgccccaa g tccagcaca 420
gatagacgtc atcggcaatg agattctgcg ccgaggctgc catgtgaacg atcgtgacgg 480
gctgaccgac atgacactgc tccactatgc gtgcaaagct gggggccacg gaggcgggga 540
ccccgcggca gccgtgcgcc tctcgcagca gctgctggcg ctgggcgcag atgtgacgct 600
gcgcagccgc tggaccaaca tgaacgcgct tccactacgc gcctatattt atgtgcccga 660
cctcgtgcgt gtgctgctga aggggtgcgag gccgcgagtg gtgaactcca cgtgcagtga 720
cttcaaccac ggctcagccc tgcacatcgc tgcttccagc ctgtgcctgg gcgcgcgcaa 780
atgtttgctg gagcacggcg ccaaccctgc gctgaggaat cgaaaaggac aggtgccggc 840
ggaggtgggt ccagatccta tggacatgtc cctggacaag gcagagggcg cactggtggc 900
caaggagctg cggacgcttc tgggaagagg agtgccacta tcttgccgcc tccccaaagt 960
cacgctaccc aactatgaca acgtcccagg caatctcatg cttagcgcac tgggcttgcg 1020
cctgggagac cgcgtgctgc tggatggcca gaagacgggc aactgcggt tctgtgggac 1080
cacggagttt gccagcggcc agtgggtggg cgtggagctg gacgaacctg agggcaagaa 1140
cgatggcagc gttggggcg ttcgggtact catctgccct cccaagcagg gtctctttgc 1200
ctccgtgtcc aagatctcca aggcagtggc cgcaccccc tctctgtca cctccacacc 1260
ccggaacccc cggatggact tctcccgct caccggcaaa ggccgcaggg aacacaaagg 1320
caagaagaag acccatcat ccccatctct gggcagcttg cagcagcgtg acggggccaa 1380
ggctgaggtt ggagaccagg tccctgtcgc gggccagaag caggggatcg tgcgttcta 1440
cggaagaca gactttgccc caggttactg gtatggcatt gagctggacc agcccacagg 1500
caagcatgat ggctctgtct tccgtgtccg gtacttca tggccggtcc actgattccc cggggacag 1620
cttcgcacca gcatcccgta ttcagaggat atcaagtgc atagcgcag cccaaacgca ccttaccac 1680
cgttggagcc aaaaagtgc atcaagtgc atcaagtgc atcaagtgc atcaagtgc 1740
agtccggacc ccaaaggaca ttgcatcaga gggcgggagat gcagtcttag aggccttgga cacctgacaa 1800
ctgggtcccc tggatgctga gggcgggagat catctcctga cacccgagga gccctgagtc accctgagat 1860
agagacagag tccccactag ccagagtaga caccctgtt gaccctgtt agccagccct atgctaccca 1920
agagattccc agtaacacat ctcccataat aacccccaaa tacagacccc atgtcaccca 1980
ggccccatta cctgagtag caccttcagg ctagtcccta tccccaaacc ctacagcag 2040
gaaagagatt taacagattt ccataatcacc ccaaatgatg gtgacctct ccacataatg 2100
attcccagat gaacattctt gaatcaccca accctggatc agaaacctcc ccattaacaa 2160
cattacaaca ttaagtcctc ttgaaataaaa cataggtcac accccaaaag caaaagagta 2220
acactgcccc atgtcattgt tccccattta acatcagtc acatcagtc acatcagtc 2280
acagacattc tgaagccctt agattccaac cctcaatca gagacttcc cagacccact gtcaccccta 2400
atgggtaccc tcttatccct caagaagaaa cccaccataa gctaattaca ggacccccca agtcttcta 2460
agacccttgt accaaaacag cctcaagaaa cccccagtgc cttgtatgaa gcccaccca catggccac 2520
atcccaagtc cctcaagaaa tcccagaaaa ttctctatct ttttaagtaac gacttcccc 2580
agctcctgtg ctggccagac tcccagaaaa ggaggcccca ttctaggact ctgggggatcc caaacctag 2640
tttgggggac cccaaaattt ccctgtgccc tcaagtcta cagccctag aagaccccaa 2700
agtacacacg tcccgaactc ccaaactatg gaatccaaa tcccaggga atccaaatt 2760
tgccgtaact cctaggacc cccaggaaac ccaatcatga ggtccttgtg cctgggtatg 2820
tgaaaatcca atcccaagtc tgcattccag gctcccagac acctcaagcc ctattcacag 2880
aggagactgc agtcaggata ggaattccca tccctggaaa ctggagaatt tcaatgcccc 2940
gcaccaggaa accccacaca accaaaattc aagagcccca gccctaaggg aaccccaaat 3000
gagtccatgg gtttcaagac taaatggaag gcccgaagg cctgagggga tctcaaatcc 3060
cctaaagcct ccatctctaa acgttctagt cactggcctc aaaggacccc acagcacctg 3120
tggaaccccc acagctcgag ggagaacctg aaggccagg gggtcaggg cggacctggg 3180
ggccagacca ccaaggacag ctacagactg ccccttca ctatggccaa aaaaaaa 3240
tgactcctgt cctcttcaat aaagacgttt

```

&lt;210&gt; 48

&lt;211&gt; 1748

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1309364CB1

&lt;400&gt; 48

```

cgccatgttt gaaaagtgat gacgggttgac gtttgcgtgat ttttgacttt gctttagact 60
gctccccgaa ctgcgcgtct tcctgtcggc ggccggcact gtaggtgagc gcgagaggac 120
ggaggaagga agcctgcaga cagacgcctt ctccatccca aggcgcgggc aggtgcggg 180
acgctggggc tggcgggtgt ttogtcgtgc tcagcgggtg gaggaggcgg aagaaccag 240
agcctgggag attaacagga aacttccaag atggaaactt tgtctttccc cagatataat 300
gtagctgaga ttgtgattca tattcgcaat aagatcttaa caggagctga tggtaaaac 360
ctcaccaaga atgatcttta tccaaatcca aagcctgaag tcttgcacat gatctacatg 420
agagccttac aaatagtata tggaattcga ctggaacatt ttacatgat gccagtgaac 480
tctgaagtca tgtatccaca tttaatggaa ggcttcttac cattcagcaa tttagtact 540
catctggact cttttttgcc tatctgccgg gtgaatgact ttgagactgc tgatattcta 600
tgtccaaaag caaacaggac aagtcgggtt ttaagtggca ttatcaactt tattcacttc 660
agagaagcat gccgtgaaac gtatatggaa tttctttggc aatataaact ctctgcggac 720
aaaaagcaac agttaaacgc cgcacaccag gaggcattaa tgaaactgga gagacttgat 780
tctgttccag ttgaagagca agaagagttc aagcagcttt cagatgggat tcaggagcta 840
caacaatcac taaatcagga ttttcatcaa aaaacgatag tgctgcaaga gggaaatccc 900
caaaagaagt caaatatttc agagaaaacc aagcgtttga atgaactaaa attgttggtg 960
gtttctttga aagaaatata agagagtttg aaaacaaaaa ttgtggattc tccagagaag 1020
ttaaagaatt ataaagaaaa aatgaaagat acggtccaga agcttaaaaa tgccagacaa 1080
gaagtgggtg gaatatatga aatctatgga gactcagttg actgctgcc ttcatgtcag 1140
ttggaagtgc agttatatca aaagaaaata caggaccttt cagataatag ggaaaaatta 1200
gccagtatct taaaggagag cctgaacttg gaggaccaa ttgagagtga tgagtcagaa 1260
ctgaagaaat tgaagactga agaaaattcg ttcaaaagac tgatgattgt gaagaaggaa 1320
aaacttgcca cgcacaatt caaaataaat aagaagcatg aagatgttaa gcaatacaaa 1380
cgcacagtaa ttgaggattg caataaagtt caagaaaaaa gaggtgctgt ctatgaacga 1440
gtaaccacaa ttaatcaaga aatccaaaaa attaaacttg gaattcaaca actaaaagat 1500
gctgctgaaa gggagaaact gaagtcccag gaaatatttc taaacttgaa aactgctttg 1560
gagaaatacc acgacggtat tgaaaaggca gcagaggact cctatgctaa gatagatgag 1620
aagacagctg aactgaagag gaagtgttc aaaatgtcaa cctgattaac aaattacat 1680
gtctttttgt aaatggcttg ccatctttta atttctatt tagaaagaaa agttgaagcg 1740
aatggaag 1748

```

<210> 49  
 <211> 2163  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1315267CB1

```

<400> 49
gatgaattga gggaacagcc ttgtaaaatc aggaaagccg tccaaaagag cacttctgaa 60
aatcagactg aatggaatgc acgtgacgat gaaggtgttc caaatagtga cagtagcact 120
gactctgagg aacagcttga tgttaccata aaaccatcga ctgaggatag agagaggggc 180
atcagcagca gagaggatag ccacaaagtc tgtgatgata aggggccttt taaggacacc 240
aggacccaag aagataaaaag gagagatgtt gatctggatt tgtctgataa agattacagt 300
agcgatgagt ctatcatgga aagcataaaa ctgagccctc ctagatcctca 360
tccctaagtc tgagtaaaat ggactttgat gatgaaagaa cttggactga ccttgaagag 420
aatttgtgta accatgatgt tgttcttggg aatgaatcca cttatgggac gccgcagaca 480
tgctacccta ataatgaaat aggtatcctg gacaaaacaa taaaagggaa gattgcacca 540
gtcaagaggg gagaagactt gagcaagtcc aggaggagca gaagtcctcc tacatcggag 600
ctgatgatga aattctttcc ttctttgaaa ccaaaaccaa agtcagattc acacttggga 660
aatgaactca agttaaacat aagtcaagac caaccacctg gtgacaatgc tcgatcccag 720
gttttgagag agaaaattat tgaattggaa acagaaatag aaaagttaa agctgagaac 780
gcatctttag ctaaacttcg cattgaacga gaaagtgcct tggaaaaact caggaaagaa 840
attgcagact tcgaacaaca gaaagcaaaa gaattagctc gaatagaaga gtttaaaaag 900
gaggagatga ggaagctaca aaaggaacgt aaagtttttg aaaagtatac tacagctgca 960
agaacttttc cagataaaaa ggaacgtgaa gaaatacaga ctttaaaaca gcaaatagca 1020
gatttacggg aagatttgaa aagaaaggaa accaaatggt caagtacaca cagccgtctc 1080
agaagccaga tacaatgtt agtcagagag aacacagacc tccgggaaga aataaaagtg 1140
attgaaaagt tccgactgga tgccctggaag agagcagaag ccatagagag cagcctcgag 1200
gtggagaaga aggaagcgt tgcgaacaca tctgttcgat ttcaaaacag tcagatttct 1260
tcaggaaccc aggtagaaaa atacaagaaa aattatcttc caatgcaagg caatccacct 1320
cgaagatcca agtctgcacc tctcgtgat ttaggcaatt tggataaggg acaagctgcc 1380
tctcccaggg agccacttga accactgaac ctgaatataa agaggaggag 1440
gaagaccaag acatacagg agaaatcagt catcctgatg gaaaggtgga aaaggtttat 1500
aagaatgggt gccgtgttat actgtttccc aatggaactc gaaaggaagt gagtgcagat 1560

```

```

gggaagacca tcaactgtcac tttctttaat ggtgacgtga agcagggtcat gccagaccaa 1620
agagtgatct actactatgc agctgccag accactcaca cgacataccc ggagggactg 1680
gaagtcttac atttctcaag tggacaaata gaaaaacatt acccagatgg aagaaaagaa 1740
atcacgtttc ctgaccagac tgttaaaaac ttatttcctg atggacaaga agaaagcatt 1800
ttcccagatg gtacaattgt cagagtacaa cgtgatggca acaaaactcat agagtttaat 1860
aatggccaaa gagaactaca tactgccag ttcaagagac gggaataccc agatggcact 1920
gttaaaaccg tatatgcaaa cggtcacaa gaaacgaagt acagatccgg tcggataaga 1980
gttaaggaca aggagggtaa tgtgctaatt gacacggagc tgtgacgac ctcattgtgat 2040
catgaagtaa cagtaactga ctttttatgt taaaaaatgt acatttactg tggattctgt 2100
ttaattttatt gtgtatgtgt ggggaaaaga ttggattcta aaataaaagt ttaccctgtg 2160
gca 2163

```

<210> 50  
 <211> 1615  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1403289CB1

```

<400> 50
gtgggttcaga ggcagcttct agacctgcag gagggagatt gtattcagag gaagagcatc 60
atthttggcaa catctgaaag tgaaaacgga agccagaaac acttggccag ccctggggga 120
tttttttctt ctatgcctct ctggtggaat gacatttgct gtgtaggcat ctttctcttg 180
actgtatttc ttggccttga agagtactga gtttaaaaag acagtatgtg acagtccatg 240
gaaattgcct cttctgtgaa atctcgccac ctgctccgaa gacatgttgt tgtctcccaa 300
attctcctta tccaccattc acatacgact gacggccaaa ggattgcttc gaaaccttcg 360
acttcttcca gggtttagga gaagcactgt tgttttccac acagttgaaa agagcaggca 420
aaagaatcct cgaagcttat gtatccagcc acagacagct cccgatgcgc tgccccctga 480
gaaaacactt gaattgacgc aatataaaac aaaatgtgaa aaccaaagtg gatttatcct 540
gcagctcaag cagcttcttg cctgtggtaa taccaagttt gaggcattga cagttgtgat 600
tcagcacctg ctgtctgagc gggaggaagc actgaaacaa cacaaaacc tatctcaaga 660
acttgtaaac ctccggggag agctagtcac tgcttcaacc acctgtgaga aattagaaaa 720
agccaggaat gagttacaaa cagtgtatga agcattcgtc cagcagcacc aggctgaaaa 780
aacagaaaga gagaatcggc ttaaagagtt ttacaccagg gagtatgaaa agcttcggga 840
cacttacatt gaagaagcag agaagtacaa aatgcaattg caagagcagt ttgacaactt 900
aaatgctcgc catgaaacct ctaagttgga aattgaagct agccactcag agaaacttga 960
attgctaaag aaggcctatg aagcctccct ttcagaaatt aagaaaggcc atgaaataga 1020
aaagaaatcg ctgaagatt tactttctga gaagcaggaa tcgctagaga agcaaataca 1080
tgatctgaag agtgaaaatg atgctttaaa tgaaaaattg aaatcagaag aacaaaaaag 1140
aagagcaaga gaaaaagcaa atttgaaaaa tcctcagatc atgtatctag aacaggagtt 1200
agaaagcctg aaagctgtgt tagagatcaa gaatgagaaa ctgcatcaac aggacatcaa 1260
gttaaatgaaa atggagaaac tgggtggaca caacacagca ttggttgaca aattgaagcg 1320
tttcagcagc gagaatgaag aattgaaagc tcggatggac aagcacatgg caatctcaag 1380
gcagcttttc acggagcagg ctgttctgca agagtgcgtg gagaaggagt cgaaagtcaa 1440
caagcgactc tctatggaaa acgaggagct tctgtggaaa ctgcacaatg gggacctgtg 1500
tagccccaag agatcccca catcctccgc catccttttg cagtcaccaa ggaattcggg 1560
ctccttacat agccccagca tttcacccag atgacacgtc cccaaagtcc acaga 1615

```

<210> 51  
 <211> 1356  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1607607CB1

```

<400> 51
ccaccccgagg aaccgggaag tggaggagga ggcgcggcgg cggcgggcgg ggcggtctgcg 60
gtggccaagc aggcagatac tgccagaccc gttcccgga gctgtcttg gtttgggggc 120
gggagacagg ctgagccgcc tgggcggcct ggctgtacg gggcggggga ggccatggcc 180
tcggctgagt tgcaggggaa gtaccagaag ctggctcagg agtactcgaa gcttcgggct 240
cagaatcagg ttctgaaaaa aggtgttgtg gatgaacaag caaattctgc agctttaaag 300
gagcaactga aaatgaagga tcagtcattg agaaaactac aacaggaaat ggacagtttg 360
acatttcgaa atctgcagct tgccaagagg gtagaactac ttcaagatga actagctcta 420

```



```

agtgaaccac gaggaagaa aaacaagaaa agtggagaat cttcttctca gttgagtcaa 480
gagcagaaga gtgtctttga tgaagatctg caaaagaaga tagaagagaa tgaacgggtg 540
catatacaat tttttgaagc tgatgagcag cacaagcatg tggaagcaga gctgaggagt 600
cgactggcca ctctggagac agaagcagcc cagcaccaag ctgtggttga cggctcacc 660
cggaagtaca tggaaacat tgagaagctg cagaacgaca aggctaaact agaagtgaaa 720
tctcagactc tagaaaagga agccaaggaa tgtcgacttc gaacggaaga atgtcaatta 780
cagttaaaga ctcttcatga agatttgtca ggtagattag aggaatcctt atcaatcatc 840
aatgaaaaag taccttttaa tgatacaaaa tatagtcagt acaacgctct gaacgttcca 900
ctccacaata ggagacacca gctgaagatg cgagatattg ctgggcaggc cctggctttt 960
gttcaggatc ttgtgacggc tcttctaaac tttcatacct acacagaaca gaggattcaa 1020
atttttctctg ttgattctgc cattgacact atatctccat tgaatcagaa gttctcacia 1080
taccttcatg aaaatgcgtc ctatgtccgc cctcttgagg aaggaaatgct tcattttattt 1140
gaaagtatca ctgaggatag tgtgactgtc ttggagacaa ctgtgaaatt gaaactttt 1200
tcagaacact taacctccta catatgtttt cttagggaaga ttcttcccta tcagttaaaa 1260
aggtagttac ccccgagggc cagggaactt ggggaattgt ggggtgaacc tgatctggct 1320
ggcgtaata aaatattaac atgtgaaaga aaaaaa 1356

```

&lt;210&gt; 52

&lt;211&gt; 1268

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1660025CB1

&lt;400&gt; 52

```

gatggaagta agccttgagt attgagggtt tcagaggggg tgcattctacc ccagccccag 60
ggagtgggga ggcggaaga ggacctgcgg caggccctct tcggcagctc ctccggcccc 120
gtttccctcg gcgtgctact gtgcgctcga tccagacca tggggaagcg ggacaatcgg 180
gtggcctata tgaacccaat agcaatggcg agatcaaggg gtccaatcca gtcttcaggg 240
ccaacaatac aggattatct gaatcgacca aggcctacct gggaagaagt aaaagagcaa 300
ctagaaaaga aaaagaaagg ctccaaggct ttggctgaat ttgaagaaaa aatgaatgag 360
aactggaaga aagaactgga aaaacacagg gagaaattgt taagtggaag tgagagctca 420
tccaaaaaaa gacagagaaa gaaaaaagaa aagaagaaat ctggtaggta ttcattctct 480
tcttcatcaa gctctgattc ttccagcagt tcttctgatt ctgaagatga ggataagaaa 540
caaggaaaac ggagaaagaa aaagaagaac cgttcacata aatcttctga aagctccatg 600
tcagaacact aatcagacag taaggatagt ttaaaaaaga aaaagaagtc aaaagatgga 660
actgagaaaag aaaaggatat taaaggactc agcaaaaaga gaaagatgta ttctgaagat 720
aaacctttat catctgagtc cttgtcagaa tcagagtata ttgaggaggt gcgagcaaaa 780
aagaagaaaa gcagtgaaga acgagaaaaa gcaacagaaa aaacaaaaaa gaaaaagaag 840
cataagaaac acagtaagaa gaagaaaaag aaggctgcta gttcaagtcc tgactacca 900
taacattaag aaaaatcagg attcccttat aaagaaagt caatgtctga ggaaatttca 960
actgtgaaaa ctacaacata ttactaaaa tgcattgaatt ttcttgtttt tagaattatt 1020
cctggactat tcagtagcca ctcatatgcc actgtgtgaa agggccataa atgttgacct 1080
ctgcttgaac atctattttt ttctcttcca gtgcttgata actctgggag ataatacact 1140
gcagtcgtac tagtggttaa gatatttggg aataaaaatta atacttttga ctagaagcgt 1200
ctaaggataa accaacagaa attgaatctg gatacatctt taagatgtaa tcagaaatga 1260
ccagatgg 1268

```

&lt;210&gt; 53

&lt;211&gt; 2554

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1796836CB1

&lt;400&gt; 53

```

cggaaatcaa cttccggggg cagaggtgtt cgaagccggg tgggtgcgtg gctaccccaa 60
ctgtgtggc tgggcccggg tctccctca agggcctggg gccgtgcctc ggggtgtaagc 120
gtaggggtct gtgtgctggg ggtggctcac cgggcagcgt ggggtgagcgg cgcagcggcg 180
gcagcggaga gcgagagagg ggagcaggtg ccacttgaag aatggatgat gatgattttg 240
gtggttttga ggctgcggag actttttagt tggaagtgg tgaacccaa acaacattct 300
ctgctattcc ttgggctgcc tttcctgcag tatctggagt ccacttttca ccatctctc 360
ctgagattgt actggacgct gaccactctt cttccattgg ctgcctctct tctgatgcca 420

```

ttattttcatc	accagagaat	acacatgcag	caaatagcat	tgtgagtcaa	actattccaa	480
aagcacagat	tcagcaatca	acacacactc	atctggatat	ctcacttttt	ccattggggt	540
taactgatga	aaaaagtaat	ggaacaattg	cccttgtgga	tgattctgag	gatcctggag	600
ccaatgtatc	taacatacag	cttcagcaaa	aaatttcaag	tctggagatt	aaactcaaag	660
tatctgaa	agaaaaacag	agaattaaac	aggatgtgga	atcattgatg	aaaagcata	720
atgtcttaga	aaaaggcttt	ctaaaagaaa	aagagcaaga	ggccatttct	tttcaagata	780
gatacaaaaga	acttcaggaa	aaacataaac	aagaattgga	agacatgagg	aaagctgggtc	840
acgaagccct	cagcattatt	gtggatgaat	ataaggcact	actgcagtct	tcagttaaagc	900
aacaagtaga	agctattgaa	aaacagtcac	tttctgcaat	tgagaaacag	gcacacaagt	960
gtgaggagtt	gctaaatgct	cagcatcaga	ggctccttga	aatgctagat	acagagaagg	1020
aactgttaaa	agaaaaaata	aaggaaagctt	tgattcagca	atctcaagaa	cagaaggaaa	1080
tattggaaaa	gtgtttggag	gaagaaaggtc	aaagaaataa	agaggcatta	gtatccgctg	1140
caaagcttga	aaaagaagca	atgaaggatg	cagtttttaa	agtcgtagaa	gaagaaagaa	1200
aaaattttaga	aaaagcgcat	gctgaagaaa	gggaattatg	gaagacagaa	catgcaaaag	1260
atcaagaaaa	agtatctcag	gaaattcaaa	aagctataca	agaacaaaga	aaaataagtc	1320
aggaaactgt	taaggcagca	ataatagaag	agcagaaacg	aagtgaagaa	gctgtggaag	1380
aggcagtgaa	agaacaaga	gatgaattga	tagagtatat	aaaagaacag	aaaaggctcg	1440
atcaagtcac	ccgccaaaga	agcctgtcca	gtttggaact	gttccctctcc	tgtgcacaga	1500
aacagtttaag	tgctttaata	gctacggaac	cagttgacat	tgaataaaaa	gaacatgaca	1560
aaccacacact	ggcattggat	aaatcatatt	acaccttcaa	aatacacact	ctgaattata	1620
aagatgtgtt	tgttttcttt	ccaaatcatg	tagaattgat	ttccagttca	aggataaacc	1680
aaaacaatat	ttagaactat	caagtgatct	aatttatatt	cttttggttt	cttctttaca	1740
tttactgtta	ttttattatt	attagtagta	gcagcaacag	agtatgatat	gacccaaaag	1800
ccattgtaaa	gtgccacatt	accaaaatta	attaagtaaa	ctttatagcc	tgtgggagtc	1860
tattatata	tattttgcaa	aagtagtaaa	tatattattg	tttcatgatg	actcttgatg	1920
agatgctaga	atgtaaccat	acatttatct	tattttgagg	atagaaatag	catggatttc	1980
aacatcactt	atttatctgt	ataattggaa	ataaaacacc	gatatgatag	agaatcattc	2040
cggcattacc	taacctcttc	tcagtttgga	tctatgtatt	ttcattggtc	ttcattggtc	2100
aaacaataca	attaaaagca	ctaaagatta	ttatattaat	tcaactttga	tctgatata	2160
cacttaaaact	aaaggggtgt	gtgtggtgta	tgcttggttc	ctatttctgc	tctttaaaga	2220
tactttgaat	caataaaaacc	attagtctac	aatcaaat	gtgaacttaa	tctctagaaa	2280
gagaatataa	ctcagccatt	tataggaatt	taggttcaag	tacaggatat	atgaaatctt	2340
ttccagttat	acttaattgt	acttaattca	caggcaggat	gcttcaatgc	aaaatcatga	2400
atattttttaa	ttcaaaacta	aatgtcatt	aatatgtatg	tatgcaaatg	ttttatctta	2460
ttttctgaaa	tgcatctact	ttcatgggct	ttgtacgttt	ctgagatttc	tcagtgtaat	2520
aaaaagagct	cccaaactta	aaaaaaaaaa	aaaa			2554

&lt;210&gt; 54

&lt;211&gt; 1216

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2880670CB1

&lt;400&gt; 54

cgctgttgcc	cttagggagc	gctgtggggc	tgctgggggt	ggggggccga	agcgccagag	60
atggctgctc	agcgagggat	gccagctcc	gccgtgaggg	tccgtggaaga	ggcgttgggc	120
atgggtttga	cggcagccgg	ggacgcgagg	gacacggcgg	acgcgggtg	ggctgagggc	180
gcctactacc	tggaacaggt	caccataact	gaagcatctg	aagatgacta	tgaatatgaa	240
gagataccag	atgacaattt	tagcatccca	gaaggtgaag	aagatctggc	aaaagcaatt	300
cagatggccc	agaacagggc	tacagatact	gaaatttttg	aacggaaaac	agttcttcct	360
tcaaagcatg	cagtacctga	agtaatagaa	gactttctct	gcaatttctt	gatcaaatg	420
ggaatgacca	gaactcttga	ttgctttcag	tctgaatggt	atgagttaat	acagaaagga	480
gtgactgaac	ttagaactgt	tgggaatggt	ccagatgtct	acacccagat	tatgcttttg	540
gaaaatgaga	acaaaaattt	aaagaaagat	ttgaagcact	acaaacaagc	agctgagtat	600
gttatttttt	aatgacatt	ttcttctttt	tcttttgga	taaataaaag	agttgagtga	660
agctgatata	tgtaatatat	cagagcctta	atttttgaaa	actgaatttt	tctagtgtga	720
agaatgtga	gaggcttcat	tagcaaatta	attaacaga	tgatcagaac	tatcacaatt	780
ataacttacc	aacaagaagg	gaatgcaggt	agttgtttag	gagatggtac	attttttgata	840
taaaattcac	ttccttgtgt	atttgatagt	cttttcatgg	tttataacat	tttctcctgt	900
aaagataggc	taatttctga	aataataatt	aaatttatag	aaagccgaga	ggaaattgct	960
agtttatctc	tggtagagga	atttctgtat	ttgaaaattc	tccagaagga	ataatataaa	1020
ctgtggactt	tgggtgataa	tgatatgtag	gttcgtcagt	tgtaaacaaa	tgtatctctc	1080
tgttgggggc	tattgataat	ggggaaggct	gtgcatgtgt	gggagtagga	ggtgtatggg	1140
acatctctgt	accttctaatt	caattttgct	atgaacttaa	aactgctcta	gaaataaagt	1200

ttattataaaaa aaaaaa

1216

&lt;210&gt; 55

&lt;211&gt; 1457

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2913976CB1

&lt;400&gt; 55

gaattacagg	tttgagccac	tgcacctggc	caggaacctg	ttttttaacc	ctggjgcctt	60
aacagaggct	gtgtagaaat	ggggggcgtc	acaggccaag	ggcttagtcc	acttcagtcc	120
agcatccatg	gggaagcctg	gttgcaggag	agtccttcta	agtaggacaa	ggagcagtgg	180
ctgattgagg	gaccccgatc	agctgccact	gtgaccttcc	tgctgagaga	cttacgcagg	240
tcccttgctt	ccctgggcct	tggtttccct	ctcgggtacaa	tgccggcta	tgtacctgtt	300
accaccacaga	tgaggaacag	agctgggagg	gctgcatggg	gtccacttgg	atgactcccc	360
gtctgcatcc	ccacaggccc	cagcgggagg	accttcagtc	cagcccggtc	ccctcaggcc	420
catggaggaa	gagctgccac	ctcccccggc	agaacctgtt	gagaaagggg	catccacaga	480
catctgtgcc	ttctgccaca	agaccgtgtt	cccccgagag	ctggctgtgg	aggccatgaa	540
gaggcagtag	catgccccagt	gcttcacgtg	ccgcacctgc	cgccgcccagc	tggctgggca	600
gagcttctac	cagaaggatg	ggcgacccct	ctgcgaaccc	tgctaccagg	acacactgga	660
gaggtgcccgc	aagtgtggcg	aggtgggtccg	ggaccacatc	atcaggggccc	tgggcccaggc	720
cttccacccc	tcctgcttca	cgtgtgtgac	ctgcgcccgg	tgcatgtggg	atgagagctt	780
tgccctggggc	agccagaacg	aggtgtactg	cctggacgac	ttctacagga	aattcgcccc	840
cgtctgcagc	atctgtgaaa	atcccatcat	ccctcgggat	gggaaagatg	ccttcaaaat	900
cgaatgcatg	ggaagaaact	tcctatgaaa	ttgctacagg	tgtgaggact	gcaggatcct	960
cctgtctgtc	gagccacagg	accaaggtcg	ctacccccctg	aacaaccatc	tcttctgcaa	1020
gccatgccat	gtgaagcggg	gtgctgcggg	gtgctgctga	gagtgcgccg	tgggcagtga	1080
acagaccact	agccccggct	ggggcccttc	acccgccttg	gcctcccaaa	gtgctggaat	1140
tacaggtgtg	agccactgtg	cccagctgag	taaatttctt	gattgcacag	aatgtacggt	1200
gatattggcg	gacttaaggga	catcgaattg	tttatcagga	ataaagtatt	atgtgtgttt	1260
tctgcggccc	tggataatgc	tgtagcattc	agggtcgatt	gagtaaaaaa	aatgtagaga	1320
tggtttccctg	tttgtttttg	cttctagttt	tcattcattt	gctattttatt	ctcttctggc	1380
tttgcttgtg	tatgcatata	tataaaaacca	ttattattat	tatcttttagt	ttctagtggg	1440
aggcttttaa	aaaaaaa					1457

&lt;210&gt; 56

&lt;211&gt; 1636

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 3092084CB1

&lt;400&gt; 56

ggcgggagct	caggaccggc	gccttctcct	tgcttctggg	ggtcgtggcc	ttgctcccgc	60
tgtgcgggaa	agaatccag	gcccttccac	gcgcgtgtgg	gtgcgggggc	ccgaagtgc	120
tcgtgggttc	ccgctaggtc	tccgctgggg	caggaaccgg	aatcatgggt	gggaccacca	180
gcacccgccc	ggtcaccttc	gaggcggagc	agaatgagaa	catcacctg	gtgaagggca	240
tccggctttc	ggaaaatgtg	attgatcgaa	tgaaggaaatc	ctctccatct	ggttcgaagt	300
ctcagcggta	ttctgggtgt	tatgggtgct	cagtttctga	tgaagaattg	aaaagaagag	360
tagctgagga	gctggcattg	gagcaagcca	agaaagaatc	cgaagatcag	aaacgactaa	420
agcaagccaa	agagctggac	cgagagaggg	ctgctgccaa	tgagcagtta	accagagcca	480
tccttcggga	gaggatatgt	agcgaggagg	aacgcgctaa	ggcaaagcac	ctggctaggc	540
agctggaaga	gaaagaccga	gtgctaaaga	agcaggatgc	attctacaaa	gaacagctgg	600
ctagactgga	ggagaggagc	tcagagttct	acagagtcac	cactgaacaa	tatcagaaag	660
ctgctgaaga	ggtggaagca	aagttcaagc	gatatgagtc	tcattccagtc	tgtgctgatc	720
tgcaggccaa	aattcttcag	tgttaccgtg	agaacaccca	ccagaccctc	aaatgctccg	780
ctctggccac	ccagtatatg	cactgtgtca	atcatgccaa	acagagcatg	cttgagaagg	840
gaggataaaa	acttttcagaa	tgagcaaaaac	accatcaacg	tttaattccag	agatggaaca	900
ttttttttcc	tagtgagaaa	acaacccatt	tgaagagaag	accactaatg	agaagaccac	960
taaagagaga	catcaagaat	ggattcagca	gaatcatttc	acgtttttgaa	cagcagcagt	1020
ttgaagggcc	aaagccttga	tcagggatca	gtcattaaag	gacactcttg	agtattagta	1080
aacctcttta	tgatgattaa	aagagaaggg	cagccctctc	caccttttgg	tactttctat	1140

```

tcaacttgca ctgaccataa aatgtttctc ttctgaacaa gcccacatcat ttggtgaacc 1200
tocaccctaa caaagtagga tgggggtggg ggctaaatta attggagtgg ggcgaggaga 1260
gagccagaaa acatagatcc gagggcagca gtgctgggtg gagagagcca gaaaacagat 1320
ctggaggcag cagtgcctgga tggaaattgtc taggctgtgg catgttggtt ttgtctttct 1380
tttctccttt gattatgtaa gagctatttc attataactt attatggtga ttatacaggc 1440
aagaagacaa aaaggagaga aaatgtacct cttctactgg aataatgttt atgattacaa 1500
gtgagataag gtatTTTTat caatatgaag gcaaccttgg ctgataaaac ctctatagtg 1560
aatactcaca tctttacttc actcactatc aataataaat atattttctg acaaagaaaa 1620
aaaaaaaaaa aaaaaa
1636

```

&lt;210&gt; 57

&lt;211&gt; 1742

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 3882482CB1

&lt;400&gt; 57

```

gctgccatgg caacactacg ccgccgccgc cttttcacgc acgtcgcgag ctaacggact 60
cggcgccggc ggccggccgg gcctgcgcc caccgcacc ccatctggac cgcctcgtg 120
aatgtgcccg gacctgcgcc ttctgggtct ctgaaagaag atgaatttgg ctgagatttg 180
tgataatgca aagaaaggaa gagaatatgc ccttcttggg aattacgact catcaatggt 240
atattaccag ggggtgatgc agcagattca gagacattgc cagtcagtca gagatccagc 300
tatcaaaggc aaatggcaac aggttcggca ggaattattg gaggaatatg aacaagttaa 360
aagtattgtc agcacttttag aaagttttta aattgacaag cctccagatt tccctgtgtc 420
ctgtcaagat gaaccattta gagatcctgc tgtttggcca cccctgttc ctgcagaaca 480
cagagctcca cctcagatca ggcgtcccaa tcgagaagta agacctctga ggaaagaaat 540
ggcaggagta ggagcccggg gacctgtagg ccgagcacat cctatatcaa agagtgaana 600
gccttctaca agtagggaca aggactatag agcaagaggg agagatgaca agggaaggaa 660
gaatatgcaa gatggtgcaa gtaatggtga aatgccaaaa tttgatgggt ctgggttatga 720
taaggatctg gtggaagccc ttgaaagaga cattgtatcc aggaatccta gcattcattg 780
ggatgacata gcagatctgg aagaagctaa gaagttgcta agggaagctg ttgttcttcc 840
aatgtggatg cctgactttt tcaaagggat tagaaggcca tgggaagggt tactgatggt 900
tggaccccca ggcactggta aaactatgct agctaaagct gttgccactg aatgtggtac 960
aacattcttc aacgtttcgt cttctacact gacatctaaa tacagagggt aatctgagaa 1020
gttagttcgt ctgttgtttg agatggctag attttatgcc cctaccacga tcttcattga 1080
tgagatagat tctatctgca gtcgaaaggg aacctctgat gaacatgagg caagtcgcag 1140
ggtcaagtct gaactgctca ttcagatgga tggagttgga ggagctttag aaaatgatga 1200
tccttccaaa atggttatgg tattggctgc tactaatttc ccgtgggaca ttgatgaagc 1260
tttgcgaaga aggttagaaa aaaggatata taccctctc ccaacagcaa aaggaagagc 1320
tgagcttctg aagatcaacc ttctgtagg gctattctgg cgaattagat cctgatattc aactggaaga 1380
tatagccgag aagattgagg gctattctgg tgctgacatc actaatgttt gcagggatgc 1440
ctctttaatg gcaatgagac ggcgtatcaa tggcttaagt ccagaagaaa tccgtgcact 1500
ttctaaagag gaacttcaga tgccgtgttac caaggagac tttgaattgg ccctaaagaa 1560
aattgctaag tctgtctctg ctgcagactt ggagaagtat ggggttgatt tggttgatt 1620
tggatctgct tgaatttctg tcagctcttt aatttctggt atttttgttg ataaaatacg 1680
aagaaattcc tgcaattttt aaaaaacaag tttggaattt tttcagtgga gtggttttctg 1740
ct
1742

```

&lt;210&gt; 58

&lt;211&gt; 602

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 4933451CB1

&lt;400&gt; 58

```

ctgagttcgg cagctcacac agcacccttt tcaagttggt ttgcagtggg tgccctcttgt 60
gagacactgt ttctgagagc agcttttgtg gcattctaca gggcagattt ctgggtccac 120
ccacctctgc ctccgccatg gctgcaagaa ccgttatcat tgaccacggg tctggctttt 180
tgaaggctgg cacggccggc tggaaatgag ctcagatggt cttcccgaa atcgtgaact 240
acctaccgtg caaggagaa cctggcccca gctatgcccg taagcgtgtg agcctgggca 300
tcgacatttg ccattctgac accttttagt accccatcga gcggggccgc atctcaact 360

```

gggaggggtgt	gcagtacctc	tggtcattttg	tgttggagaa	ccacagacgg	gagcaagagg	420
tccccctgt	gatcatcacg	gagacaccct	tgagggagcc	tgcggaccga	aagaagatgt	480
cattccttgg	aaccctgcag	gggacagttt	ttccaggggtg	gcctatcatt	ggggatatgag	540
tggtctgactg	ccatctccat	gcctgagagc	actgattttt	tcatttggcat	ttccccctggt	600
tt						602

&lt;210&gt; 59

&lt;211&gt; 3237

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5043904CB1

&lt;400&gt; 59

gaaaatgtct	gatgctcagg	gcagctacaa	actggatgaa	gctcaggctg	tcttgagaga	60
aacaaaagcc	atcaaaaagg	ctattacctg	tggggaaaag	gaaaagcaag	atctcattaa	120
gagccttgcc	atgttgaagg	acggcttctg	cactgacagg	gggtctcact	cagacctgtg	180
gtccagcagc	agctctctgg	agagttcgag	tttcccgccta	ccgaaacagt	acctggatgt	240
gagctoccag	acagacatct	cgggaagctt	cggcatcaac	agcaacaatc	agttggcaga	300
gaaggtcaga	ttgcgcttcc	gatatgaaga	ggctaagaga	aggatcgcca	acctgaagat	360
ccagctggcc	aagcttgaca	gtgaggcctg	gcctgggggtg	ctggactcag	agagggaccg	420
ctcgatcctt	atcaacgaga	aggaggagct	gctgaaggag	atgcgcttca	tcagcccccg	480
caagtggacc	caggggggagg	tggagcagct	ggagatggcc	cgggaagcggc	tggaaaagga	540
cctgcaggca	gcccgggaca	cccagagcaa	ggcgcctgacg	gagagggttaa	agttaaacag	600
taagaggaac	cagcttctga	gagaacttga	ggaagccacc	cggcagggtgg	caactctgca	660
ctcccagctg	aaaagtctct	caagcagcat	gcagtcctctg	tcctcaggca	gcagcccccg	720
atccctcacg	tccagccggg	gctccctggt	tgcattccagc	ctggactcct	ccacttcagc	780
cagcttcact	gacctctact	atgacctctt	tgagcagctg	gactcagagc	tgcagagcaa	840
ggtggagttc	ctgctcctgg	agggggccac	cggcttcccg	ccctcaggct	gcataccacc	900
catccacgag	gatgaggtgg	ccaagaccct	gaaggcagag	ggaggtggcc	gcctgcaggc	960
tctgcgttcc	ctgtctggca	ccccaaagtc	catgacctcc	ctatccccac	gttccctctt	1020
ctcctcccc	tccccacct	gttccccctt	catggctgac	ccccctctgg	ctgggtgatgc	1080
cttctcaac	tccttggagt	ttgaagacc	ggagctgagt	gccactcttt	gtgaactgag	1140
ccttggtaac	agcgcccagg	aaagataccg	gctggaggaa	ccaggaaacgg	agggcaagca	1200
gctggggcaa	gctgtgaata	cggcccaggg	gtgtggcctg	aaagtggcct	gtgtctcagc	1260
cgcggtatcg	gacgagtcag	tggctggaga	cagtgggtgtg	tacgaggtt	cggtgcagag	1320
actgggtgct	tcagaagctg	ctgcatttga	cagtgcagaa	tcgggaagcag	tgggtgcgac	1380
ccgaattcag	attgccctga	agtatgatga	gaagaataag	caatttgcaa	tattaatcat	1440
ccagctgagt	aacctttctg	ctctgttgca	gcaacaagac	cagaaagtga	atatcccggt	1500
ggctgtcctt	ccttgcctctg	aaagcacaac	ctgcctgttc	cggacccggc	ctctggagcg	1560
ctcagacact	ctagtgttca	atgaggtgtt	ctgggtatcc	atgtcctatc	cagcccttca	1620
ccagaagacc	ttaagagtcg	atgtctgtac	caccgacagg	agccatctgg	aagagtgcct	1680
gggaggcgcc	cagatcagcc	tggcggaggt	ctgccggctt	ggggagaggt	cgactcgctg	1740
gtacaacctt	ctcagctaca	aatacttgaa	gaagcagagc	agggagctca	agccagtggg	1800
agttatggcc	cctgcctcag	ggcctgccag	cacggacgct	gtgtctgctc	tgttggaaac	1860
gacagcagtg	gagctggaga	agaggcagga	ggcgaggagc	agcacacaga	cactggaaga	1920
cagctggagg	tatgaggaga	ccagtgcagaa	tgaggcagta	gccgagggaag	aggaggagga	1980
ggtggaggag	gaggagggag	aagaggatgt	tttcaccgag	aaagcctcac	ctgatatgga	2040
tgggtaccca	gcattaaagg	tggacaaaaga	gaccaacacg	gagacccccg	ccccatcccc	2100
cacagtgggtg	cgacctaaagg	accggagagt	gggcaccccg	tcccaggggc	catttcttccg	2160
agggagcacc	atcatccgct	ctaagacctt	ctccccagga	ccccagagcc	agtacgtgtg	2220
ccggctgaat	cggagtgcata	gtgacagctc	cactctgtcc	aaaaagccac	cttttgttccg	2280
aaactccctg	gagcgacgca	gcgtccgcat	gaagcggcct	tcctcgggtca	agtcgctgcg	2340
ctccgagcgt	ctgatccgta	cctcgctgga	cctggagttta	gacctgcagg	cgacaagaac	2400
ctggcacagc	caactgacct	aggagatctc	ggtgctgaag	gagctcaagg	agcagctgga	2460
acaagccaag	agccacgggg	agaaggagct	gccacagtgg	ttgcgtgagg	acgagcgttt	2520
cgccttgctg	ctgaggatgc	tggagaagcg	gcagatggac	cgagcgggagc	acaaggggtga	2580
gcttcagaca	gacaagatga	tgaggggcagc	tgccaaggat	gtgcacaggc	tccgaggcca	2640
gagctgtaag	gaacccccag	aagttcagtc	tttcaggggag	aagatggcat	ttttcaccgg	2700
gcctcggtg	aatatcccag	ctctctctgc	agatgacgtc	taatcgccag	aaaagtattt	2760
cctttgttcc	actgaccagg	ctgtgaacat	tgactgtggc	ttaaagttatt	tatgtggtgt	2820
tatatgaagg	tactgagtc	caagtcctct	agtgcctctt	ttggtttgaa	gatgaaccga	2880
cttttttagtt	tgggtccctac	tgttgttatt	aaaaacagaa	caaaaaacaaa	acacacacac	2940
acacaaaaaac	agaaacaaaa	aaaaccagca	ttaaaataat	aagattgtat	agtttgtata	3000
tttaggagtg	tatttttggg	aaagaaattt	taaatagaact	aaagcagtat	tgagttgctg	3060

ctcttcttaa	aatcggttag	atctttttttg	gtttgtacag	ctccaccttt	tagaggctctt	3120
actgcaataa	gaagtaatgc	ctgggtgggac	ggtaatccta	ataggacgct	ccgcacttgt	3180
cacagtacag	ctaatttttc	ctagttaaca	tattttgtac	aatatttaga	atgcacg	3237

&lt;210&gt; 60

&lt;211&gt; 3640

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5202390CB1

&lt;400&gt; 60

acctggaata	aaaaatccct	atcggtggtgt	tgttggtgtgg	cctgttcctg	aaaacattga	60
aatcactgta	acactttttaa	aggatcctca	tgcggaagaa	tttgaagaca	aagagtggac	120
atltgtcata	gaaaatgaat	ccccttctgg	tcgaaggaaa	gctcttgcta	ctagcagcat	180
caatatgaaa	cagtatgcaa	gccctatgcc	aactcagact	gatgtcaagt	taaaattcaa	240
gccattatct	aaaaaagttg	tatctgccgc	tcttcagttt	tcattatctt	gcatttttct	300
gagggaaagga	aaagccacag	atgaagacat	gcaaagtttg	gctagtttga	tgagtatgaa	360
gcaggctgac	attggcaatt	tagatgactt	cgaagaagat	aatgaagatg	atgatgagaa	420
caagagtgaac	caagaagaaa	aggcagctaa	aattacagag	cttatcaaca	aacttaactt	480
tttggtatgaa	gcagaaaagg	acttggccac	cgtgaattca	aatccatttg	atgatcctga	540
tgctgcagaa	ttaaattccat	ttggagatcc	tgactcagaa	gaacctatca	ctgaaacagc	600
ttcacctaga	aaaacagaag	actctttttaa	taataacagc	tataatccct	ttaaagaggt	660
gcagactcca	cagtatttga	acccattcga	tgagccagaa	gcattttgtga	ccataaagga	720
ttctcctccc	cagctctaca	aaagaaaaaa	tataagacct	gtggatatga	gcaagtacct	780
ctatgtgat	agttctaaaa	ctgaagaaga	agaattggat	gaatcaaatc	ctttttatga	840
acctaaatca	actcctcctc	caaataattt	ggtaaattcct	gttcaagaac	tagaaactga	900
aaggcgagtg	aaaagaaaagg	ccccggctcc	accagtcctc	tcaccaaaaa	caggagtatt	960
aaatgaaaac	acagtttctg	caggaaaaga	tctctctact	tctcctaagc	caagccctat	1020
accaagtctc	gttttggggc	gaaagccaaa	tgctagtccag	tctttgcttg	tatggtgtaa	1080
agaagttaca	agaactacc	gaggagtaaa	aatcaccaat	tttactacat	cgtggagaaa	1140
tggtttatct	ttttgtgcaa	tattacacca	ctttagacca	gatttaattg	actacaagtc	1200
tctgaatcct	caagatatta	aagagaacaa	caaaaaggca	taogatggat	ttgccagcat	1260
aggaatttcc	cgattatttg	aaccttctga	tatggtatta	ttagcaattc	ctgataaact	1320
gactgttatg	acttatctct	atcaaaataag	ggcacatttc	agtggccaag	aactaaatgt	1380
cgttcagata	gaggaaaaca	gcagtaaaag	cacatataaa	gttggaaact	atgaaacaga	1440
tacaaacagt	tctgttgatc	aagaaaaatt	ctatgcagag	cttagtgatc	tgaagcggga	1500
gcctgaacta	caacagccta	tcagcggagc	agtagacttc	ttatcacagg	atgactctgt	1560
atltgttaaa	gatagcgggg	ttggagagtc	agaaagttag	catcaaaact	ctgatgatca	1620
ccttagtcca	agcacagcct	ccccttactg	tcgcaggact	aaaagtgaac	cagaacccca	1680
gaagtctcag	cagagctctg	gaaggacttc	aggatctgat	gacctgggaa	tatgttccaa	1740
tacagattca	acccaagcac	aggttttgtt	aggcaaaaag	agactattga	aagctgagac	1800
tttagaattg	agtgacttat	atgttagtga	taagaagaag	gatattgtct	caccctttat	1860
ttgtgaggag	acagatgaac	aaaagcttca	aactctagac	atcggtagta	acttgagaaa	1920
agaaaaatta	gagaattcca	gatccttaga	atgcagatca	gatccagaat	ctcctatcaa	1980
aaaaacaagt	ttatctccta	cttctaaact	tggaactca	tatagtagag	atctagacct	2040
tgctaagaaa	aaacatgctt	ccctgaggya	gacggagtct	gatccagatg	ctgatagaac	2100
cacttttaaat	catgcagatc	attcatcaaa	aatagtccag	catcgattgt	tatctagaca	2160
agaagaactt	aaggaaaagag	caagagttct	gcttagagcaa	gcaagaagag	atgcagcctt	2220
aaaggcgggg	aataagcaca	ataccaacac	agccacccca	ttctgcaaca	ggcagctaag	2280
tgatcagcaa	gatgaagagc	gacgtcggca	gctgagagag	agagctcgtc	agctaatagc	2340
agaagctcga	tctggagtga	agatgtcaga	acttcccagc	tatggtgaaa	tggtctgcaga	2400
aaagttgaaa	gaaaggtcaa	aggcatctgg	agatgaaaaat	gataatattg	agatagatac	2460
taacgaggag	atcccctgaag	gctttgttgt	aggaggtgga	gatgaactta	ctaacttaga	2520
aatgacctt	gatactcccg	aacaaaacag	taagttggtg	gacttgaagc	tgaagaagct	2580
cctagaagtt	cagccacagg	tggaacattc	accctccagt	gctgcccaga	aagctgtaac	2640
tgagagctca	gagcaggaca	tgaaaagtgg	cacagaagat	ctccggactg	aacgattaca	2700
aaaaacaaca	gaacgtttta	gaaatcctgt	tgtgttcagc	aaagattcta	cagtcagaaa	2760
aactcaactt	cagtcatttca	gccaatatat	tgagaataga	ccagagatga	aaaggcagag	2820
atcaatacag	gaagatacaa	agaaaggaaa	tgaggagaag	gcagcgataa	ctgaaactca	2880
gaggaagcca	tcagaagatg	aagtgtctaa	taaagggttc	aaagacacca	gtcagtatgt	2940
agtaggagaa	ttggcagcac	tagagaatga	gcaaaagcaa	attgacaccc	gtgccgcgct	3000
ggtggagaag	cgccttcgct	atctcatgga	cacaggaagg	aacacagaag	agaagaagc	3060
tatgatgcag	gaatggttta	tgtaggttaa	taagaaaaat	gccttaataa	ggagaatgaa	3120
tcagctctct	cttctggaaa	aagaacatga	tttagaacga	cggtatgagc	tgctgaaccg	3180

```

ggaattgagg gcaatgctag ccattgaaga ctggcagaag accgaggccc agaagcgacg 3240
cgaacagctt ctgctagatg agctggtggc cctggtgaac aagcgcgatg cgctcgtcag 3300
ggacctggac ggcgaggaga agcaggccga agaagaagat gagcatttgg agcgaactct 3360
ggagcaaaac aaaggcaaga tggccaagaa agaggagaaa tgtgttcttc agtagccatc 3420
agatcagaaa gaatctctcc caacatttta gagtcttggc tcccaaacca gaaaaagtca 3480
gactcattgt tgatttaaaa cttttaacat tttgtttggc tggattgtac tactttacct 3540
ctactttacc accaccaccc ttttctctcc tcttttccaa ataataataca gaactccaaa 3600
atagcttcat ttaaggattt ttttgtgagt taacaatttc 3640

```

```

<210> 61
<211> 2111
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 5526375CB1

```

```

<400> 61
gcaggacatg gaagaacgcg ggtcaccgga cgggggatctc ggcgggagcc tggagcaagg 60
gccagagggg ccggaacgc ccatccaggt ggtgctcagg gtacgtccca tgagcgcggc 120
cgagctgcgt cgagggcagc agagcgtgct gcaactgctca gggaccgga ctctgcaggt 180
gagtcctcca ggccgggggtc cagaagtggc gttccgcttc ggtgcgggtgc tagacgcggc 240
gcgcacgcag gaggacgtgt tccgggcgtg cggcgtgcgg cgcttggggg agctggcgct 300
gcgcgggtttc tctgcactg ttttcaacct tggccagacg ggctctggga agacctacac 360
cctgactgga cccctcccc agggggaggg ggtgcctgta ccccccagcc tggctggcat 420
catgcagagg accttcgcct ggctgttggg ccgcgtgcag cacctgggtg cccctgtcac 480
ccttcgcgcc tcttatctgg agatctacaa tgggcaggtt cgggacttgc tgagcctggg 540
gtctccccgg cccctccctg ttcgctggaa caagactcgg ggcttctatg tggagcagct 600
gcgggtggtg gaatttggga gtctggaggc cctgatggaa cttttgcaaa cgggtctcag 660
ccgtcgaagg aactcagccc acaccctgaa ccaggcctcc agccgaagcc atgcctgct 720
caccctttac atcagccgtc aaactgccc gcatatgcct tctgtggacc ctggggagcc 780
ccctgttggg gggaagctgt gctttgtgga cctggcaggc agtgagaagg tagcagccac 840
gggatcccggt ggggagctga tgcttgagg taacagcatc aaccgaagcc tgctggccct 900
gggtcactgc atctccctgc tgctggacc acagcggag cagagccaca tccctttccg 960
ggacagcaag ctacccaagt tgctggcaga ctactggga gggcgcgggg tcaccctcat 1020
ggtggcctgc gtgtccccct cagcccagtg ccttcttgag actctcagca cctgcgata 1080
tgcaagccga gctcagcggg tcaccacccg accacaggcc cccaagtctc ctgtggcaaa 1140
gcagccccag cgtttggaga cagagatgct gcagctccag gaggagaacc gtcgcctgca 1200
gttccagctg gaccaaattg actgcaaggc ctccaggctc agtggagccc ggggtggcctg 1260
ggcccagcgg aacctgtacg ggatgctaca ggagttcatg ctagagaatg agaggctcag 1320
gaaagaaaag agccagctgc agaataagcc agagctagcc cagaatgagc agcgcacct 1380
ggcccagcag gtccatgcac tagagaggcg tctctctct gactgctacc atcaccagca 1440
gggtcctggc ctgacccac cgtgtccctg ctgtatggcc ccagctcccc cttgccatgc 1500
actgccaccc ctctactcct gcccctgctg ccacatctgc ccactgtgtc gactgcccct 1560
ggcccactgg ggctgcctgc caggggagca ccacctgccc cagectctct tctgggctct 1620
gaggagtcag aaatagacca gacgtggttc ctgctctcag gaggcttcta gtctcaggag 1680
aggacggtag aagaaccatc tgctgcacac agaaggcag caagacccca cccctttatg 1740
ccatcctgac attcttggta ccttgtgggg taggcatggt ggggtgtcat ggctagcacc 1800
aggtaggtgg gacacagaga ggcaggactg aggtcaccca gactggactc tgggtctttc 1860
tacaggtggt ggaccctgag gcctcaggtg gcaggcccc atctgcccgg cccccacct 1920
gggcagcccc atgcagccct ggctctgcca agtgcccaag agagaggagt cacagtact 1980
ggactcagac ccgagtcctg gcagagatgt tgacggagga ggaggtggt ccttctgcac 2040
ctcccctgcc tgtgaggccc ccgaagacat caccagggt cagaggtggg gccgggggtc 2100
aaacctggcc c 2111

```

```

<210> 62
<211> 1389
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 5677408CB1

```

```

<400> 62
cccagactcc ttgcggactc gccgcctgat tctaggtctg tcaactactcc gagcctgtga 60

```

```

cgtttgccggc agccaggccg tcgacgatgc ccagtgaaac tctctgggaa attgcaaaag 120
ctgaagtgggaa aaaaagggga attaatggaa gtgaagggtga tggagctgaa attgcagaaa 180
aattttgtttt cttcattggc agtaaaaatg ggggaaagac tactattatt ctaagggtgtc 240
ttgacagaga tgaaccacca aaaccaacct tagcttttga atatacatat ggaagaagag 300
caaaaggggca caacacacca aaagatatcg ctcacttttg ggaactcggg ggaggaacct 360
ctttatttggga cttaatcagc atacccatca caggtgacac cttacggacg ttttctcttg 420
ttctcgttct ggatctttca aaacctaatg atctctggcc caccatggaa aatctcttgc 480
aagccacaaa aagccatgta gacaaagtga taatgaaact gggaaagaca aatgctaaag 540
cagtttctga aatgagacag aagatctgga ataatatgcc gaaggatcat cctgatcatg 600
aattaattga ccattttccg gtacctctgg tcataattgg aagtaaatat gatgttttct 660
aggattttga gtctgagaag agaaaggtaa tatgcaagac acttcgattt gttgcacatt 720
attatggagc atcattaatg tttaccagta aatcagaagc tctattacta aaaatacgtg 780
gagttatcaa ccagttggca tttggcattg acaaaagcaa atcaatatgt gtggatcaga 840
ataaaccgct gtttatcaca cgaggattgg attctttcgg tcaaatagga tctctcctg 900
ttcctgaaaa tgacattgga aagcttcatg ccactcacc tatggagtgtg tggaaaaaag 960
tgtatgaaaa gctctttcca ccaaagagta ttaacacgct gaaagatatc aaggaccctg 1020
cgagagatcc tcagtatgct gaaaatgaag tcgatgagat gagaattcag aaggatctgg 1080
aactggaaca gtacaaaaga agttcttcca agtcttggaa acaaatcgag cttgattctt 1140
gaacctattt caattattgt atatttattt ctctttttcc aaatacaaat aagatttatac 1200
tgtgaattaa ctattgtggc aatatgtgaa gaaagttaaa ctgtataatt tgttaaagga 1260
caagctggat ttcttggact agtgcacatc cctgtatata ttgaagcttt ttaaaaggaa 1320
aaattattgt agaaccacgt gtaatttttt ttaaaataaa agaatcttct actacctaca 1380
aaaaaaaaa

```

```

<210> 63
<211> 3331
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 5982278CB1

```

```

<400> 63
gtgcgcgccc gcttcgcaga gcaccgcgcc ttageccgca agttctagtt cttgctgccc 60
gtcctaaccgt cccgcagtc tgcgcagcca gccgtcccgc atgcgcgctt gggcgccgtg 120
gagcctgctg ccatgaagtc agcgagagct ggggtatact gtaggggtgc cgtgaaaaaa 180
gggtcccaaa cgaaccttaa agacccagtt ggggtatact gtaggggtgc cccactgggc 240
tttctgatc aagagtgttg catagaagtg atcaataata caactgttca gcttcatact 300
cctgagggct acagactcaa ccgaaatgga gactataagg agactcagta ttcattttaa 360
caagtatttg gcactcacac caccagaag gaactctttg atgttgtggc taatcccttg 420
gtcaatgacc tcattcatgg caaaaatgg catatggtgt gacgggaagt 480
ggaaaaaact acacaatgac tggttctcca ggggaaggag ggctgcttcc tcgttgtttg 540
gacatgatct ttaacagtat agggtcattt caagctaaac gatatgtttt caaatctaata 600
gataggaata gtatggatat acagtgtgag gttgatgcct tattagaacg tcagaaaaga 660
gaagctatgc ccaatccaaa gacttcttct agcaaacgac aagtagatcc agagtttgca 720
gatatgataa ctgtacaaga attctgcaaa gcagaaggag tgatgaaga tagtgtctat 780
gggtgtattt tctcttatat tgaaatatat aataattaca tatatgatct attggaagag 840
gtgcgcgttt atcccataaa cccaaacctc cacaatctaa attgcttctg gaagattaag 900
aaccataaca tgtatgtttg aggatgtaca gaagttgaag tgaaatctac tgaggaggct 960
tttgaagttt tctggagagg ccagaaaaag agacgtattg attaaattag ttcaggctcc tttgaatcgt 1020
gagtcacgcc gttcccatag cgtgttcaac attaaattag ttcaggctcc cttggatgca 1080
gatggagaca atgtcttaca ggaaaaagaa caaatcacta taagtcagtt gtccttggtg 1140
gatcttgctg gaagtgaag aactaaccgg accagagcag aagggaacag attacgtgaa 1200
gctggtaata ttaatcagtc actaatgacg ctaagaacat gtatggatgt cctaagagag 1260
aaccaaatgt atggaactaa caagatggtt ccatactgag attcaaagtt aacctctctg 1320
ttcaagaact actttgatgg ggaaggaaaa ttacgacaaa tctgtgtgtg gaacccaag 1380
gctgaagatt atgaagaaaa cttgcaagtc atgagatttg cggaagtgtc tcaagaagtt 1440
gaagtagcaa gacctgtaga caaggcaata tgtggtttaa cgctggggag gagatacaga 1500
aaccagcctc gaggtccagt tggaaatgaa ccatttggtt ctgacgtggg tttgcagagt 1560
tttccacctt tgcgctcatg cgaaattttg gatatcaacg atgagcagac acttccaagg 1620
ctgattgaag ccttagagaa acgacataac ttacgacaaa tgatgattga tgaagtttaa 1680
aaacaatcta atgtttttaa agctttgtta caagaatttg acaatgctgt ttaagtaaaa 1740
gaaaaccaca tgcaagggaa actaaatgaa aaggagaaga tgatctcagg acagaaattg 1800
gaaatagaac gactggaaaa gaaaaacaaa acttttagaataaagattga gatttttagag 1860
aaaacaacta ctatctatga ggaagataaa cgcaatttgc aacaggaact tgaaactcag 1920
aaccagaaac ttcagcgaca gttttctgac aaacgcagat tagaagccag gttgcaaggc 1980

```



atggtgacag	aaacgacaat	gaagtgggag	aaagaatgtg	agcgtagagt	ggcagccaaa	2040
cagctggaga	tgacagaataa	actctgggtt	aaagatgaaa	agctgaaaca	actgaaggct	2100
attgttactg	aacctaataac	tgagaagcca	gagagaccct	ctcgggagcg	agatcgagaa	2160
aaagttactc	aaagatctgt	ttctccatca	cctgtgcctt	tactctttca	acctgatcag	2220
aacgcaccac	caattcgtct	ccgacacaga	cgatcacgct	ctgcaggaga	cagatgggta	2280
gatcataagc	ccgcctctaa	catgcaaaact	gaaacagtca	tgacagccaca	tgtccctcat	2340
gccatcacag	tatctgtttg	aaatgaaaag	gcactagcta	agtgtgagaa	gtacatgctg	2400
acccaccagg	aactagcctc	cgatggggag	attgaaacta	aactaattaa	gggtgatatt	2460
tataaaacaa	ggggtgggtg	acaatctgtt	cagtttactg	atattgagac	tttaaagcaa	2520
gaatcaccaa	atggtagtgc	aaaacgaaga	tcttccacag	tagcacctgc	ccaaccagat	2580
ggtgcagagt	ctgaatggac	cgatgtagaa	acaaggtgtt	ctgtggctgt	ggagatgaga	2640
gcaggatccc	agctgggacc	tggatatcag	catcacgcac	aaccaagcg	caaaaagcca	2700
tgaactgaca	gtcccagtac	tgaaagaaca	ttttcatttg	tgtggatgat	ttctcgaaag	2760
ccatgccaga	agcagtcctt	caggtcatct	tgtagaactc	cagctttgtt	gaaaatcacg	2820
gacctcagct	acatcataca	ctgaccaga	gcaaagcttt	ccctatgggt	caaagacaa	2880
ctagtattca	acaaaccttg	tatagtgtat	gttttgccat	atttaatatt	aatagcagag	2940
gaagactcct	tttttcatca	ctgtatgaat	tttttataat	gtttttttta	aatatatttc	3000
atgtatactt	ataaaactaa	tcacacaagt	gtttgtctta	gatgattaag	gaagactata	3060
tctagatcat	gtctgatttt	ttattgtgac	ttctccagcc	ctggtctgaa	tttcttaagg	3120
ttttataaac	aaatgctgct	atattattagc	tgcaagaatg	cacttttagaa	ctatttgaca	3180
attcagactt	tcaaaataaa	gatgtaaagt	actggccaat	aataaccatt	ttaggaaggt	3240
gttttgaatt	ctgtatgtat	atattcactt	tctgacattt	agatatgcc	aaagaattaa	3300
aatcaaaagc	actaagaagt	acaaaaaaa	a			3331

&lt;210&gt; 64

&lt;211&gt; 3558

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 6437362CB1

&lt;400&gt; 64

ggcgcccgctc	ctcatcccgg	cgcttgagag	gacgcggggc	tgcgcaaatg	gcttgtcccc	60
ctctcggtct	ggaagctctt	cagccccctg	agccccagcc	gccccccgag	cccgcccttct	120
ccgaggcgca	gaagtggatt	gagcaagtaa	ctggcagaag	ttttgggtgat	aaagattttc	180
ggacagggtt	agaaaatgga	atcctcctct	cgagttgct	gaatgctata	aagccaggac	240
ttgttaaaaa	gatcaataga	ttgcctaccc	ccattgcagg	actggacaat	attatcttat	300
tcttgagagg	ttgtaaagag	ctcggcctta	aagaatctca	actttttgac	ccgagtgacc	360
tccaggatag	atccaacaga	gtaacagtca	agagccttga	ttatagtagg	aagctgaaaa	420
atgtattagt	taccatttac	tggctgggaa	aagcagcaaa	cagctgcaca	tcctacagcg	480
gaacgacact	aaacctgaag	gagtttgaag	gattgttggc	tcagatgcga	aaggacactg	540
atgacattga	aagtcctaaa	cgcagtatcc	gagacagtgg	ctacatcgac	tgctgggatt	600
ccgagcgag	cgactccctc	tctcctcctc	gccacggcag	agatgattcc	ttcgacagcc	660
tggattcctt	tggctctcgc	tctcggcaga	cgccttcacc	agatgtagtc	ctcaggggaa	720
gcacgcgatg	gagaggaagc	gactctgaat	ccgacttgcc	tcactcggaag	ctgcagatg	780
tgaagaagga	tgacatgtct	gcacggcgga	cttcccattg	tgagccgaaa	tcagcagtcg	840
cttttaacca	gtacctcccc	aacaaaagca	atcagacggc	ctacgtcccc	gcgcctctga	900
gaaagaagaa	agcagagaga	gaggaatacc	gcaagagctg	gagtaccgcc	acctccccgc	960
tgggtgggga	gaggcccttc	agatacggtc	cgagaactcc	tgtgtctgat	gacgcagaga	1020
gcaccagcat	gtttgacatg	cggtgtgagg	aggaggccgc	ggtgcagccg	cacagcaggg	1080
cccgccagga	gcagctgcag	ctgataaata	accagctgag	ggaagaggac	gacaaatggc	1140
aagatgacct	ggctcgtttg	aagagtcgta	gaagaagtgt	ttctcaggac	ttaatcaaga	1200
aagaggaaag	aaggaaaaaa	atggagaaat	tactggctgg	agaagatggg	acaagtgaac	1260
gaaggaaaag	catcaaaacc	tacagagaaa	ttgttcaaga	aaaagagcgg	agagagagag	1320
agctgcatga	agcatataag	aacgctcgg	cccaggagga	ggcagagggg	atccttcaac	1380
agtacattga	gaggttcacc	atcagtgagg	ctgttctcga	acgcttggag	atgccaaaaa	1440
ttctggaaaag	aagccattca	acagagccaa	atttatcctc	cttctgaat	gaccccaatc	1500
ccatgaaata	cctgcgga	cagtcactgc	ctccacccaa	attcactgcc	actgttgaaa	1560
ccaccattgc	tctgtccagt	gttctggata	ccagcatgtc	agcaggcagt	gggtctccaa	1620
gcacaaactgt	cactcccaaa	gcagtgccta	tgctgacacc	caagccttac	tcccagccca	1680
aaaattctca	agatgttctg	aagaccttta	aggtagacgg	gaaagtcagt	gtgaatggag	1740
agacgggttca	tagagaggag	gagaaggaaa	gagagtgtcc	cacgggtggc	cctgcccact	1800
ccttaaccaa	atcccagatg	tttgaagggt	tggccagagt	gcacgggtct	ccactggagc	1860
tgaacaaga	caacggtagc	atcgagatca	acataaagaa	gccaaactct	gttccccaa	1920
agctcgcagc	aaccactgag	aaaaacgga	cgaatagtca	agaggacaag	aatgatgggt	1980

gaaaatcaag	aaaaggggaat	atagaacttg	cctcatcaga	accacagcat	tttacaacaa	2040
ctgtgactcg	atgcagcccc	accgtggcct	ttgtggaatt	tccctccagc	ccccagctga	2100
agaatgatgt	gtcgggaagaa	aaagaccaga	agaaaccaga	aatgaaatg	agtggaaagg	2160
tggagttggt	gctgtcacaa	aaggtggtaa	agccaaaatc	tccagaacc	gaagcaacgc	2220
tgacatttcc	atctctggac	aaaatgcctg	aagccaacca	actacatttg	ccaaatctca	2280
attctcaagt	ggattctcca	agcagtgaga	agtcacctgt	tacgacacct	tttaagttct	2340
gggcatggga	cccagaagag	gagcgcaggc	gacaggaaaa	atggcaacag	gaacaggaac	2400
gtttgctcca	ggagagatac	cagaaggagc	aggacaagct	gaaagaagag	tgggaaaagg	2460
cccaaaagga	ggtggaagag	gaagaacgca	gatactatga	ggaggagcgt	aagataattg	2520
aagacactgt	ggttccattt	actgtttctt	caagtccgc	tgaccagctg	tctacctctt	2580
cctccatgac	tgaaggcagt	gggacaatga	ataagataga	cctgggaaac	tgtcaagatg	2640
aaaaacaaga	cagaagatgg	aagaaatcat	tccaggggaga	tgacagtgc	ttattgctga	2700
agactaggga	aagtgatcga	ctggaggaga	agggcagcct	aactgaaggg	gccttggctc	2760
attctgggaa	ccctgtatca	aaaggagtcc	atgaagacca	tcagctggat	accgaggctg	2820
gggccccaca	ctgtggaaca	aaccacagc	ttgctcagga	tccatccag	aatcagcaga	2880
catcaaatcc	aacgcacagt	tcagaagatg	tgaagccaaa	aacctcccg	ctggataaaa	2940
gcattaacca	tcagatcgag	tctcccagtg	aaaggcggaa	gtctataagt	ggaaagaagc	3000
tgtgctcttc	ctgtgggctt	cctttgggta	aaggagctgc	aatgatcatc	gagaccctca	3060
atctctatct	tcacatccag	tgtttcaggt	gtggaatttg	taaaggccag	cttgagatg	3120
cagtgagtgg	gacggatggt	aggattcgaa	atgggtctct	gaactgtaat	gattgctaca	3180
tgcatccag	aagtgcggg	cagcctacaa	cattgtgaca	cggctttcaa	gcttccggat	3240
cactcaccat	ttctttactg	agagtgtccc	ctggcaactg	cttaacaaaa	tcccaagctc	3300
aggggcttct	cagcatttac	ctaattcttg	aaaggctctt	ctgaaagggtg	gtatctgttc	3360
ttctgtagca	cagtgtttat	gtttttcctg	tttattgttt	tgggtttttt	tttttttttg	3420
catttgcaca	gtatacacia	aagaatatgg	ggttgtaatg	atcctgaata	gctcaaaaaa	3480
ggttttagca	tggtcaaaca	ggcttatggt	ttaaaatgtg	ttattctctt	ctttgggaat	3540
tagctaaatg	atgcaata					3558

&lt;210&gt; 65

&lt;211&gt; 5373

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 4173970CB1

&lt;400&gt; 65

cccgaactgtc	taatcattag	tctcaaaaat	tattgcttgg	tgaagctttt	ttttaaaaaa	60
aagcttcagg	aaacaaaagt	gcatttgggtg	gtatatgaag	ataacctaaa	acaggatatc	120
tgttttcattt	atgggcctttt	gtcttattct	ttgggtctgag	atgatttttt	tctgagttga	180
ctttttagggt	gccacgtgtg	ccaacgcata	cacttgccat	ggttgtacct	cccaggaac	240
ctgacagaa	ttcacaggag	aactctcctg	cccttttagg	agtgcaaaaa	gctgtgagta	300
ccagagtgcc	cactggttcc	aacagttctt	ctcagaccac	agagtgtctt	acacctgaat	360
cctgtttcgca	gactacaagc	aatgtggcct	cccaatcgat	gcctcctgtg	tatccttcag	420
ttgacattga	tgacatact	gagagcaatc	atgacacagc	attaacacta	gcttgtgcag	480
tgggtcatga	agaacttgta	tctgtgctca	ttgcacggga	tgccaaaatt	gaacacagag	540
acaaaaaagg	tttcacacca	ctaactcctg	cagcaacagc	agggcatggt	ggagtgtgtg	600
aaatcctttt	ggataaagg	ggagatatag	aagcacagtc	tgaacgaact	aaggatactc	660
cgcttttcatt	ggcatgttct	ggtggacgtc	aggaggtggt	agacttgctg	ctggctcgag	720
gtgcaaataa	agaacatagg	aacgtatctg	attatacacc	actgagtcta	gctgcgtctg	780
gaggatatgt	taatcatctt	aagattctgc	ttaattgctg	ggcagaaatt	aattcaagga	840
ctggggagtaa	actaggtatt	tctccccctga	tgttggctgc	aatgaatgga	catgttctctg	900
cagtaaaatt	gctgctcgat	atgggttcag	acattaatgc	ccaaatagag	accaatcgga	960
acacggctct	caccctggcc	tgtttccagg	gccgagcaga	agtagtgagt	ttgcttctgg	1020
accgaaaagc	caatgttgaa	cataggggcaa	agacgggtct	taccccttg	atggaagcag	1080
cttctggagg	gtatgcagag	gttggaagag	ttcttcttga	taaaggagca	gatgttaatg	1140
ctccccctgt	gccttctctca	agagatactg	ctttaacaat	agcagcagac	aaaggtcact	1200
acaaattttg	tgaactcctg	attcataggg	gagcccacat	tgatgttcgt	aacaaaaagg	1260
gaaatacgcc	actttggctg	gcataccaatg	gaggtcattt	tgatgttggtg	cagttgctag	1320
tgcaagcagg	tgctgatgtg	gatgcagcag	ataaccggaa	aatcacacct	cttatgtcag	1380
tctctcgcaa	ggctcatgta	aaagttgttc	aatatttggt	aaagggaagta	aatcagttcc	1440
cttctgatata	agaatgcatg	agatacatag	caacaattac	agataaggaa	ctgttgaaaa	1500
aatgtcatca	atgtgtcgaa	accattgtga	aggctaaaga	ccagcaagct	gcagaagcaa	1560
ataagaatgc	gagtattctt	ttaaaggaac	ttgatctgga	aaagtcaaga	gaagagagca	1620
gaaagcaggc	tcttgctgct	aaaagagaaa	aaagaaaaga	aaagagaaaa	aagaaaaaag	1680
aggaacagaa	aaggaaacag	gaagaagatg	aagaaaacaa	acctaaggag	aattcggaac	1740

taccagagga	tgaagatgaa	gaggagaaatg	atgaagatgt	ggagcaagaa	gttcccatag	1800
aacctcctag	tgcaaccacc	accactacga	ttggaatctc	tgcaacatct	gcaacattca	1860
caaatgtgtt	tgggaaaaaa	agggccaatg	tgggtgacaac	tcccagcacc	aatcggaaaa	1920
ataagaagaa	caaaacaaaa	gaaacccctc	ctacagcaca	tttaatttta	ccagaacaac	1980
atatgtcttt	agcccaacaa	aaggcgagata	aaaataaaaat	aaatggagaa	cctagagggtg	2040
gtgggtgcagg	tgggaatagt	gattcagata	acttggacag	cacagactgc	aacagtgaga	2100
gtagcagtg	tggtaaaaagc	caagagttaa	attttgtgat	ggatgtgaat	tcctctaaat	2160
acccctcact	gctccttcct	tcccaagaag	aaaagacaag	tactgtctact	tccaaaactc	2220
agacacgcta	caagacagtg	tcattggccat	taagctctcc	aaacataaag	ctgaatctca	2280
ctagccctaa	aaggggtcag	aaaagagaag	aaggggtgaa	agaagttgta	cgaagggtcaa	2340
agaaattgtc	tgttccagcc	tcagtgggtg	cgaggataat	gggaagagga	ggatgcaaca	2400
tcactgcaat	acaggatggt	actgggtgcc	atattgatgt	ggataaacia	aaagataaga	2460
atggcgagag	aatgatcaca	ataaggggtg	gcacagaatc	aacaagatat	gcagttcaac	2520
taatcaatgc	actcattcaa	gatcctgcta	aggaactgga	agacttgatt	cttaaaaaatc	2580
atatcagaac	acctgccagc	accaaatacaa	ttcatgctaa	cttctcatct	ggagttaggta	2640
ccacagcagc	ttccagtaaa	aatgcatttc	ctttgggtgc	tccaactctt	gtaacttcac	2700
aggcaacaac	gttatctacg	ttccagcccg	ctaataaact	taataagaat	gttccaacaa	2760
atgtacgttc	ttctttccca	gtttctctac	ccttagctta	tcctcaccct	cattttgccc	2820
tgtgtgctgc	tcaaaactatg	caacagattc	ggcatcctcg	cttaccocatg	ccccagtttg	2880
gaggaacctt	ctcaccttct	cctaacacat	ggggaccatt	cccagtgaag	cctgtgaatc	2940
ctggcaaacac	aaatagctct	ccaaagcata	ataacacaag	cogtctacct	aaccagaacg	3000
ggactgtttt	accctcagag	tctgctggac	tagctactgc	cagttgtcct	atcactgtct	3060
cttctgtagt	tgtgtccagt	cagcaactgt	gtgtcactaa	taccgggact	cttcatctag	3120
tcagaaagca	gttgtttgct	tgtgtgctta	agacaagtcc	tccagcaaca	gtgatttctt	3180
ctgtgacaag	cacttgtagt	tccttgctct	ctgtctctct	tgcacctatc	actagcgggc	3240
aagctcccac	cacattttct	cctgcaagta	cttctcaagc	acagctttct	tcacaaaaga	3300
tggagtcttt	ctctgctgtg	ccaccaccca	aagagaaagt	gtccacacag	gaccagcccc	3360
tggcaaacct	atgtacccca	tcttcaactg	caaacagttg	cagtagctct	gccagcaaca	3420
ccccgggagc	tccagaaact	cacccatcca	gtagtccac	tcctacttcc	agtaacacac	3480
aagaggaggg	acagccatcc	agtgtgtctg	atttaagtcc	tatgtcaatg	cctttttgcat	3540
ctaactcaga	acctgctcca	ttgactttga	catcaccag	aatgggtgct	gctgataatc	3600
aggacaccag	taattttacct	cagtttagctg	taccagcacc	tcgagtttct	catcgaatgc	3660
agcccagagg	ttcttttttac	tccatggtac	caaattgcaac	tattcaaccag	gatccccagt	3720
ctatttttgt	tacgaatcca	gttactttta	caccacctca	aggcccacca	gctgcagtg	3780
agctttcttc	agctgtgaac	attatgaatg	gttctcagat	gcacataaac	ccagcaataa	3840
agtctttgcc	acctacatct	ggcccagcca	cactttttca	tcacttcagc	agtctttttg	3900
atagtatgca	ggtgccagct	aaccagggct	ggggagatgg	tcactgtcc	tcacgagttg	3960
ctacagatgc	gttcttcact	gttcagtcag	cgttcctggg	taactcagtg	cttggacact	4020
tggaaaacat	gcacctgat	aaactcaaagg	cacctggctt	cagaccacct	ttcccagcgag	4080
tttctactag	tccagttggg	ttaccatcca	ttgacctatc	aggcagctcc	ccatcttctc	4140
cttctgctcc	tctggcaagt	ttttccggca	taccaggaac	aaggggtttc	ctgcaagggc	4200
cagctcctgt	tgggaactct	agtttcaaca	gacaacatct	ttctcccat	ccttggacaa	4260
gcgcctcaaa	ctcatgtgac	ttctctatct	catctgtttc	ttcgggatca	ttctcactc	4320
tttcagccac	ttctgcccc	ccaacgttgg	gccaaccaa	aggagttagt	gccagtcaag	4380
atcgaaagat	acctccccc	atttgaacag	agagactggc	ccgaattcgg	caaggagggt	4440
ctgttgacac	agccccggcg	gggaccagtt	ttgtcgctcc	cgttggacac	agtggaaatc	4500
ggtcaatttg	tgtaaatgct	gtgtcagaag	gcttatcagg	ttgggtcgaa	tctgtgatgg	4560
ggaaaccatcc	aatgcatcaa	caattatcag	acccaagcac	attctcccaa	catcagocaa	4620
tggagagaga	tgattctgga	atggtagccc	cctctaacat	ttttcatcag	cctatggcaa	4680
gtgggtttgt	ggatttttct	aaaggtctgc	caattttccat	gtatggaggc	accataatac	4740
cctctcatcc	tcagcttgct	gatgttccag	gagggcctct	gtttaatgga	cttcacaate	4800
cagatcctgc	ttgggaacct	atgataaaaag	ttatccaaa	ttcaactgaa	tgcactgatg	4860
cccagcagat	ttggcctggc	acgtgggcac	ctcatattgg	aaacatgcat	ctcaaatatg	4920
tcaactaagt	tagaaggtct	ttactcttta	gccttggtta	agaaacctat	gaccttggaa	4980
gaaccatggg	gatttttttt	taatgtgcct	aagaaatctt	ctctgagggt	ttagcaatgg	5040
aaatttgatt	gcccattgta	taagaacaaa	ttgatttctc	atccacctga	ttatgttctc	5100
tgggttagttt	agccattttg	aacttaagat	catatgaact	tagtgctttt	ggctaatacat	5160
acagaatact	acttgtatgc	agaagagaat	tagttgatta	catgtttcaa	ccttttaggg	5220
tgataaatac	atgtataatt	gtttacatac	ttaaaaggaa	aaagttgagt	aaattttctg	5280
tcataatagt	gctctacgta	atgtagcctg	tattaatgtg	aaatatattac	cagaatatct	5340
aataaaaaaga	tgaacagtct	ttaaaaaaa	aaa			5373

&lt;210&gt; 66

&lt;211&gt; 4333

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2772751CB1

&lt;400&gt; 66

```

gagaggggaca gaggctggag aaggatgtat ggctgcctt gggcttgtct gttccctcct 60
gagcctgagc cccttacctt cctgacccca tgaagcacac actggctctg ctggctcccc 120
tgctgggcct gggcctgggg ctggccctga gtcagctggc tgcaggggcc acagactgca 180
agttccttgg ccggcgagag cacctgacat tcacccagc agccagggcc cggtggtgg 240
ccctcgagt tcgtgcgcca ggactcctgg actccctcta tggcacctg cggcgcttc 300
tctcggtggg gcagctcaat cctttccctt cagagttggg aaaggcccta ctgaatgagc 360
tggtccctgg gacgtgagc gaggtgggtg ggtacgaggg gggctacgtg gtatgcgctg 420
tgatcgcggg cctctacctg ctgctgggtg ccactgccgg gctttgcttc tgctgctgcc 480
gctgcacacc ggcgtgcggg ggacgagtg agacagagca caaggcgctg gcctgtgagc 540
gcgcggccct catggtcttc ctgctgctga ccacctctt gctgctgatt ggtgtggtct 600
gtgcctttgt caccaaccag cgcacgcagt aacagatggg cccagcatc gaggccatgc 660
ctgagaccct gctcagcctc tggggcctgg tctctgatgt ccccaagag ctgcaggccg 720
tggtcacagca attctccctg cccaggagc aagtctcaga ggagctggat ggtgttgggtg 780
tgagcattgg gagecgatc cactctcagc tcaggagctc cgtgtacccc ctgtgtggcg 840
ccgtgggcag tttgggcccag gtcctgcagg tctccgtgca ccacctgcaa accttgaatg 900
ctacagtggg agagctgcag gccgggagc aggacctgga gccagccatc cgggaacacc 960
gggaccgcct ccttgagctg ctgcaggagg ccagggtgcca gggagattgt gcagggggcc 1020
tgagctgggg ccgcacctg gagctgggtg ctgacttcag ccagggtgcc tctgtggacc 1080
atgtcctgca ccagctaaaa ggtgtccccc aggccaaact ctocagcatg gtcaggagg 1140
agaacagcac cttcaacgcc cttccagccc tggctgccat gcagacatcc agcgtgggtg 1200
aagagctgaa gaaggcagtg gccagcagc cgggaagggt gaggacactg gctgaagggt 1260
tcccgggctt ggaggcagct tcccgcctgg cccaggcact gcaggagggt gagggagagca 1320
gccgccccta cctgcaggag gtgcagagat caggtggatc cagggtggatc gtgggtgcg 1380
tgctgtgctc cgtggtccta tctgtgtgct tctgcaacct gctgggacct aatctgggca 1440
tctggggcct gtctgccagg gacgacccca gccaccaga agccaagggc gaggtggag 1500
cccgttccct catggcaggt gtgggcctca gcttccctt tgctgcaccc ctcatcctcc 1560
tggtgttcgc caccttccct gtgggtggca acgtgcagac gctgggtgtg cagagctggg 1620
agaacagcga gctctttgag ttgacagaca cccagggaa cctgcccccg tccatgaacc 1680
tgctgcgaact tcttggcctg aggaagaaca tcagcatcca ccaagcctat cagcagtgca 1740
aggaaggggc agcgtctctg acagtcctgc agctcaacga ctctacgac ctggaggagc 1800
acctggatat caaccagtat accaacaagc tacggcagga gttgcagagc ctgaaagtag 1860
acacacagag cctggacctg ctgagctcag ccgccgcgcg ggacctggag gccctgcaga 1920
gcagtgggct ctagcgcatc cactaccccg acttccctgt tcagatccag aggccgtgg 1980
tgaagaccag catggagcag ctggcccagg agctgcaagg actggcccag gcccaagaca 2040
attctgtgct ggggcagcgg ctgcaggagg aggccaaagg actcagaaac cttaccagg 2100
agaaggtcgt ccccagcag agccttgtgg caaagctcaa cctcagcgtc agggccctgg 2160
agtccctgc cccgaatctc cagctggaga cctcagatgt cctagccaat gtcacctacc 2220
tgaaaggaga gctgcctgcc tgggcagcca ggatcctgag gaatgtgagt gagtgtttcc 2280
tggtccggga gatgggctac ttctcccagt acgtggcctg ggtgagagag gaggtgactc 2340
agcgcattgc cacctgccag cccctctccg gagccctgga caacagccgt gtgatcctgt 2400
gtgacatgat ggctgacccc tgggaatgcct tctggttctg cctggcatgg tgcaccttct 2460
tctgatccc cagcatcacc ttgcccgtca agacctccaa atacttccgt cctactccga 2520
aacgcctcag ctccaccagc tctgaggaga ctcagctctt ccacatcccc cgggttacct 2580
ccctgaagct gtagggcctt gtgggtgtag gtgacctga ggctgcctgt cctccccctt 2640
gatttagcct gggccacagg acttcggtag ctcttgcccc agagcccagg ctggcatcca 2700
ggcctggact gtccccagt ceggttacc tggtccccc atgggacttc tgtgcagctg 2760
ctttctgctc acgaccccca tcattcacgc ccttaccccc atgctggtgg catcctcaca 2880
cagagccagc aagtccctcc aggtgtcacc tggacctgg ggtgggacag aggcctcgtc 2940
ggaagagcct gttctccacc tgctggagcc cctgccaag cctccccctg ccaagcctcc 3000
caaccccact ccccttcccg tgtgtcttcc ccctgcccc catctggcct cccccctggc 3060
ccccacttcc ctcttatgce cttcctggcc ctttgccttc gagaccaccc tcccttagtc cctcttcac 3120
catacttcca ctgctacctt gctggcccca gagaccaccc tgcccaacca aaccactcag 3180
gtaacgccac taatcaggca ggggccacca tggcctaggt ctgggctggc tgcaggccct 3240
gcctcatggc ctctgagccc tccactgccc cagggccttg ggccctctgc agatctcatc 3300
caggatattt tgggtgccag tgggtgagc gaggcctgtc tgaaggccga gctccctgtc 3360
ctgcacccaa gttagaaatg ggggtaccag cctctgtagt ctctctgagt gctggctccc 3420
aagggaaggga cctgggacct gggccacagt gggggccttg ccttacctct tcagaaggaa 3480
gcattctcca cagccccac ccaactttct taggagtgat ctggtggcca gaacaggatt 3540
ttgcacggcc ctttttatcc tgcgcagtgt gcctagggtc atccccagcc catcctgtg 3600
tcagccctga gtgctggaca ctgcgttcca gaaatgagga agaggagaga gaagagatgg 3660
acagacctca gatccattaa agtgttctca cttccctgag acttggttct gggctcctaa 3720

```

aaccaggttt	cctaggtctgg	gaccctgtac	atagttggtg	tttaatgagt	gtttatggag	3780
aggagagttc	taaggtcacc	tctggctgca	ggcatccagg	gattattcca	gcaatctgca	3840
ggtagggagt	gggtcccagc	ctggggagcct	gctgtcagga	gcaggcagac	ctggactcac	3900
agcctggctg	tgatgcttgt	tcgctcagct	tctccattta	tgagatgggg	agaatagtca	3960
cagcctcctc	aaagggttgt	gaaaatcaaa	tgtgataatt	tgtggaaagc	ccttagcagt	4020
ggcctggcac	aaaacaaatg	ctcagtggat	ggaagctgcc	tattattatt	gtcgttgttg	4080
ttgtttgcca	tgactgctct	gggcccgggg	tagagctagc	atccgggcat	gtacgaggga	4140
agagggaggc	aggcctctat	tcaaaggcag	aaattccttt	aagattgtgg	tctgctgggt	4200
ttcagggagt	gtctgtgttg	tttgtttttg	tttgtttgtt	tgttttgaga	cagggtctcg	4260
ctctgtcacc	caggctggag	tgtagtgggtg	cagtcttggc	tcaactgcaac	ctccacctcc	4320
tgggctcagc	gat					4333

&lt;210&gt; 67

&lt;211&gt; 2213

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2793768CB1

&lt;400&gt; 67

cggacctccc	gcgcgccccg	cacccgaccg	gctcagccgc	cggcagcgta	acacgcccta	60
cgctcgcttg	ctcgccggcc	tcagggcagg	caggcgggcg	cgggagaccc	cgccggggcc	120
gagacttggg	gcgggcgacg	aggaccaggt	tacggcctcc	tcgccatgtc	ctcggcctgc	180
gacgcgggcg	accactaccc	cctgcacctc	ctagtctgga	aaaacgacta	ccggcagctc	240
gagaaggagc	tgcagggcca	gaatgtggag	gctgtggacc	cacgaggctcg	aacattattg	300
catcttgctg	tttccttggg	acatttggaa	tctgctcgag	tcttactccg	acataaagca	360
gatgtgacaa	aagaaaatcg	ccagggatgg	acagttttac	atgaggctgt	gagcactggc	420
gatcctgaga	tgggtgtacac	agttctccaa	catcgagact	accacaacac	atccatggcc	480
cttgagggag	ttcctgagct	gctccaaaaa	attctcagag	ctccggattt	ctatgtgcag	540
atgaaatggg	aattcaccag	ctgggtgccc	ttggtttcta	gaatatgccc	aatgatgtc	600
tgtcgcatct	ggaaaagtgg	tgccaaaactg	cgcgtcgata	tcacattgct	gggatttgaa	660
aacatgagct	ggataagagg	gaggcgtagt	tttataattta	agggagaaga	caactgggcg	720
gagttaaatg	aagtcaacca	tgatgacaaa	gtgggtcacca	ccgaacgctt	cgacctttcc	780
caagaaatgg	agcgcctcac	tctggacttg	atgaagccaa	aaagcagggga	agttgagcgg	840
cggctcacia	gcctgtcat	taacaccagc	ctcgatacta	aaaataattgc	ttttgaaaga	900
actaaaatccg	gattctgggg	ctggaggaca	gataaagcag	aagttgttaa	tggtttacgaa	960
gcaaaggttt	acacagtaaa	caatgtgaat	gtgatcacca	aaatacgcac	agaacatctg	1020
accgaggagg	aaaaaaagag	atataaagca	gacaggaacc	cgctggaatc	tttgctggga	1080
actgtggaac	accaattttg	tgcacaaggg	gacctcacca	cggaaatgtg	tactgcaaac	1140
aaccccacag	ccatcacgcc	tgatgagtac	ttcaatgaag	agtttgatct	gaaagacagg	1200
gacattggaa	ggccgaaaga	gctgacgatt	agaacacaga	agttttaaagc	aatgttgtgg	1260
atgtgtgaag	agttttccct	ctctctgggtg	gagcaggtca	ttcccatcat	tgacctaatg	1320
gctcgaaacga	gtgctcattt	tgcaagactg	agagatttca	tcaaattgga	attcccacct	1380
ggatttccctg	tcaaaataga	aattcccttg	tttcatgtct	taaatgcacg	gattacattt	1440
ggaaaatgtta	atggctgtag	cactgccgaa	gaatctgtat	ctcaaaatgt	ggaagggacc	1500
caggctgatt	cagcttccca	catcacaaac	tttgaggttg	atcaatctgt	gtttgaaatt	1560
cccgaatctt	actatgttca	agacaatggc	agaaatgtgc	atttgcaaga	tgaagattac	1620
gagataaatgc	agtttgccat	ccagcaaaag	ctgctggagt	ccagcaggag	ccaggaactt	1680
tcaggaccag	cttcgaatgg	agggatcagc	cagacaaaca	cctatgacgc	ccagtatgag	1740
agggccatcc	aggagagcct	cctcaccagg	acagaaggcc	tgtgccccag	cgccctgagc	1800
gagacaagcc	gttttgataa	tgacttgcag	ctagccatgg	agctctctgc	caaagagctg	1860
gaggaatggg	agctccggct	ccaggaggaa	gaggctgagc	tccagcaagt	cttacagctg	1920
tcactcactg	acaaatagac	ctttcagcct	gtgagcctct	gcacaaagca	gaggctgtgg	1980
gctgtcacag	atgctgtgtc	aaccagggcc	ctagggctaa	gggcctgcac	cttgctgtga	2040
tgcagcaggc	aacaactgcc	ccttctttat	gcagagggtgc	agaaccaggg	actcatgggc	2100
ccatccaggc	tgctccctgg	cgtggagaag	ggaccaggga	ttgcaggccc	catctccagg	2160
ctaaggggag	gagagcatca	tcactttcca	ttagctgtat	tggcttgacg	gtc	2213

&lt;210&gt; 68

&lt;211&gt; 1142

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 3035248CB1

&lt;400&gt; 68

```

gggctggcga tccggatccc gcagggcgcgc tggctgcgct gcccggtgt ctgtcgtcat 60
ggtggggccc tgggtgtatc tgggtggcggc agttttgtct atcggcctga tcctcttcct 120
gactcgcagc cggggtcggg cggcagcagc tgacggagaa ccactgcaca atgaggaaga 180
gagggcagga gcaggccagg taggccgctc tttgccccag gagtctgaag aacagagaac 240
tggaagcaga ccccggcgtc ggagggactt gggcagccgt ctacaggccc agcgtcgagc 300
ccagcgagtg gcctgggaag acggggatga gaatgtgggt caaactgtta ttcagccca 360
ggaggaagaa ggcattgaga agccagcaga agttcaccca acagggaaaa ttggagccaa 420
gaaactacgg aagctagagg aaaaacaggc tcgaaaggct cagcgagagg cagaggaggc 480
tgaacgtgaa gaacggaaac gcctagagtc ccaacgtgag gccgaatgga agaaggaaga 540
ggaacggcct cgcctgaagg aagaacagaa ggaggaggaa gagaggaagg ctcaggagga 600
gcaggcccgg cgggatcacg aggagtacct gaaactgaag gaggccttcg tggtagaaga 660
agaagggtgt agcgaaacca tgactgagga gcagtctcac agcttcctga cagaattcat 720
caattacatc aagaagtcca aggttgtgct tttggaagat ctggctttcc agatgggcct 780
aaggactcag gacgccataa accgcataca ggacctgctg acggagggga ctctaacagg 840
tgtgattgac gaccggggca agtttatcta cataaccca gaggaactgg ctgccgtggc 900
caatttcata cgacagcggg gccgggtgtc catcacagag cttgccagg ccagcaactc 960
cctcatctcc tggggccagg acctccctgc ccaggcttca gcctgactcc agtccttcct 1020
tgagtgtatc ctgtggccta catgtgtctt catccttccc taatgccgtc ttggggcagg 1080
gatggaatat gaccagaaag ttgtggattt aaaggcctgt gaatactgaa aaaaaaaaaa 1140
gg                                                    1142

```